



3 2044 106 421 084

F9120

W. G. FARLOW.

There is said to be
a 2d Edn. of this work
dated 1824. See List of
Aris. works in *Icones Selectae*
Vol. II. From this one might
infer that the two editions
were identical in contents.
Pritzel, however, gives the
number of pages as different
in the two editions. The only
copy of second edit. which
I have yet seen (NM. 1817)
is that in the library of the
Mass. Hart. Soc. In that
copy both volumes are
bound in one and Vol. I
has the date 1824 and
the contents of it are identical
with that of Vol. I, 1815 but
the plates I-IV are bound
with the V-VIII of Vol. II
at the end of the whole
volume. In the volume
of the Mass. Hart. Soc. →
the Vol. II is dated 1818

but in plates, etc. That some
are not colored are the
same as in the original
edition of Vol. II.

Apparently the Mass.

Hart. Soc. volume is
the Editio Nova of Vol. I
but the Vol. II is the
original and not
the Editio Nova. This
volume belonged to

Mr. C. J. Sprague who
presented it to John
L. Russell in 1873

and was bequeathed
by him to the Mass.

Hart. Soc.

Fig. was the genuine
Vol. II of the Editio Nova identical
with the original Vol. II
1873 as above. The Editio
Nova have only one volume

Digitized by the Internet Archive
in 2016



Indp

OBSERVATIONES
MYCOLOGICÆ.

Auctore

E. M. FRIES.

Cum Tab. IV æncis.

H A V N I Æ.

Sumptibus Gerhardi Bonnier.

MDCCCXV.

1252 N. ALLEN ST.

MICROLOGICAL

1252 N. ALLEN ST.

OBSERVATIONES MYCOLOGICÆ.

AMANITA. PERS.

1. *Amanita incarnata*, pileo campanulato villoso albo, lamellis carneis, stipite solido nudo albo.

A. incarnata. Pers. Syn. p. 248.

Agaricus incarnatus. Batsch. El. fung. p. 51.

Ag. bombycinus. Schæff. Fung. Bav. t. 98.

Fungus magnus esculentus e volva erumpens, pileolo villoso albo, lamellis carneis, pediculo cylindrico albo. Mich. Gen. t. 76. f. c.

Habitat ad truncos arborum in Italia, Germania, Suecia.

Descr. Fungus nondum explicatus, volva involucratus, ovatus totus albus. Volva dein

laxa suberecta 1-2 uncialis membranacea viscosa laciniata albida v. flavescent. *Stipes* 3-6 uncialis crassit. digiti minimi solidus teres adscendens curvatus albus glaber, sursum nudus attenuatus. *Pileus* carnosus junior globosus, mox campanulatus hemisphaericus demum convexus subumbonatus 3-7 uncial. squamuloso-villosus fericeus, rarius apice calvus, albus interdum nigro-punctatus. *Caro* alba. *Lamellae* juniores albidæ mox incarnatæ liberæ lato-lanceolatæ 2-4 natæ tandemque deliquescentes.

2. *Amanita pusilla*, minor, Pileo hemisphaerico subpiloso incarnato-albido, lamellis roseis, stipite fistuloso nudo albido.

A. pusilla. *Perf. Syn.* p. 249. *Obs. myc.*

2. p. 36. t. 4. f. 4.

Agaricus volvaceus. *Bull. Champ.* t. 30.

β. minor.

Ad vias, in hortis &c. Etiam in *Svecia* legi.

Obs. Descr. dedit *Peersoon* l. c. Hæc et antecedens species *Coprinis* affines; lamellas enim semper deliquescere et in hæc nonnunquam nigrescentes, observavi.

3. *Amanita fulva*, pileo planiusculo glabro striato volvaque fulvis, lamellis stipiteque fistuloso nudo pallidis.

A. spadicea. β . *fulva.* *Perf. Syn.* p. 248.
Albert. et Schweitn. Consp. Fung. Nisk. p. 141.

Agaricus fulvus. *Schæff. Fung.* t. 95.

Agaricus trilobus. *Bolt. Fung.* t. 38, f. 2.
Abbild. der Schwäm. t. 9.

Agaricus vaginatus. *Bull. Champ.* t. 512.
 f. F. G. H. L. N. O.

In filvis frequens. Etiam in paludibus turfosis.

Deser. *Stipes* fistulosus v. substantia spongiosa intus fartus 4–5 uncial. crass. pennæ anserinæ l. cygneæ fragilis æqualis l. fursum attenuatus cæpius albus, interdum flavesceus, lævis l. squamulosus. *Volva* vaginalis elongata lævis glabra fulvo-rubra. *Annulum*, qui in *A. livida* juniore adest, nunquam in hac vidi. *Pileus* carnosus 1–2 uncial. junior campanulatus mox explanatus, umbone parvo subobsoleto, fulvus nudus l. squamis majusculis albis vestitus, margine striatus. *Lamellæ* liberæ latiusculæ 2-natæ l. integræ candidæ carne pilei latiores.

Obs. *Aman. livida* et *spadicea* α P., omnino sunt varietates, inter quas nulli dantur limites. *Pileus* campanulatus et planiusculus ab ætate solum pendet. Ex hac specie, plurimis sub nominibus ab Auctoribus descripta, *Aman. fulva* etiam sine dubio originem ducit; sed ob magni-

tudinem minorem, stipitem nudum, pilei volvæque colorem constantem distinxit.

4. *Amanita nitida*, candida; pileo convexo nitido albo-maculato, lamellis subadnatis, stipite annulato basi bulboso.

In pinetis *Sveciæ*.

Descript. *Stipes* 2-3 unc. initio crassior deinde elongato-angustatus solidus candidus lævis glaber, sed ex *annulo* albo laxo reflexo sublacero ad apicem stipitis inserto fibrillosus nonnunquam, basi bulbosus volvatus. *Volva* albida mollis oblitterata submarginata glabra, diam. interdum ultra uncial. *Pileus* carnosus junior subcampulatus obtusus, deinde convexus, tandemque planus candidus nitens fere flavescens $1\frac{1}{2}$ - $2\frac{1}{4}$ unc. non striatus, squamulis majusculis irregularibus sordidis l. pallidis irregulariter sparsis exasperatus. *Caro* alba. *Lamellæ* albæ numerosæ tenuēs tenaces inæquales subadfixæ.

Obs. *A. virescenti* P. proxima; sed abunde distincta. Cum *A. bulbosa* P. et *umbrina* P. in Nov. Fl. Svec. p. 17 descriptis nullam habet affinitatem.

AGARICUS. LINN. PERS.

Fam. I. LEPLOTA.

150 *Agaricus caperatus*, subsolitarius, pileo carnosio convexo rugoso pallefcente-citrino, lamellis ochraceo-argillaceis, stipite solido supra annulum obliquum squamuloso.

Species non differt:

Agaricus caperatus. Pers. Obs. myc. I. p. 48 descr. Ejusdem Syn. p. 273.

In ficcis Specie cum *A. alutario* Pers. copiose.

Descriptio. Solitarius magnus. Stipes solidus l. a vermibus erosus 3-6 unc. subæqualis crassitie digiti intermedi albidus, supra annulum obliquum album subevanescentem evidenter albido-squamosus. Pileus carnosus convexus demum planiusculus 4 unc. et ultra glaber disco lævis, marginem versus lacunoso-fulcatus dilute flavus. Lamellæ versus stipitem rotundatæ leviter adfixæ 4-natæ ochraceo-argillacæ.

Obs. Fungus noster a *Persoonii* differt magnitudine, stipite subæquali et lamellis rotundato-adnexis.

6. *Agaricus olivaceo-albus*, subsolitarius, pileo carnosio convexulo subumbonato fusco-oli-

vaceo, lamellis candidis adnatis, stipite annuloque fugaci viscoso albidis.

Ag. brunneus. Schæff. F. t. 312. (var.)

In pinetis Sveciæ, Germaniæ.

Descr. *Stipes* solidus 2-5 unc. penna asperina crassior subæqualis interdum flexuosus firmus albidus l. fusco-maculato-reticulatus subviscosus. *Annulus* albus fugax viscosus cortinæ haud dissimilis. *Pileus* carnosus primum convexus demum planiusculus umbonatus $\frac{1}{3}$ -2 unc. viscidus fusco-olivaceus, rarius olivacco-brunneus, umbone prominente obscuriori, margine striatulo dilutiori. *Caro* albida. *Lamellæ* albæ totæ adnatæ interdum fere decurrentes firmæ 2-natæ carne pilei latiores.

Obs. *A. brunneus* P. longe alia est species. — Ad Cortinarias ob lamellas candidas immutabiles hæc species referri nequit.

Fam. II. CORTINARIA.

7. *Agaricus scaber*, gregarius cinereo-fuliginosus, pileo squamoso subcampanulato, lamellis obscurioribus, stipite fibrilloso dilutiori cylindrico.

A. scaber. Sowerb. Fung. t. 207.

In silvaticis Angliæ, Sveciæ, subæspitosis.

Descr. *Stripes* solidus subbipollicaris crass. digiti minimi sursum subangustatus cylindricus pallide cinerascenti-fordidus, squamis obscurioribus fibrillofis vestitus. *Pileus* campanulatus l. convexus 1-2 unc. carnosus dilute cinereo-fuligineus ficcus, squamulis minutis scabris obscurioribus exasperatus, margine deflexo regulari. *Caro* albida. *Lamellæ* cinero-fuligineæ adnatæ subalternæ carne pilei sublatores. *Cortina* fugax albida.

8. Ut *Novitiæ Floræ Sveciæ*; nec a *Cel. Swartz* in *Vet. Acad. Handl.* descriptæ, hoc loco nominandæ:

1. *A. nudus.* *Pers. Syn.* p. 278. *Bull. Herb.* t. 439. f. A.

In pinetis.

2. *A. violaceo-cinereus.* *Pers. Syn.* p. 279. *Schæff. Fung.* t. 3.

In filvis frondosis.

3. *A. argentatus.* *Pers. Syn.* p. 286.

In filvis muscolis acerosis.

4. *A. sanguineus.* *Pers. Syn.* p. 289. *Sowerb.* f. 43.

In pinetis. Descripti in *Nov. Fl. Svec.* p. 18.

5. *A. albobrunneus.* *Pers. Syn.* p. 293. *Schæff. Fung.* t. 38.

In filvis.

6. *A. fulgens.* Pers. Syn. p. 294. Schæff.
Fung. t. 24.

In betuletis meum legi fungum.

7. *A. subrepandus.* Pers. Syn. p. 300.
In fagetis.

Ceterum *A. Monachella.* Schæff. t. 66
aliique in parte secunda describendi.

Fam. III. G Y M N O P U S.

+ Pileo albedo.

9. *Agaricus candidus*, pileo carnosio
sericeo-fibroso albo-nitido, lamellis albedo-brun-
nescentibus subliberis, stipite solido brevi.

Agar. candidus. Batsch. El. fung. p. 133.
t. 20. f. 106.

In ericetosis sylvaticis circa Temsis in Suecia
vulgaris.

Descr. *Stipes* solidus uncialis crassiuscu-
lus glaber lævis l. subtriatus candidus, semper
subæqualis et rarissime deorsum subincrassatus.
Pileus carnosus convexus albedo-pallescent 1-2
uncialis sericeo-fibrosus, vix umbonatus, demum
rimoso-repandus. *Caro* alba. *Lamellæ* albidæ
mox fordidæ cinerascetes, sed simul parum in

ochraceum colorem vergentes semiovatae 2 - 4
natae liberae.

Obs. Quamvis fungus meus stipite sub-
aquali et pileo vix umbonato recedat, secundum
iconem tamen idem omnino fungus. — Affines
etiam sunt:

- 1) *Agar. decumbens* P., sed stipites decumbens
bulbosus, lamellae adnexae magis ochra-
cae &c.
- 2) *Ag. argillaceus* P., sed differt. stipite tenui-
ori 1 lin. crasso, pileo acute umbonato,
lamellis adnexis &c.
- 3) *Ag. tricholoma*, *Albert. et Schwein.*
stipite quoque tenuiori, lamellis decur-
rentibus &c.

10. *Agaricus albus*, totus albus, pileo
carnoso glabro convexo, lamellis totis adnatis,
stipite subsolido brevi.

Agar. albus. *Pers. Syn.* p. 363. *Albert.*
et Schwein. l. c. p. 181.

Ag. virgineus. *Batsch. El. fung.* p. 39.
t. 3. f. 12.

In fagetis &c. *Sueciae*, *Germaniae*.

Descr. Solitarius, totus albus. *Stipes*
brevis uncialis, raro subelongatus, $\frac{3}{4}$ unc. crassus
solidus, medio interdum cavus, glaber laevis sub-
irregularis nunc rectus nunc adscendens, nunc

furfum nunc deorfum subattenuatus. *Pileus* convexus carnosus 2-3 unc. demum subplaniusculus lævis glaber, margine deflexo l. involuto glabro, interdum repando. *Lamellæ* distinctæ latiusculæ 2-natæ interdum flexuofæ, "ferrulatæ *Perf.*" totæ adfixæ.

In pinetis sæpius minor.

II. *Agaricus inamoenus*, lacteus graveolens, pileo planiusculo subumbonato carnosio glabro, lamellis adnatis, stipite solido elongato tenaci.

In pinetis, præfertim abietinis *Sveciæ* frequens.

De scr. — Odor et forma *A. sulphurei*, cui perquam affinis et in cujus consortio crescit. Distinctam tamen esse speciem, vix est quod moneam. — Totus albus. *Stipes* 2-4 unc. solidus subæqualis l. fursum angustatus lævis glaber, basi pubescens tenax. *Pileus* junior convexus dein planiusculus sæpe umbonatus glaber lævis albus uncialis et ultra, margine patente. *Caro* alba. *Lamellæ* alternæ firmæ subadnatæ, carne pilei latiores. Odor ingratus.

Agar. graveolens. Schrad. Spic. p. 120. stipite fragillimo tantummodo differt.

12. *Agaricus erythropus. γ. rhodopus*, pileo subcarnoso convexo glabro lammellisque

latisimis liberis albis, stipite fistuloso glabro roseo.

Ag. erythropus. *Perf. Syn.* p. 367. var.

In fagetis umbrosis Octobri.

Descr. *Stipes* fistulosus $1\frac{1}{2}$ –3 unc. crassit. pennæ gallinæ teres glaber sursum attenuatus subfragilis roseus lævis, interne ad latera stipitis cavi subvillosus. *Pileus* subcarnosus convexus vertice demum planiusculus albus $\frac{3}{4}$ – $1\frac{1}{2}$ unc. latus glaber lævis l. margine demum substriatus. *Lamellæ* latisimæ candidæ liberæ distinctæ 2–4 natæ.

Sequenti, etiam in Svecia a me primo lecto, vix affinis. Hujus descriptionem etiam dabo.

13. *Agaricus confluens*, cæspitosus pileo subcarnoso convexo glabro lamellisque angustis liberis albidis, stipite fistuloso subcompresso rubro pulverulento-tomentoso.

Ag. confluens. *Perf. Obs. myc.* I. p. 8. *Perf. Syn.* p. 368. *Id. et descr. f.* p. 15. t. 5. f. 1. *Albert. et Schweitn.* l. c. p. 184. α . et β . rufo-cinnamomeus.

Ag. tremulus et dispar. *Batsch. El. fung.* p. 104, 210.

In fagetis et pinetis frequens. Etiam in Svecia.

Descr. Cæspitosus. *Stipes* 3–5 uncialis

crass. pennæ gallinæ majoris, nunc teres nunc compressus, in speciminibus teretibus fistulosus, in compressis solidus, superne sæpe parum incrassatus l. dilatatus rufus, ubique tomento pulverulento subcinerascienti adpresso vestitus, basi cum pluribus confluentibus. *Pileus* subcarnosus fere membranaceo-coriaceus tenax flexilis convexus l. campanulatus albus explanatus 1-2 unc. glaber lævis. — Sæpe totus rufescit. *Lamellæ* confertæ angustæ, sed carne pilei sublatiores liberæ albidæ polynatæ.

14. *Agaricus xanthopus*, pileo membranaceo-carnoso planiusculo glabro substriato, lamellisque liberis albidis, stipite fistuloso fulvo-flavo glabro.

In pinetis, Septembri.

Descr. Priori valde affinis, sed optime distinctus. *Stipes* fistulosus 3-5 unc. subæqualis teres l. sursum subattenuatus glaber crass. pennæ gallinæ et ultra erectus flavo-fulvus l. rufescenti-flavus unicolor glaber lævis. *Pileus* planiusculus tenax subcarnosus, margine membranaceus patens substriatus, albus l. pallidus 1-2 uncialis glaber. Ut antecedens species nonnunquam rufescit, hæc flavescit. *Lamellæ* albæ liberæ inæquales carne pilei latiores approximatae. — Etiam *Mycenis* affinis est.

Obs. Ad plurimas novas species sæpius attuli; cui proximæ sint e notis; expertus enim novi, quantam in fungis determinandis affert utilitatem.

15. *Agaricus fordarius*, albidus, pileo carnosio convexo glabro, lamellis longissime decurrentibus stipite solido.

Ag. fordarius. *Perf. Syn.* p. 370.

In silvaticis.

Descr. *Stipes* 1 - 1 $\frac{1}{4}$ unc. solidus crassit. digiti minimi, vix ultra, subæqualis l. deorsum incrassatus glaber lævis albidus compactus elasticus. *Pileus* carnosus compactus hemisphæricus unicolor fordide albus lævis 1 unc. et ultra, margine deflexo primum involuto tomentoso, demum glabro acuto. *Lamellæ* latiusculæ congestæ albidæ 4-natæ longissime decurrentes.

Affinis est Ag. mollis, *Bolt.* t. 40 et vix specie differt.

16. *Agaricus cretaceus*, cæspitosus albus, pileo carnosio planiusculo lamellis subconfertis adnexus, stipite longiusculo solido subtomentoso. *Perf. Syn.* p. 369.

β . *ambiguus*, pileo subexcentrico.

In fagetis ad ligna putridissima.

Descr. *Stipites* numerosissimi cæspitosi basi connati radicati, radice albida lignum intraure,

curvati adscendentes subflexuosi 1-2½ unciales, crass. 2 lin., quisque lævis glaber basi albo-villosus candidus flexilis nec fragilis solidus. *Pileus* carnosus difformis subexcentricus junior convexus demum planiusculus glaber lævis pallescenti-albus l. candidus 1-2 unc., margine deflexo l. patente. *Lamellæ* totæ adnatæ tenues confertæ angustæ candidæ, carne pilei alba tenaci sublatores.

Pileus tactus rarius rufescit. — Ad *A. lignatilem* etiam accedit.

17. *Agaricus repens*, pileo carnosio planiusculo cinerascenti-albo, lamellis basi truncatis latis albis, stipite albido compresso cavo, radiculis longissimis repentibus.

In fagetis, *Augusto*.

Stripes 2-3 uncial. cavus glaber lævis subæqualis l. basi subincrassatus albidus difformis, compressus sæpe curvatus longitudinaliter fissilis, deorsum albido-pubescentis radiculis simplicissimis pedalibus et ultra albis horizontaliter repentibus auctus. Variat teres et bifidus. *Pileus* 3-6 uncialis subcarnosus fere pellucidus glaber planus exalbido-cinerascens tenax margine membranaceo patente acuto. *Lamellæ* vix adnatæ, sed prope stipitem oblique truncatæ 4-8 natæ albæ latæ (carne pilei 8ties latiores).

Singularis species, *Ag. compresso* proximus;

sed differt *stipite* sæpe tereti, nunquam torto, cavo, *pileo* majori plano tenaci regulari dilutiori et radicibus repentibus &c.

18. *Agaricus discoxanthus*, subsolitarius, pileo carnosio planiusculo viscoso albido, disco flavescente, lamellis adnatis fuscescentibus latiusculis, stipite cavo squamuloso albo.

Descr. *Stipes* cavus 2 unc. crass. pennæ anserinæ æqualis albus rectus, totus albido-farinulentus l. obsolete squamulosus. *Pileus* planus orbicularis valde viscosus carnosulus $1\frac{1}{2}$ - 2 unc. albus vertice obsolete subumbonatus flavescent, margine patente subintegro nonnunquam striato. Variat, licet raro, pileo unicolori albo et 3 unc. *Lamellæ* albido-fuscescentes leviter adnatæ 4-natæ decurrentes.

Differunt sequentes species maxime affines:

1. *Ag. eburneus*, Pers. Swartz., gregarius, pileo subcarnosio convexiusculo viscoso unicolori albido margine lævi, lamellis decurrentibus candidis latiusculis, stipite solido sub pileo farinulento.
2. *Ag. discoides*, Pers., subæspitosus, pileo carnosio planiusculo viscoso albido, disco flavescente, lamellis decurrentibus albidis tenuibus, stipite subsolido squamuloso.

Non multum forte erraverit, qui has, ut subspecies, conjunxerit.

** Pileo cinerascenti.

19. *Agaricus umbrosus*, pileo carnosomembranaceo planiusculo rugoso-squamuloso nigrescenti-umbrino margine piloso, lamellis confertis latissimis liberis roseis, stipite solido albido.

A. umbrosus. *Perf. Syn.* p. 353. *Ic. et descr.* 1. p. 8. n. 9. t. 2. f. 5, 6.

In ligno putrido fagineo locis umbrosis. *Octobri.*

Descr. *Stipes* 2-3 unc. crassit. pennæ anserinæ cylindricus solidus albidus subsquamulosus. *Pileus* subcarnosus planiusculus rugosus squamulosus 2-3 unc. umbrinus, margine integro villoso. *Caro* tenuis superne pileo concolor. *Lamellæ* latissimæ liberæ 4-natæ roseæ, juniores dilutiores.

An ab *Ag. pluteo* P. vere distinctus? In hujus consortio sequentem legimus *A. plutei varietatem*:

Ag. pluteus. β. permutatus, minor, pileo glabro flavescente, lamellis flavescenti-roseis, stipite flavescente striato. — Pileum etiam vidi cinereum sed lamellas et stipitem flavescentes.

20. *Agaricus agathosmus*, pileo convexiusculo carnosopapilloso svaveolenti, cinereo, lamellis albis decurrentibus, stipite solido albo.

In silvis abietinis densis, una cum *Ag. inamoeno*, sed rarior. *Septembr. Octobr.*

Descr. *Stipes* solidus 2 uncialis crassit. pennæ anserinæ æqualis glaber basi albido pubescens lavis albus rectus, raro curvatus, elasticus, demum cavus. Variat sursum subattenuatus et squamuloso-pubescens. *Pileus* carnosus convexus demum planiusculus 1 - 1½ unc., interdum subviscosus, papillis minutissimis dilutionibus oblitus. *Lamellæ* albæ 2-natæ decurrentes distinctæ latiusculæ. *Odor* gratissimus.

Diu pro varietate *Agar. pustulati* P. habui, a quo caractere ægre distinguitur, sed odore gratissimo, colore et magnitudine satis differre videtur. Ceterum nullo modo *Ag. eburneo* affinis, ut de fungo hoc attulerunt observatores diligentissimi de *Albertini* et de *Schweiniz*. — Noster *Ag. fragranti* potius affinis et hinc summa differentia apparet!

21. *Agaricus tumidus*, pileo carnosio convexo-plano lævi cinerascenti, lamellis distinctis albis latis emarginato-adnexis, stipite longo tumido.

Ag. tumidus. *Perf. Syn.* p. 350.

In pinetis *Sveciæ*, *Germaniæ* rarior, *Septembri*.

Descr. *Stipes* ventricosus crassus difformis

subradicatus, radice deorsum subangustata præ-
morsa glabra, carnosus solidus glaber striatus
pallidus albus, semel subflavescentis visus, subcur-
vatus firmus 3-4-6 uncialis, 1 unc. crassus.
Pileus carnosus convexus mox planus, raro sub-
umbonatus cinereus, glaber lævis 3-4 uncialis,
sæpe difformis et demum margine rimosus. *Caro*
alba. *Lamellæ* $\frac{1}{2}$ unc. totæ distinctæ albæ emar-
ginatæ, denticulo stipiti adfixæ.

22. *Agaricus lacchnopus*, pileo carnosio re-
pando convexiusculo cinereo, lamellis candidis
angustissimis subvenosis, stipite villosio subcavo
cinerascens.

In fagetis umbrosissimis ad ligna putre-
scentia. *Initio Octobr.*

Distincta species, si qua; nulli alio mihi
noto affinis.

Gregarius l. cæspitosus carnosus firmus. *Sti-
pes* solidus l. cavus, intus medulla spongiosa cen-
tro fartus 1-1 $\frac{1}{2}$ unc. undique pube tenui obli-
tus curvatus adscendens crassit. fere digiti minimi
substriatus cinerascens. *Pileus* irregularis subho-
rizontalis convexiusculus, nunc integer nunc di-
midatus nunc centralis nunc excentricus, loba-
tus lobis rotundatis firmis minime flexibilibus
marginalibus, sed interdum usque ad stipitem
fissus, cinereus furfuraceo-pubescentis l. glaber

1 - 1 $\frac{1}{2}$ unc. margine revoluta repando. *Lamellæ* angustissimæ marginem pilei versus subvenosæ, prope stipitem incrassato-adfixæ candidæ inæquales sæpe dichotomæ.

23. *Agaricus lampropus*, pileo subcarnoso convexo cinereo-griseo fibrilloso, lamellis albidis denticulo-adnatis, stipite nitido coeruleo fistuloso.

Ag. columbarius. *Bull. Champ.* t. 408. f. 1.

In silvaticis frondosis.

Descr. *Stipes* fistulosus 1-2 unc. 2-3 lin. crassus subæqualis glaber lævis nitidus coeruleus l. cyanescens teres, demum nonnunquam compressus. *Pileus* uncialis et ultra tenue carnosus convexus, raro umbilicatus, cinereo-griseus fibrillosus. *Lamellæ* latæ distinctæ pallidæ fordide-albæ l. cinerascens denticulo adfixæ.

Ag. atro-cyaneo P., cyanipede &c. affinis, sed colore distinctus. Etiam satis diversus videtur *Ag. calumbarius*, *Sowerb.* t. 161.

24. *Agaricus polius*, cæspitosus, pileo subcarnoso convexo dilute cinerascens glabro, lamellis albidis adnatis, stipite solido albo.

In fagetis inter folia dejecta, æstate et autumno.

Descr. *Stipites* plurimi (40-50) basi connati cæspitosi solidi 2-3 unc. crassit. vix

pennæ anserinæ æquales flexuosi glabri læves albi. *Pileus* primo convexus mox planiusculus subcarnosus glaber orbicularis dilute cinerascens $\frac{1}{2}$ - 2 unc., raro subumbonatus. *Lamellæ* 4-natæ albidæ adfixæ. Odor et sapor nullus.

Ag. canus, *Schum. Sall.*, hujus esset loci, nisi *pileus* sericeus.

25. *Agaricus lanuginosus*, pileo carnoso convexo squamuloso-villoso griseo-murino, lamellis adnatis cinerascens, stipite solido obscure cinereo.

Ag. lanuginosus. *Bull. Ch. t.* 370.

Juxta vias in silvis *Svecicis* meum fungum haud raro legi.

Descr. *Stipes* solidus brevis (subuncialis) æqualis rectus teres firmulus plus minusve obscure cinereus lævis l. tenuissime fibrilloso-squamulosus. *Pileus* carnosus convexus griseo-murinus 1 - 1 $\frac{1}{2}$ unc. villosus-squamulosus. *Caro* diluta. *Lamellæ* 2-4 natæ carne pilei latiores cinerascens totæ adnatæ.

26. *Agaricus mammosus*, pileo subcarnoso convexo umbonato fuscescenti-griseo glabro, lamellis levissime adnatis griseis, stipite fistuloso concolori.

Ag. mammosus. *Linn. Svec.* 2. n. 1218. (quoad descr.)

In silvis frequens.

Descr. *Stipes* fistulosus elongatus 2-4 unc. teres subfragilis striatulus glaber subæqualis pileo concolor. *Pileus* primo conicus dein convexus umbonatus glaber lævis fuscescenti-griseus l. subfuscus vertice carnosus totus explanatus $\frac{1}{2}$ - $\frac{1}{3}$ unc. *Lamellæ* 2-4 natæ griseæ distinctæ subliberæ l. levissime adnatæ, margine rarius crenatæ.

Obs. Agarici cinerei numerosissimi et hinc caute determinandi. — Ex sexcentis aliis, tales nonnullos modo attuli, qui rariores sunt et ab aliis proprietate quadam singulari distinguuntur. Ut Novitiæ Fl. Svecicæ nominandi *Agaricus nebularis*, *obscurus* &c. ex hac sectione.

*** Pileo pallescenti-flavescente, virescente et cinereo-flavo.

27. *Agaricus lepideus*, pileo carnosocompacto convexo subsquamoso flavescente, lamellis albis ferratis crenatisque decurrentibus, stipite solido squamoso.

Ag. squamosus. Schæff. Fung. t. 29, 30. optime. Dickf. Crypt. Brit. 2. p. 24. Hudf. Angl. 2. p. 614.

In truncis pineis Sveciæ, Germaniæ, Angliæ, solitarie.

Obf. Cum *A. squamoso* *Perf. Syn.* p. 409 non confundendus.

Descr. *Stipes* durus sublignosus solidus uncialis et ultra crass. digiti minimi æqualis radicatus squamis fuscis erectis vestitus albido-fuscescens. *Pileus* compactus carnosus convexo-pulvinatus durus crassus fordide flavesceus sub-squamosus squamis obscurioribus adpressis, margine deflexo submembranaceo. *Lamellæ* latiusculæ 4-natæ albe longissime decurrentes margine serratæ. *Pileus* $1\frac{1}{2}$ - 2 unc. latus.

Ad *A. tigrinum*, Bull. t. 70 *Cel. Pearson*, dubie licet, retulit et cum hoc miro modo *stipite* duro squamoso, *pileo* squamoso, *lamellis* decurrentibus ferratis albis, magnitudine, loco natali &c. convenit, sed *pileus* in *lepideo* constanter convexus flavesceus crassior, squamæ minus pilosæ, *stipes* firmior &c.

28. *Agaricus corticatus*, *pileo* carnosio convexo lævi firmo pallescente, *lamellis* distinctis liberis *stipite*que solido, cortice fuscescente sub-villoso-rugoso oblecto, albis.

In graminosis, *Augusto*.

Descr. Odor subfortis gratus. *Stipes* 2-3 unc., penna anserina tenuior, solidus, substantia interna fibrosa strictus firmus glaber lævis; sed cortice crassiusculo dissolubili albido fusce-

sciente deorsum præcipue villoso-rugoso vestitus. *Pileus* carnosus convexus, demum planiusculus glaber læviusculus 1-2 unc. subhumidus flavescenti-pallidus, raro subumbonatus, margine striato. *Caro* alba. *Lamellæ* distinctæ 2-natæ firmæ latæ liberæ, prope stipitem truncato-dilatatæ, albæ.

Notabilis sane species, odore, stipite corticato sfictisfimo, lamellis latis liberis et nonnunquam obbifidis.

29. *Agaricus penetrans*, pileo carnosio convexo-plano glabro flavescenti, lamellis adnatis albidis crebris, stipite folido albo striato, radice albo-villofa.

Truncos putridos pineos in *Suecia* penetrat.

Defcr. *Stipes* 2-3 unc. 4-5 lin. crassus æqualis folidus glaber albus striatus radicans, radice elongata æquali albo-villofa. *Pileus* subcarnosus convexus l. subplanus tenax fictus rugulosus glaber 1-2 unc. pallide flavus margine demum fissus. *Lamellæ* adfixæ pallidæ albæ l. flavescentes inæquales confertæ.

Varietatem *A. peronati* Sow. esse suspicatus sum, donec hunc invenerim.

A. fusipede ceterum proximus, quamvis ab hoc etiam abunde differt.

30. *Agaricus peronatus*, pileo carnosio-

membranaceo convexo glabro flavescenti, lamellis liberis subochraceis distinctis, stipite solido albidulo, deorsum flavescenti-strigoso radicato.

α) *A. peronatus*. *Perf. Syn.* p. 331. *Bolt. F.* t. 58. *Sowerb. F.* t. 37.

β) *approximatus*, pileo ochraceo, lamellis liberis subdistantibus stipiti approximatis, stipite albo-strigoso.

γ) *remotus*, pileo flavescenti, lamellis liberis a stipite remotis, stipite flavo- l. albo-strigoso.

δ) *apricarius*, lamellis stipitem subattingentibus, stipite pallide strigoso.

In Germania, Anglia, Svecia, frequens.

α, β. Inter folia dejecta faginea præcipue legi, locis subhumidis γ. Eisdem locis siccioribus. δ. Semel visus in ericeto aprico ad folia betulina.

Descr. Conveniunt omnes allatae, hujus sine dubio loci, varietates:

Stipite solido 2-3 unc. crassit. pennæ gallinæ l. anserinæ firmo subelastico laevi l. striatulo pallescenti-albido, rarius sordide ochraceo sursum recto subglabro basi subcurvato radicato subincrassato tomentoso, tomento erecto strigoso elongato conferro pallido. *Pileo* carnosio membranaceo fere coriaceo flaccido subumbonato; juniore convexo demum explanato glabro sæpius rugu-

loso flavescente l. pallide ochraceo 1-2 unc., margine primum inflexo, dein patente. *Carne* tenaci tenui alba. *Lamellis* liberis, sed ad stipitem plus minusve approximatis, 4-8 natis carne pilei latioribus firmis pileo subconcoloribus.

α. Distinguitur: lamellis stipitem attingentibus et tomento stipitis flavo.

β. Lamellis pileo subobscurioribus ochraceis, stipite interdum substipuloso et in solidis demum compresso, utrinque dilatato, basi albo-villoso.

γ. Lamellis a stipite remotis albido-ochraceis, stipite superne tomento adpresso vestito, inferne tomentoso; tomento nunc pallide flavo, nunc albido.

δ. Lamellis stipitem attingentibus, stipite albo-strigoso.

31. *Agaricus flavovirens*, pileo carnoso subdepresso flavescente, squamis minutis nigrescentibus, lamellis stipiteque substipuloso concolore.

Ag. flavovirens. *Perf. Syn.* p. 319. *Abb. der Schw.* 3. t. 24.

A. aureus. *Schaff. Fung.* t. 41.

In ligno putrido pineo Germaniæ, *Suecia* &c.

Descr. *Stipes* 2-3 unc. pennæ anserinæ crassit. subcentralis glaber lævis l. squamulis ci-

nerascentibus vestitus, teres subfistulosus citrinus subcurvatus deorsum attenuatus. *Pileus* carnosus subdifformis flavus 3 unc. primo convexiusculus demum et depressus, in quo statu cum *A. trichæo*, *bello* &c. non confundendus, squamulis cinereo-nigris obfuso, marginem deflexum l. patentem versus rarioribus. *Lamellæ* citrinæ 4-8 natæ adfixæ demum subliberæ.

32. *Agaricus luridus*. *Perf. Syn.* p. 321. *Schæff.* t. 69.

Ad hunc, et secundum iconem *Schæfferi* et descriptionem *Persoonii* sequens fungus referri debet:

♂. *spodoxanthus*, lamellis subtotis adnatis, stipite cavo.

In ligno fagineo putrido loco umbroso, semel in Svecia lectus, sed copiose.

Descr. Truncigenus gregarius l. cæspitosus: *Stipes* subcavus $1\frac{1}{2}$ - 2 unc. crassit. pennæ gallinæ subæqualis subtenax teres flavescens, superne subsquamofus farinulentus, interdum subcompresus. *Pileus* convexus carnosus demum planiusculus $\frac{1}{3}$ - $\frac{2}{3}$ unc. lævis virescenti-cinereo-flavescens, margine lævi subpatente. *Lamellæ* 4-8 natæ cinereo-flavescentes nonnunquam virescentes confertæ totæ adnexæ.

Agaricus cerineus, Floræ Sveciæ Novitia, et

nonnullæ aliæ novæ species huic proximæ, omnino etiam differunt.

**** Pileo rubro, rufo &c.

33. *Agaricus pratenfis*, pileo carnosio convexo glabro rubescenti, lamellis subobscurioribus distantibus decurrentibus, stipite solido deorsum attenuato.

A. pratenfis. *Perf. Syn.* p. 304. *Comm. in Schæff. Fung.* p. 124.

A. fulvosus. *Bolt. F.* 2. f. 56.

A. miniatus. *Sæwerb. F.* 1. 141.

In muscosis *Sveciæ*, *Germaniæ*, *Galliæ*, *Angliæ* &c.

Deser. *Stipes* 1 - 1½ uncialis basi attenuatus crassit. pennæ gallinæ tranferinæ solidus firmus teres glaber lævis. *Pileus* varius, campanulato-dilatatus subconvexus l. planiusculus subumbonatus glaber lævis nitidus 1 - 1½ unc. varie fulvo-rubescens. *Lamellæ* subcinnamomeo-rubræ (obscuriores pileo) decurrentes firmæ distantes carue pilei pallida vertice angustiores, margine latiores.

34. *Agaricus imbricatus*, pileo carnosio convexiusculo subumbonato, squamulis villosis imbricatis, spadiceo-rufo, lamellis albidis adnatis, stipite subsquamuloso.

In filvis. Octobri.

Descr. *Stipes* solidus l. sæpius cævus firmus rectus teres $2-2\frac{1}{2}$ unc. longus $\frac{1}{2}$ unc. crassus subsquamosus l. glaber fordidus albidus l. rufescens. *Pileus* carnosus convexus demum magis explanatus subumbonatus umbrino-rufus (fere *Hydn. imbric.* sed color magis rufus) squamulis villosis subimbricatis minoribus dense obsitus 2-3 unc. subæqualis firmus, carne alba. *Lamellæ* albidæ 4-natæ firmæ adnatæ tenuissime emarginatæ.

35. *Agaricus picreus*, cæspitosus, pileo carnosio convexo cinnamomeo glabro, lamellis flavis emarginato-adnatis, stipite glabro fulvo-umbrino.

Ag. picreus. *Perf. Syn.* p. 315. *Ic. et descr. Fung.* p. 14. t. 4. f. 7.

In truncis, præsertim pineis, Germaniæ, Sueciæ.

Descr. Gregarius cæspitosus. *Stipes* 2 unc. crassit. pennæ gallinæ l. parum crassior fistulosus fulvo-umbrinus subæqualis cylindricus glaber lævis adscendens. *Pileus* subcarnosus convexus glaber cinnamomeus lævis 1-2 uncial. ficcus l. interdum humidus margine integro. *Lamellæ* flavæ emarginatæ, adfixæ tamen leviter, subangustæ, sed carne pilei latiores, approximata 4-natæ. — Sapor nunc amarus nunc etiam nullus.

IV. MYCENA.

A) Graveolentes.

Obs. Familia hæc, a *Michaelio* constituta, maxime naturalis et species affines conjungit; hinc a me restituta.

36. *Agaricus scorodoni* (*Schæfferi*) graveolens, pileo subcarnoso planiusculo lamellisque crispis albidis, stipite brevi rufescenti glabro.

A. *Schæfferi*. *Perf. Obs. myc.* 2. p. 55.

A. *alliatus*. *Perf. Syn.* p. 373. *Swariz in Vet. Ac. Handl.* 1808. p. 91. *Schæff. F.* t. 99.

A. *alliaceus*. *Sw. Bot.* n. 179.

Fungus non vescus exilis albicans pediculo spadiceo, odore allii. D. *Breynii*. *Mich. Gen.* t. 77. f. 2.

In ericetosis frequens.

Obs. Descr. dederunt *Swariz* l. c. *Perfoon* l. c.

37. *Agaricus cepaceus* (*Michelii*) graveolens, pileo submembranaceo convexo luteo, lamellis albis, stipite mediocri lutescenti glabro.

Fungus parvus luteus odore Allii, lamellis albis, pediculo superna parte concolore. *Mich. Gen.* l. c. t. 98. f. 5.

Habitat in Italia in fimo per agros fuso-
autumno. *Michelius*.

Obs. Non vidi, sed autoritate *Michellii* attuli, cum quoad descriptionem, iconem et locum natalem omnino a sequenti differre videtur. *Stipes* deorsum incrassatus 2-3 unc. *Pileus* $\frac{1}{3}$ - $1\frac{1}{2}$ unc. (*Quoad Ic.*)

38. *Agaricus porreus* (*Scopoli*), graveolens, pileo submembranaceo convexo lamellisque fusco-flavescentibus, stipite longiusculo rufescenti basi piloso.

Ag. porreus. *Perf. Syn.* p. 376. *Swariz.* l. c. p. 92.

A. Scopoli. *Perf. Obs. myc.* 2. p. 55.

A. alliaceus. *Scop. carn.* 2. p. 454. *Sow. Fung.* t. 81.

In nemoribus Carnioliae, Angliae, Daniae, Sueciae inter folia dijecta.

Obs. *A. allicinus* (*Bulliardii*). *Schum. Sall. Bull.* t. 158, 524. f. 1, a me non visus; nec ab antecedenti et sequenti sufficientes invenire potui characteres.

39. *Agaricus alliaceus* (*Jacquini*) graveolens, pileo membranaceo convexo fuscescenti, lamellis albidis, stipite elongato nigrescenti-glaucobasi strigoso.

A. alliaceus. *Perf. Syn.* p. 375. *Obs. myc.* 2. p. 54. *Jacq. Obs. Bor.* p. 100. *Fl. Austr.*

t. 82. *Fl. Dan.* t. 1251. *Schum. Sall.* 2. p. 270.
Albert. et Schwein. l. c. p. 190.

Fungus alpinus odore et sapore Allii &c.
Mich. Gen. p. 144. t. 78. f. 4.

In fagetis umbrosis subudis. Etiam in ligno putrido.

Hæc, ut etiam omnes sequentes *Mycenæ* species, a me in *Svecia* primo lecta. — Hujus descriptionem dare superfluum foret, cum variis l. c. exstat; sed nomen specif. emendavi &c.

40. *Agaricus leptocephalus* (*Perfoonii*) graveolens cinereus, pileo membranaceo umbonato fulcato repando, lamellis crassiusculis adnatis, stipite brevi glabro.

A. leptocephalus. *Perf. Syn.* p. 381. *lc. et descr.* t. 12. f. 4.

Ad truncos in umbrosis *Sveciæ*, *Germaniæ*.

Descr. Stipes fistulosus uncialis l. parum ultra, crass. vix pennæ gallinæ æqualis glaber lævis cinereus. *Pileus* membranaceus umbonatus, umbone obscuriori cinereus opacus glaber striatus, striis profundis obscurioribus, margine subrepando; totus explan. $\frac{2}{3}$ unc. &c. *Lamellæ* obscure glauco-cinereæ crassæ distinctæ, in meis speciminibus totæ fere adnatæ. Odor fortis nitrosus.

Obs. *A. epipterygium* Pers. semper graveolentem inveni (tacent tamen omnes fere Auctores) et hinc ad hanc familiam pertinet.

B) I n o d o r i .

† Stipite lactescente.

(Etiam hæc Sectio naturalis.)

41. *Agaricus crocatus*, subsolitarius, pileo membranaceo cinereo obselete striato, lamellis albis stipiteque succo croceo refertis.

Ag. crocatus. Pers. Syn. p. 380. Schrad. Spic. p. 127. Schum. Sall. 2. p. 272.

β. *truncigenus*, pileo minori cinereo-rubro.

Inter folia dijecta faginea locis umbrosis humidis in Germania, Dania, *Svecia australi*.

β. Ad truncos fagineos.

Descr. *Stipes* 3-5 unc. crasfit. culmi gramin. rectus strictus l. flexuosus croceo-ruber lævis glaber, *radice* obliqua repente fibrilloso-villosa. *Pileus* campanulatus membranaceus papillatus glaber dilute cinereus striatus, vertice lævi subcarnoso obscure rubicundo. Diamet. infer. 4-5 lin., pilei explanati uncialis. *Lamellæ* distinctæ 2-natæ candidæ basi attenuato adfixæ et, stipitis instar, succo croceo refertæ.

Var. β . colore, magnitudine et loco natali tantum differt.

42. *Agaricus hamatopus*, cæspitosus, pileo carnosio-membranaceo rufescenti striato, lamellis albis stipiteque succo atro-sanguineo refertis.

Ag. hamatopus. *Perf. Syn.* p. 379. *Obs. myc.* 2. p. 56. *Albert. et Schwein.* p. 192.

Ad truncos putridos, præsertim fagineos, Germaniæ, Sveciæ.

Descr. Cæspitosus truncigenus totus fere lacte sanguineo-purpureo refertus. *Stipites* numerosissimi confertim nascentes fistulosi 2-4 unc. crassit. pennæ gallinæ adscendentes glabri deorsum pubescentes, rupti lactescentes. *Pileus* submembranaceus, disco pseudo-umbonato subcaroso, campanulatus demum convexus uncialis glaber striatus pallido-roseus l. exalbido-rufescens. *Lamellæ* leviter adfixæ, denticulo decurrente, 2-natæ albidæ.

43. *Agaricus sanguinolentus*, subsolitarius, pileo membranaceo rubello striato, lamellis concoloribus atro-purpureo-marginatis, stipite succum dilute rubrum stillante.

Am. sanguinolenta. *Albert. et Schwein.* l. c. p. 196.

In Germania, Svecia. Meum fungum in paludosis sphagnosis foliis abietinis adhærentem legi.

Descr. Terrestris solitarius l. gregarius. *Stipes* fistulosus elongatus 4-5 unc. filiformis strictus glaber tenax laevis, basi dilutior subfibrillosus. *Pileus* vere membranaceus late campanulatus obtusus dilute roseus, disco subobscuriori exsuccus glaber striatus, striis saepe obscurioribus 2-3 lin. altus, sed explan. $\frac{1}{4}$ - $\frac{1}{2}$ unc. *Lamellae* distinctae dilutiores 2-natae subliberae, margine atropurpureo, interdum concolori. *Succus* stipitis dilute roseus.

44. *Agaricus galopus*, subsolitarius, pileo membranaceo nigrescenti-glaucostriato, lamellis albis, stipite succum album secernente.

A. galopus. Pers. Syn. p. 379. Obs. myc. 2. p. 56. *Alberi. et Schwein.* p. 192.

A. lactescens. Schrad. spic. p. 128. n. 35.

In silvis muscosis praesertim humentibus Germaniae, Sueciae. Ad ligna putrida, raro.

Descr. Solitarius saepius epigeus, sed etiam epixylus. *Stipes* fistulosus 1-3 unc. gracilis basi subincrassatus, radice pubescente auctus, adscendens glaber laevis cinereo-nigricans sursum dilutior. *Pileus* membranaceus campanulatus 3-4 lin. striatus glaber "pruinatus Pers." colore plus minusve nigrescenti-cinereo l. glaucescente. *Lamellae* liberae albae 2-3 natae marginem versus latiores interdum cum pileo, ut stipites semper, lactescentes.

Obf. Lamellas nunquam arcuato-adnexas, ut apud *Perfoonium*; sed semper liberas cum *Schradero* inveni.

++ Stipite exsucco.

* Lamellis albis, pileo discolori.

45. *Agaricus lævigatus*, subgregarius, pileo membranaceo campanulato lævigato dealbato-cinereo, lamellis albis, stipite basi fibrilloso.

Ag. lævigatus. *Perf. Syn.* p. 380. *Albert. et Schwein.* p. 192.

In pinetis muscosis humentibus passim; etiam in *Suecia*.

Descr. Gregarius terrestris. *Stipes* fistulosus elongatus 2-4 unc. crassit. $1\frac{1}{2}$ - 2 lin. rectus subtenax albidus l. cinerascens albus lævis glaber basi piloso-fibrillosus. *Pileus* membranaceus campanulatus 4-6 lin. latus obtusus nec papillatus, omnino fere lævis exalbido-cinerascens, vertice subobscuriori. *Lamellæ* albidæ latiusculæ 2-natæ adfixæ.

46. *Agaricus pilosus*, solitarius, pileo membranaceo subhemisphærico papillato striato, lamellis stipiteque basi villoso-albidis.

Ag. pilosus. *Perf. Syn.* p. 380. *Albert. et Schwein.* l. c. p. 193. *Börsch. El. fung.* p. 67. t. 1. f. 2. non differt.

Ag. filopes. *Bull. Champ.* t. 320.

A. membranaceus. *Hoffm. Nomencl.* t. 6. f. 1.

In silvaticis intra folia dijecta faginea &c. —
Etiam in *Suecia* non infrequens.

Descr. Subsolitarius terrestris. *Stipes* fistulosus elongatus 2-4 unc. gracilis strictus subæqualis albido-cinereus lævis glaber, basi villo albido radicans. *Pileus* membranaceus, junior campanulatus mox hemisphaericus umbonatus striatus glaber albido-cinereus semiuncialis et infra. *Lamellæ* albæ distinctæ alternæ, breviores adfixæ.

47. *Agaricus conigenus*, gregarius, pileo subcarnoso difformi planiusculo umbonato lævigato, lamellis albis confertis, stipite brevi radiculis albis.

Fungus parvus in conis abietis dejectis nascens. *Buxb. Cent.* 1. p. 36. t. LVII. f. 2.

α) *fuligineo-lividus*. Ag. conigen. *Perf. Syn.* p. 388.

β) *pallens*. *Albert. et Schwein.* l. c. p. 195.

In Strobilis *Pini Abietis* et *sylvestris* in *Suecia*, *Germania* &c.

Obs. Fungus *Svecicus*, quem describo, ad β pertinet. *Buxbaumius* utrumque describit. Icon Ejus, ab omnibus neglecta, bona.

Descr. *Stipes* fistulosus $\frac{1}{2}$ - $1\frac{1}{2}$ unc. crassit. vix pennæ gallinæ æqualis albus, pulvere conca-

lore conspersus, basi fibrillis longis rectis albis squamulas strobilorum adhæret. *Pileus* subcarnosus tennis convexus subumbonatus demum planiusculus subdifformis angulatus l. depressus lavis l. margine tenuissime striatus $\frac{1}{2}$ unc. pallide cinerascens-rufescens (var. β) et cinereo-lividus P. (var. α .) *Perf.* *Lamellæ* confertæ candidæ angustæ liberæ quidem, sed stipitem fere attingentes.

Obs. Præter colorem pallentem β . ab α . nullo modo differt.

48. *Agaricus vulgaris*, gregarius, pileo membranaceo papillato, demum subumbilicato subviscoso, cinereo striato, lamellis albidis, stipite crassiusculo viscoso.

Ag. vulgaris. *Perf. Syn.* p. 394. *Albert. et Schwein.* p. 198.

In silvis, præsertim *Svecicis*, autumnno pluvioso, sat copiose.

Descr. Gregarius confertus parvus terrestris rarior epixylus. *Stipes* 1 - 2 unc. fistulosus $1\frac{1}{2}$ - 2 lin. crassus deorsum subincrassatus viscosus cinereus inferne pilis albis folia picea adhæret. *Pileus* membranaceus subviscosus, junior campanulatus mox planiusculus subumbilicatus cinereus striatus 2 - 3 lin. *Lamellæ* distinctæ albidæ 2-natæ uncinato-adfixæ; etiam subdecurrentes in pileo umbilicato, hinc omphaliis affinis.

Ad hanc sectionem pertinent e Svecicis *Ag. galericularius, polygrammus, spinipes, astivus* &c.

**** Lamellis albis, pileoque concolore.**

49. *Agaricus lacteus*, subgregarius albus, pileo membranaceo campanulato obtuso striato, lamellis subliberis, stipite longiusculo basi hirsuto.

Ag. lacteus. Pers. Syn. p. 394. Albert. et Schwein. l. c. p. 198.

β *pirhyus*, pilis radicalibus densis subradiatis. *Pers. l. c.*

Habitat in pinetis pasfim. Frequens in *Svecia*,

Descr. Subgregarius epigæus, raro epixylus, totus albus. *Stipes* fistulosus 2-3 unc. subfiliformis fragilis basi parum crassior adscendens hirsuto-radicatus, pilis densis albis folia pinea adnatus, superne glaber lævis. *Pileus* membranaceus campanulatus obtusus, rarissime umbonatus, striatus glaber $\frac{1}{3}$ - $\frac{1}{2}$ unc. haud ultra. Variat viscosus. *Lamellæ* candidæ subliberæ distinctæ alternæ l. æquales.

Lamellæ subadnatæ secund. Persoon.

50. *Agaricus tuberosus*, gregarius albus, pileo subcarnoso convexo lævi, lamellis adnatis, stipite brevi Sclerotiis adnato.

Ag. tuberosus. Pers. Syn. p. 374. Albert.

et Schwein. l. c. p. 190. Bull. t. 256 (in Scler. cornuto Fr.) et t. 522 (in Scl. Fungorum P.)

Ag. bulbosus. Schum. Siell. 2. p. 307.

Ag. amanitæ. Batseb. El. fung. Cont. 1. p. 109. t. 18. f. 93. Pers. Obs. myc. 2. p. 52. ~~æ~~ uberosus et β . Sclerotii.

Ag. alumnus. Bolt. Fung. p. 155. t. 155.

β . cirratus, stipite deorsum piloso. Pers. Syn. l. c. β . Obs. myc. l. c. γ . Alb. l. c. β .

Habitat intra Agaricos putridos et muscos in Sclerotio muscorum, fungorum, cornuto &c. — Passim in Suecia.

Descr. Gregarius confertus parvus in fungis parasiticus. Sripes subfibrillosus tenuis $\frac{1}{2}$ unct. et ultra subæqualis $\frac{1}{2}$ lin. crassus varie flexus plus minusve glaber basi variis Sclerotiis innatus. Pileus 1–2 lin. latus convexus tenuissime carnosus lævis glaber albus papillatus. Lamellæ alternæ adnatæ tenues subdistinctæ candidæ.

Obs. Quod de basi hujus fungi polymorpha attulerunt Auctores, ad Sclerotia, quibus innascitur, pertinet. Hinc etiam duæ varietates in Obs. myc. l. c. Vide alio loc.

51. *Agaricus stylobates*, solitarius albidus, pileo membranaceo convexo striato, lamellis plano-adscendentibus, stipite basi membranulæ orbiculari imposito.

Ag. stylobates. *Perf. Syn.* p. 390. t. 5. f. 4.
Schum. Sall. 2. p. 267. *Albert. et Schwein. l. c.*
 p. 196.

Habitat in Germania, Svecia, Dania. A me
 variis locis lectus ex. gr. in foliis arborum, vac-
 cinii vitis id. et in *Rhizomorpha subcorticali* et
 ligno putrido et præsertim copiose in pinetis ad
 terram supra folia pinea putrescentia.

Descr. Solitarius, rarissime gregarius.
Stipes fistulosus uncialis et ultra, in speciminibus
 epipsyllis fere capillaris, in epigæis firmior et
 sæpe transversim corrugatus, glaber albus, basi
 membranula orbiculari convexa pilosa 1-2 lin.
 albida l. fuscescenti. *Pileus* membranaceus con-
 vexus 3 lin. glaber striatus pellucidus albidus, de-
 mum cinerascens. *Lamellæ* albidæ plano-adscen-
 dentes demum subliberæ 2-4 natæ semiobovatæ.

Agar. nanus. Bull. Specificè vix differt.
 Sequentem hujus subspeciem, valde notabilem, in
 trunco putrido semel (18. Jul.) legi.

52. *Agaricus dilatatus*, solitarius, pileo
 planiusculo membranaceo vertice lævi, lamellis
 liberis, basi membranulæ orbiculari imposito.

In trunco betulino.

Descr. Species vix est distincta, sed nec
 pro mera pror's varietas habenda. *Stipes* $1\frac{1}{2}$ unc.
 fistulosus fragilis lævis, nec corrugatus, glaber

albus deorsum subincrassatus et membranula orbiculari concolore striata glabra convexa 2-lineari impositus. *Pileus* convexo-planiusculus membranaceus albidus uncialis, vertice lævi margine subdeflexo sulcato-striato, striis subramosis. *Lamellæ* albæ distinctæ semilanceolatæ 4-natæ liberæ.

53. *Agaricus integrellus*, fascicularis candidus, pileo membranaceo hemisphærico striato, lamellis decurrentibus distantibus subintegræ, stipite deorsum pubescente.

Ag. integrellus. Pers. Syn. p. 393. Ic. et descr. Fung. p. 54. t. 13. f. 5. Schum. Sall. 2. p. 336.

β. lignatilis. Albert. et Schwein. l. c. p. 197.

In umbrosis supra folia dijecta et ramulos putrescentes, rarior in Germania, Dania, Suecia.

Descr. Fungus polymorphus, totus semper candidus, fascicularis l. saltem gregarius. — *Stipes* fistulosus sæpe uncialis et ultra $\frac{1}{2}$ — $1\frac{1}{2}$ lin. crass., suberectis firmulus basi tenuissime incrassatus et pubescens. *Pileus* membranaceus junior conicus, mox hemisphæricus subdifformis basi 2–3 lin. latus, sed explanatus interdum fere semiunc. glaber tenuissime striatus. *Lamellæ* integræ distantes rugæformes subdecurrentes marginem sæpe vix attingentes.

Obs. Descr. nostra a Persoonii parum recedit, sed idem omnino fungus.

*** Lamellis albis, margine discoloribus.

54. *Agaricus rubromarginatus*, pileo campanulato - convexo membranaceo obtuso cinerascenti - pallido, lamellis adnatis albis purpureo-marginatis, stipite exsucco.

In pinetis *Sveciæ* supra folia et strobilos *Pini Abietis*.

Descr. Species maxime distincta, et *Ag. sanguinolento*, cui descriptio proxima, ne affinis quidem. *Stipes* fistulosus 1 - 2 unc. erectus l. basi curvatus subtenuis albidus æqualis glaber lævis basi fibrillosus exsuccus. *Pileus* junior campanulatus mox convexus non papillatus cinerascenti - pallidus, rarius rufescenti - cinereus $\frac{1}{2}$ unc. et ultra. *Lamellæ* distinctæ firmæ totæ adnexæ 2 - natæ albidæ, margine purpureo - rubræ.

**** Lamellis roseis, rubris &c.

55. *Agaricus rhodellus*, gregarius roseus, pileo campanulato papillato striato, stipite filiformi pallescente fistuloso, basi fibrilloso.

Agar. roseus. Pers. Syn. p. 393. t. 5. f. 3.

Ag. fistulosus. Bull. t. 518 inter Litt. D.

& P.

In pinetis præcipue abietinis. — Etiam in *Sveciæ*, cum *strobilino*, qui ad hanc sectionem pertinet.

Descr. Gregarius terrestris. *Stipes* fistulosus 1-3 uncial. strictus lævis glaber pallide rubicundus, basi villis albidis vestitus. *Pileus* membr. campanulatus papillatus demum convexus roseus striatus glaber, explan. diam. $\frac{1}{4}$ - $\frac{1}{2}$ unc. *Lamellæ* dilutiones distantes 2-4 natae arcuato-adfixæ.

Nomen triviale, ne cum *A. roseo* Sowerb., Bull., Witter., Swartz. in *Vet. Ac. Handl.* 1808. p. 84 l. cum *A. roseo* Schaff., Pers. Obs. myc. l. cum *rosaceo* P. et *rosello* Batsch. confundetur, licet invitus, mutavi.

56. *Agaricus pterigenus*, subgregarius dilute roseus, pileo convexo obtuso lævigato, stipite capillari pallescente solido, basi glabro. *Icon. vidi Mscr. ad Flor. Dan. apud Cel. Prof. Hornemann.*

In stipidibus dejectis, præsertim Filicibus, locis umbrocis humidis *Sveciæ*, *Daniæ*.

Descr. Subgregarius tenuissimus. *Stipes* 1-3 unc. capillo tenuior solidus (saltem ob tenuitatem solidus videtur) æqualis tenax flexilis dilutissime roseus, basi glaberrimus. *Pileus* tenerissime membranaceus junior subglobosus deinde convexus 1-2 lin. glaber sublævis non papillatus, quin potius demum subumbilicatus, sed *Omphal.* species non est, dilute roseus. *Lamellæ* paucæ distinctæ concolores latiusculæ adnatæ.

***** Lamellis flavescentibus, citrinis
ochraceis et rubiginosis.

57. *Agaricus citrinellus*, gregarius, pileo hemisphærico papillato striato stipiteque breviusculo basi piloso citrinis, lamellis uncinatis dilutioribus.

Ag. citrinellus. *Perf. Syn.* p. 384. *Ic. et descr. Fung.* p. 44. t. 11. f. 3.

Ag. tenellus. *Batsch. El. fung. Cont.* 1. p. 103. t. 18. f. 88.

In pinetis; rarior in *Svecia*.

Descr. Parum quidem fungus noster et a *Perfoonii* et a *Batschii* recedit, sed a neutro differt. *Stipes* fistulosus 1-2 unc. filiformis æqualis glaber l. subviscosus rectus lævis citrinus deorsum subobscurior. *Pileus* membranaceus striatus citrinus vertice striisque obscurioribus 2-3 lin. glaber, junior conicus dein late campanulatus. *Lamellæ* albido-citrinæ integræ l. 2-natæ arcuato-adnexæ.

58. *Agaricus elongatus*, solitarius, pileo hemisphærico lævi flavescenti, lamellis adnatis subconcoloribus, stipite longissimo fibrilloso.

Ag. elongatus. *Perf. Syn.* p. 384. *Ic. et descr.* p. 3. t. 1. f. 4.

In paludosis inter *Sphagna*; passim in *Svecia*.

Descr. Solitarius terrestris. *Stipes* 4-6 unc. fistulosus crass. pennæ gallinæ rufospadiceus, superne glaber lævis, deorsum villoso-fibrillosus radicans. *Pileus* convexus flavescent 3-4 lin. latus lævis glaber. *Lamellæ* latiusculæ concoloræ basi rotundatæ adnatæ.

Cum varietatibus *Agar. hypnorum* præcipue *γ. Sphagnorum* et *δ. rubiginoso* (*Ag. rubig. Pers.*) non confundendus, a quibus pileo lævi mox distinguitur.

Obs. In fungum nostrum omnino quadrat descriptio *Persoonii*, sed pileum virescentem ut l. c. depictus, nondum vidi. Pictoris forte culpa.

59. *Agaricus carpophikus*, subgregarius, pileo convexo lævi pallescente, lamellis subliberis ochraceis, stipite brevi solido.

Bull. t. 24. f. 2. subsimilis, sed aplo major.

In pericarpis nucum *Fagi* dejectis semiputridis, in umbrosis humidis in *Svecia*.

Descr. Solitarius l. gregarius. *Stipes* solidus? uncialis capillaris rectiusculus glaber lævis pallidus l. subalutaceus, basi fibrillis adfixus. *Pileus* membranaceus convexus 2 lin. latus lævis glaber obtusus fordide pallescens l. pallescenti-albidus. *Lamellæ* ochraceæ latiusculæ liberæ l. levisime adnatæ.

60. *Agaricus tener*, subsolitarius ochraceo-ferrugineus pileo conico glabro striatulo, lamellis adscendentibus, stipite longiusculo fistuloso glabro.

Ag. tener. *Perf. Syn.* p. 386. *Schæff. F.* t. 70. *Sowerb. F.* t. 33.

Habitat in pratis et graminosis; passim in *Svecia*.

Descr. Solitarius l. sparsus, terrestris tener fragilis. *Stipes* fistulosus 2-4 unc. 1 lin. crassus subæqualis l. deorsum tenuissime incrassatus cylindricus lævis glaberrimus striatulus obtusus flavo-ferrugineus 3-5 lin. altus, explan. uncialis. *Lamellæ* subobscuriores cinnamomeæ 2-4 natæ adscendentes.

61. *Agaricus pilosellus*, subsolitarius ochraceo-ferrugineus, pileo conico piloso læviusculo, lamellis adnatis, stipite longiusculo fistuloso villoso.

Ag. pilosellus. *Perf. Syn.* p. 387.

Ad truncos dijectos, raro. In *Svecia* truncos fagineos putridissimos incolit solitarie; in abietino gregarium legerunt in Germania *Albert. et Schwein.* l. c. p. 194.

Descr. Solitarius fragilis truncigenus. *Stipes* fistulosus 1-2 unc. filiformis rectus ochraceo-ferrugineus basi subradicatus, villo denso

tenui albido undique vestitus. *Pileus* conicus l. campanulatus membranaceus ochraceo-ferrugineus superne piloso læviusculo 3 lin. altus et latus. *Lamellæ* subconcolores latiusculæ 4-natæ adfixæ; dentic. decurrente.

***** *Lamellis cyaneis, cinereis &c.*

62. *Agaricus Batschii*, solitarius atrocyaneus, pileo cuculato striato, margine lamellis dilutioribus, stipite mediocri striato.

Ag. atrocyaneus. *Batsch. El. fung. Cont. 1. p. 101. t. 18. f. 87. (a Persoonio ad Ag. polygrammum relatus).*

In graminosis Germaniæ, Sveciæ rarissime.

Descri. Solitarius terrestris. *Stipes* fistulosus uncialis æqualis glaber striatus obscure cyaneus violascentis, vix radicans. *Pileus* cuculatus l. subconicus obtusus nec umbonatus membranaceus glaber profunde striatus 3-4 alt. 2 lat. cyaneus demum subpruinatus, margine dilutiori. *Lamellæ* 2-4 natæ rectæ cyaneæ lanceolatæ subliberæ, margine albescunt.

Obs. Cum *A. atrocyaneo* Pers. non confundendus.

63. *Agaricus griseus*, solitarius cinereus, pileo convexo striato, margine concolori subpatente, lamellis canis, stipite mediocri lavi. *Fries.*

β. *Ag. griseus.* *Batsch. El. fung. Cont. 1.*
p. 87. f. 80.

In pineti ad Temsiö (16. Aug.) semel vidi.

Descr. *Stipes* fistulosus 1-2 unc. crassiusculus subfragilis glaber lævis canescenti-cinereus. *Pileus* convexus demum explanatus membranaceus striatus glaber obscure cinereus, exsiccatus canus, minime umbonatus $\frac{1}{4}$ unc. latus. *Lamellæ* latiusculæ canæ 2-natæ adfixæ. *Stipes* basi pubescens.

Obs. De *Coprinis*, Genere distincto
Conf. *Link in Schrad. n. Journ. 3. p. 12. t. 1.*
f. 16, 17, in secunda parte.

V. P R A T E L L A.

64. *Agaricus stipatus*, cæspitosus, pileo convexo lævigato fuscescente-livido, exsiccatione ruguloso pallescente, lamellis adnatis confertis incarnato-fuscis, stipite albicante lævi subnudo.

Ag. stipatus. *Pers. Syn. p. 423.*

Ag. spadiceus. *Schaff. Fung. t. 237.*

Ad truncos passim. In *Svecia* præcipue ad truncos fagineos.

Descr. Cæspitosus fragilis truncigenus. *Stipites* plurimi conferti 2-3 unciales fistulosi

erecti l. curvati cylindrici aquose pallidi glabri læves nudī. *Pileus* membranaceus, vertice subcarnoso, convexus $\frac{1}{2}$ - 1 uncial. glaber lævis margine striatulo plus minusve fuscescens lividus l. rufescenti-ispadiceus, siccatus expallens rugulosus. *Lamellæ* confertæ incarnato-fuscæ latiusculæ 4-natæ adfixæ subnebulosæ pulverulentæ.

65. *Agaricus aleuriatus*, subsolitarius, pileo subconico, demum explanato subdepresso, subviscoso, lamellis subliberis ochraceo-cinereis, stipite albido farinoso nudo.

Agar. farinosus. *Gmel. Syst. Linn. II. 2. P. 1414.*

Ag. farinulentus. *Schaff. Fung. t. 205.*

In ligno putrido fagineo, in *Svecia*, *Germania* &c.

Descr. Solitarius l. gregarius, nec cæspitosus. *Stipes* fistulosus $1\frac{1}{2}$ l. $2\frac{1}{2}$ uncial; crass. pennæ gallinæ l. anserinæ, rectus l. curvatus albus æqualis, basi interdum incrassatus, glaber, sed farina albida undique conspersus. *Pileus* junior conicus mox campanulatus demum planiusculus, rarius planus, vertice subdepresso, glaber subviscosus tenuissime carnosus dilute cinereus, vertice obscuriori subolivaceo fusco, 1 - 2 unc. læviusculus, margine striato. Colorem pilei etiam subroseo-cinerascentem et flavescentem vidi. *Lamellæ* sub-

renues, sed carne pilei multo latiores, inæquales stipitem approximatae, sed rotundato-liberae dilutæ ochraceo-cinereæ demum obscuriores.

Obs. E *synopsi Fungor.* hanc speciem exclusit et in *Comment. Fung. Schæff.* pro varietate *Ag. cernui* habendam esse suspicatus Cel. *Persoon.* Nomina illa: *farinosus*, *farinaeus*, *farinatus*, *farinulentus* &c. nimiam habent similitudinem et de cetero pluribus apud Auctores communia sunt.

66. *Agaricus cernuus*, subgregarius, pileo hemisphærico albido, lamellis adnatis cinereo-nigricantibus, stipite albido nudo lævi subcurvato.

Ag. cernuus. *Fl. Dan.* t. 1008.

Ad truncorum radices in *Svecia*, *Dania* &c. In fimo. *Vabl.*

Descr. Alia dubia Mycologis species, ab *A. aleuriato* Fr. vere distincta. Gregarius. *Stipes* fistulosus fragilis 2 uncial., crass. pennæ gallinæ, arcuato-adscedens nudus lævis albus, absque pulvere. *Pileus* membranaceus hemisphæricus uncialis subfragilis albido-l. flavescenti-cinereus glaber læviusculus, ob stipitem sæpe curvatum nonnunquam cernuus. *Lamellæ* adnatæ cinereo-nigricantes, juniores dilutiores, 4-8 natæ distinctæ.

Etiam hujus nomen triviale non optimum.

67. *Agaricus sarcocephalus*, subsolitarius, pileo subhemisphærico carnoso flavescente, lamellis cinereo-fuligineis adnatis, stipite longissimo ventricoso glabro.

In silvis Sveciæ, rarior. Augusto legimus.

Descr. Solitarius, raro gregarius, terrestris. Stipes fistulosus strictus 4-6 unc., penna gallina parum crassior, glaber flavescens rugato-flexuoso-ventricoso. Pileus conico-convexus deinde hemisphæricus convexus carnosus glaber flavescens sublævis l. margine rugulosus, explanatus ultra uncial. Caro alba a vermibus sæpe erosa. Lamellæ cinereo-fuliginæ margine albicantes latiusculæ 2-4 natæ adfixæ denticulo decurrente.

Vera omnino Pratellæ species. A *Persoonianis* distinctissima.

68. *Agaricus polytrichi*, subgregarius, pileo convexo subcarnoso flavescente, lamellis adnatis albidis dein cinerascanti-fuligineis, stipite longissimo æquali fibrilloso subannulato.

Ag. mutabilis. *Batsch. El. fung. Cont. 2. t. 38. f. 208 fungum meum bene refert, sed ab A. caudicino, ad quem retulit Cel. Persoon, distinctissimus.*

In humidis Sveciæ inter *Polytricha* &c.

Descr. Stipes fistulosus 2-3 unc., crassit. pennæ gallinæ, subæqualis flavescens lævis glaber

deorsum fibrillosus. *Annulus* albus fugacissimus. *Pileus* convexus fere membranaceus tenuissime carnosus $\frac{1}{2}$ - 1 unc. lævis glaber flavesceus, margine deflexo. *Lamellæ* primo albidae dein cinerascens fuligineo-nebulosæ latiusculæ adfixæ.

O b s. In animadversariis etiam descripsi varietatem *stipite flexuoso et lamellis subliberis*.

69. *Agaricus pascuus*, subsolarius, pileo membranaceo conico dein dilatato cinereo-umbrino, lamellis liberis latiusculis incarnatis, stipite glabro nudo æquali.

Ag. pascuus. *Pers. Syn.* p. 427. *Albert. et Schwein.* l. c. p. 207.

Ag. pyramidatus. *Schæff. Fung.* t. 229.

Ag. irregularis. *Bolt. Fung.* t. 13.

In graminosis et pascuis; passim in *Suecia*, *Germania*, *Anglia* &c.

Descr. *Stipes* fistulosus 2-3 unc., crassit. pennæ gallinæ l. ultra, fragillimus subæqualis cylindricus glaber lævis pileo concolor l. dilutior cinerascens. *Pileus* membranaceus conicus mox campanulatus subconvexus obtusus subflexuosus subaquosus fragilis glaber striatus spadiceus l. sæpius fuligineo-pallens (vid. *Alb. et Schw.* l. c.) explanatus 1 unc. *Lamellæ* latiusculæ 2-natæ

subliberæ exalbido-rosæ demum incarnato-rufæ
margine subdenticulatæ.

Vix specie differt:

70. *Agaricus fuliginus*, gregarius, pileo
e campanulato-convexo cinereo-livido, lamellis
liberis tenuioribus, stipite glabro nudo deorsum
incrassato.

Ag. fuliginus. *Pers. Syn.* p. 427.

Habitat quoque in consortio prioris.

Descr. A præcedenti, cui proximus, dif-
fert fungus noster: Stipite subbrevisiori deorsum in-
crassato cinereo, pileo læviusculo sed demum sub-
ruguloso, striis tamen obscurioribus sæpe picto,
tenuioribus roseis et integerrimis lamellis.

71. *Agaricus montanus*, gregarius, pi-
leo subcarnoso convexo fuscescente, lamellis sub-
decurrentibus umbrinis, stipite brevi fibrilloso
æquali.

Ag. montanus. *Pers. Syn.* p. 428 (exclus.
var.) *Obs. Myc.* 1. p. 9.

β. *xylarius*, truncigenus, stipite fuscescenti-
albido, lamellis adnatis.

γ? *coriarius*, cæspitosus, pileo repando, sti-
pite farinoso, lamellis rubris.

Ag. montanus. *Albert. et Schwein.* l. c.
p. 207. γ γ.

Ipse var. β . tantummodo in lignis putridissimis ad terram locis sylvaticis dejectis legi.

Varietas β . *Persf.*, me iudice, vix hujus loci. Vide inter *Omphalias*.

Obs. Descriptio *Persoonii* in fungum meum β . bene quadrat, ut vix sit distinguendus nisi loco natali et lamellis minus decurrentibus.

Var. γ . vero dubia. *Ag. obtusus*, quoad locum natalem propius accedit, sed stipite deorsum incrassato &c. differt. An var. β . ad hanc potius referenda?

Descr. β . *Stipites* plures gregarii, rarius vero cæspitosi, unciales nec ultra subrecti fuscescentes subalbidi æquales, crass. 1 lin. subfibrillosi. *Pileus* membranaceo-subcarnosus $\frac{1}{2}$ unc. hemisphæricus obtusus absque umbone, demum plano-convexus fuscescens l. spadiceus (exsiccatus fere canescens) glaber subhumidus lævis margine striato. *Lamellæ* lariusculæ 4-natæ umbrinæ nebulosæ totæ adnexæ, raro tamen decurrentes.

72 *Agaricus renax*, subsolitarius, pileo subcarnoso campanulato flavo, lamellis adnatis nebulosis, stipite longiusculo glabro nudo.

In silvis inter folia *Pini Abietis*, Septembri.

Descr. Species optime distincta a reliquis, ut e descriptione patebit. *Stipes* substulosus rectus 2-4 uncialis cylindricus subæqua-

lis, interdum, forte monstrositas, medio angustatus, glaber lævis albido-fuscus aliusve coloris. *Pileus* subcarnosus campanulatus flavus l. ochraceus, margine aquose-flavescens glaber lævis explanatus 1½ - 2 uncialis. *Lamellæ* latiusculæ 4-natæ affixæ nebulosæ *subvirescentes*. *Substantia* stipitis et pilei tenax!

VI. LACTARIUS.

* Pileo albido.

73. *Agaricus pergamenus*, pileo planiusculo subdepresso candido subcoriaceo, lamellis fordide albidis, stipite obscuriori.

Agar. pergamenus. Swartz in Vet. Acad. Handl. 1809. p. 90.

Ag. urens. Schum. Sell. 2. p. 343.

Ag. piperatus. Batsch. El. fung. Cont. 1. p. 34. t. 13. f. 59. in fungum meum bene quadrat.

In Svecia, Dania, Germania locis umbrosis subhumidis.

Obs. Descriptionem optimam dedit *Illustr. Swartz l. c.* Stipes variat exalbido-coerulescens, spadiceus et badius. Hinc synonymon *Schumacheri* huc etiam retuli.

Hujus loci e Svecicis *Ag. Listeri* Sowerb.,
controversus P.

** Pileo pallido, cinerascenti, flavescenti, &c.

74. *Agaricus pubescens* pileo albido azono subumbilicato pubescente, lamellis adnatis albo-lactescentibus, stipite pubescente.

Ag. pubescens. Schrad. *Spic.* p. 122.

In muscosis rarissime. Semel in *Suecia* ad Femsjö legi.

Descr. Stipes breviusculus uncialis, crass. digiti, solidus teres interdum parum compressus subæqualis carnosus lævis albidus, pubescentia concolori molli brevi subadpressa ubique tectus. *Pileus* sub-2-uncialis carnosus planus, vertice depressus, albidus lævis fuscus pubescens, margine deflexo subinvoluta vilioso. *Caro* alba. *Lamellæ* albæ confertæ affixæ nec decurrentes carne pilei angustiores.

Obs. *Ag. piperati* L. (*torminosi* Auct.), varietas esse videtur.

75. *Agaricus pallidus*, pileo incarnato pallido umbilicato subviscoso, lamellis subdecurrentibus albo-lactescentibus, stipite glabro, basi subpubescente.

Ag. pallidus. *Pérf. Syn.* p. 431. *Albert. et Schwein.* l. c. p. 209.

Ag. incarnatus. *Pérf. disp. Fung.* p. 64.

In silvis umbrosis, præsertim betulinis frequens. Inter *Svecicos* tamen non attulit *Celeb. Swartz* l. c.

76. *Agaricus intermedius*, pileo flavescenti luteo umbilicato viscoso margine involuto tomentoso, lamellis dilutioribus exalbido-flavo-lactescentibus, stipite deorsum pubescente æquali.

Agar. thejogalus. *Swartz.* l. c. p. 91. (*exclus. syn.*)

Ag. thejogalus. β. luteus. *Alb. et Schw.* l. c. p. 208. (*Descriptio optima*)

Ag. crinitus. *Schæff. Fung.* t. 228.

Ag. scrobiculatus. *Schæff.* l. c. t. 227.

In silvis abietinis muscosis humidis.

Descr. *Stripes* nunc solidus, nunc cavus $1\frac{1}{2}$ – $2\frac{1}{2}$ unc. crasfit. $\frac{1}{2}$ unc. æqualis teres subfirmus flavescent l. albidus sæpius maculis pallidioribus pictus glaber deorsum pubescenti-radicatus. *Pileus* carnosus, junior convexiusculus demum subinfundibuliformis latissimus (3–12 unc., ut optime monuerunt *Albert. et Schwein.* l. c.) late luteus l. flavescent, recens viscosus villo ochraceo-fulvo adglutinato, margine, ut in *torminoso*, involuto tomentoso. *Substantia* alba. *Lac album*,

mox sulphureum. *Lamellæ* 4-natæ affixæ angustatæ approximatae exalbido-flavescentes.

In betuletis siccis legimus varietatem *expal-
lentem*, cujus

Descr. *Stipes* cavus vix uncialis albus subæqualis l. deorsum angustatus pubescens. *Pileus* carnosus 4-5 unc. subsiccus fere infundibuliformis subglaber albidus, margine patente l. erecto, rarius deflexo, et vix involuto. *Lac* omnino prioris. *Lamellæ* 6-8 natæ subsimplices albidae decurrentes.

Obs. Medius omnino inter *Ag. torminosum* Schæff. (*piperatum* Linn.) et *thejogalum* Bulliard. Ab utroque vero, ut videtur, distinctissimus. Ad illum, me iudice, propius accedit et ad quem quoque icones *Schæfferi* referendas esse suspicatus est *Persoon* in Obs. Myc. 2. p. 41, cum quo etiam optime comparavit illustr. *Swartz* et hinc fungum ejus haud dubius retuli. *Ag.* vero *thejogalus*, quem in Dania vidi, habitu et colore quam maxime differt; sed ob lactis colorem vnic affinis. Synonyma hujus, quæ vera videntur:

Ag. thejogalus. *Perf. Syn.* p. 431. *Bull. Herb.* t. 567. f. 2. *Schum. Sall.* p. 347.

77. *Agaricus flexuosus*, pileo rufo-cinerascente subpallido lobato-flexuoso duro azono,

lamellis albo-lactescentibus flavis, stipite brevissimo cinerascente glabro.

Ag. azonatus. *Bull.* l. c. t. 559. f. 1.

In filvis. In *Svecia* legi.

Descr. Gregarius. *Stipes* solidus brevis uncialis, crass. fere unciali, firmus carnosus, deorsum subangustatus difformis, vix teres, glaber lavis cinerascens. *Pileus* carnosus firmus undulatus flexuoso-lobatus rufo-cinerascens azonus 2-4 uncial. glaber lavis fuscus. *Lamellæ* flavæ totæ adnatæ polynatæ interdum furcatæ.

Vix differre videtur *A. flexuosus* Pers. Syn. p. 430. Quam plurima alia Synonyma, licet dubia, afferre possem; sed species plurimæ affines, satis tamen distinctæ, *acris*, *luridus* &c. ex figuris difficiliter, quam e descriptionibus determinantur.

*** Pileo umbrino aut fuliginoso nigrescente.

78. *Agaricus Necator*, pileo ex olivaceo-umbrino, disco subdepresso zonato, margine tomentoso-involuto, lamellis albolactescentibus, stipite subolivaceo.

Ag. *Necator*. Pers. Syn. p. 435. *Bull. Herb.* t. 43. *Albert. et Schwein.* p. 210.

In Gallia, Germania passim, in *Svecia* rarior.

Minime varietas *A. piperati*. L.

Descr. *Stipes* brevis uncialis, crass. digiti intermedi, firmus solidus, raro demum cavus glaber lactifluus fuligineo-umbrinus l. subolivaceus laxis elasticus intus carnosus albus. *Pileus* carnosus 2-4 uncialis convexiusculus, vertice depresso olivaceo-umbrino l. subfuligineo-olivaceo, laxis glaber concentrice subzonatus, margine tomentoso involuto. *Caro* alba firma parce lactescens. *Lamellæ* pallido-albidæ adfixæ sæpe dichotomæ 2-4 natæ.

79. *Agaricus blennius*, pileo cinereo-olivaceo plano depresso viscoso margine patente, lamellis albo-lactescentibus, stipite cinerascente.

Krappf. Esb. Schw. I. t. vi. f. 1, 2.

In fagetis *Sveciæ* non infrequens.

Descr. *Stipes* solidus $1\frac{1}{4}$ unc. deorsum angustatus glaber rectus l. basi interdum incurvus elasticus laxis pallidus cinereus lactescens. Variat stipite æquali. *Pileus* carnosus primo planiusculus dein vertice depressus cinereo-olivaceus valde viscosus 2-4 unc.; ceterum nec villosus nec striatus, margine glabro patente. Siccus observatur interdum coelo sereno, sed humectatus vox viscosus. *Caro* superne cinerascens, inferne alba lactiflua. *Lac* album acre immutabile. *Lamellæ* adnatæ albæ inæquales.

Species vere distincta et priori vix affinis.

Figura citata fungum meum bene refert. — Hujus etiam loci esse videntur:

Agar. xylophilus β . viscosus. *Perf. Syn.*

P. 438 et

Ag. curtipes. *Schum. Sall. 2. p. 343.*

80. *Agaricus trivialis*, pileo obscure cinerascanti-incarnato depresso subviscoso, lamellis albo-lactescentibus, stipite elongato albidō.

Krapf. Esb. Schw. I. t. 5. f. 1, 2, 3.

In pinetis sat frequens.

Deser. *Stipes* elongatus 2–4 uncial. crassus cavus subæqualis l. inferne parum crassior subfragilis lævis superne glaber deorsum pubescens. *Pileus* planiusculus, vertice depresso, subinfundibuliformis carnosus 2–4 unc. lævis ficcus l. viscosus, quod a coeli temperie pendet, obscure cinereo-incarnatus, margine subreflexo. *Lamellæ* albidæ subdecurrentes inæquales, venis sæpe connexis, interdum dichotomæ carne pilei basi angustiores, apice latiores. Variat lamellis adnatis. *Lac*, ut in priori, album subacre, non vero acerrimum, immutabile.

Varietatem l. potius monstrositatem, pileo depresso piloso-squamuloso, succo aquoso, stipite rufescente subvillosa eisdem locis semel observavimus.

Ag. pallido proximus, sed colore et magnitudine valde diversus. Ab *A. lurido* lacte immutabili pluribusque aliis differt characteribus. Habitu quodam triviali semper distinguitur.

81. *Agaricus acris*, pileo cinereo-fuligineo azono planiusculo, lamellis exalbido-rufescenti-lactescentibus demum alutaceis, stipite cinerascente.

Agar. acris. *Perf. Syn.* p. 437. *Alb. et Schwein.* l. c. p. 216. *Bolt. Fung.* 2. t. 60.

Ag. opacus. *Batsch.* f. 68.

In silvis *Sueciæ*, *Germaniæ*, *Daniæ*, *Angliæ* &c. minime rarus.

Descr. *Stipes* nunc cavus nunc solidus 2-3 unc. æqualis, crass. pennæ anserinæ l. digiti minimi, lævis subelasticus pallide cinerascens glaber basi subpubescens. *Pileus* carnosus planus subdepressus azonus cinerascens cinereo-lividus l. fuliginosus, vertice nonnunquam obscuriori fuligineo, 2-3 unc. lævis glaberrimus, margine junioris reflexo mox patente. *Caro* pallida alba. *Lamellæ* adnatæ alutacæ l. ochraceæ subdistantes inæquales. *Lac* album demum rufescenti-lateritium.

Ab *Ag. lurido* B. Swartz l. c. p. 212, cui affinis, differt: *stipite* subcavo elastico, *pileo* azono cinereo-livido, *lamellis* mox alutaceis.

Inter plures novas hujus, quas legimus,

varietates, unam, quam diximus *Agalum*, describamus: *Stipes* elongatus 1-uncialis, crass. $\frac{1}{2}$ - $\frac{3}{4}$ unc., subæqualis sæpe curvatus subsolidus intus spongiosus glaber lævis. *Pileus* carnosus 3-4 unc. planiusculus subdepressus lævis ficcus glaberrimus, margine concolore sæpe flexuoso. *Lamellæ* subcroceæ substeriles 4-8 natæ adfixæ carne pilei marginem versus latiores. *Odor* non ingratus! *Sapor* et *Lac* fere nullum! et hujus tamen omnino loci: colore pilei et substantia ab *Agar. flexuoso*, antea descripto, quam maxime differt.

82. *Agaricus pyrogatus*, pileo plumbeo-livido subzonato planiuscolo, lamellis distinctis albido-lactescentibus pallidis stipite recto fuliginoso. — *Perf. Syn.* p. 436 *cum synonym.*

In silvis *Svecicis* sequentes hujus legimus varietates:

α) *obsoletum*: pileo obsoleto-zonato, stipite cavo.

Descr. *Stipes* 2-unc. et ultra rectus, crass. digiti minoris, subcavus cinereo-fuliginosus lævis l. sublacunofus, interdum scrobiculatus glaberrimus. *Pileus* carnosus plumbeo-lividus planus interdum excentricus, vertice subdepresso, 2-3 unc. glaber lævis; margine patente, zonis obsoletis l. nullis.

Lac album acre immutabile. *Lamellæ* pallidæ
l. albidæ adfixæ inæquales polynatæ subdecur-
rentes carne pilei æquales.

- β) *disfimilem*: pileo azono, stipite solido breviori.
Differt: *Stipite* crass. pennæ cygneæ basi sub-
pubescente læviusculo elastico, *Pileo* subcarnoso
cinereo l. spadiceo-cinereo depresso excavato
fere infundibuliformi, zonis nullis. *Lamellæ*
sunt flavæ decurrentes 4-natæ carne pilei latio-
res. — Pileus in hac varietate subrugulosus.

Agar. plumbeus P. ab hac variet. vix nisi
magnitudine differt.

- γ) *elongatum*: pileo azono, stipite demum cavo
longissimo.

In hoc fungo *Stipes* junior solidus mox
cavus 4-6 unc. sursum reticulatus, olivaceus
l. olivaceo-plumbeus. *Pileus* 2 unc. vere car-
nosus planiusculus depressus nigricanti-fuscus
rugosus, margine duplicato-crenatus l. flexu-
oso-lobatus. *Lamellæ* albæ parum decurrentes
4-natæ.

Inter *Polytricha* &c. sæpius viget.

VII. R U S S U L A .

(Distinctum forte Genus. Conf. *Link.*
in *Schrad. Journ.* l. c.)

* Pileo albido.

83. *Agaricus galochrous*, pileo albo margine lævi, lamellis stipiteque mediocri concoloribus. (*Russula galochroa*. Fr.)

Agar. pectinaceus. *Bull.* t. 509. f. L.

Ag. lacteus. *Perf. Syn.* p. 439 (*sed minime confundendus cum lacteo Ejusd.* l. c. p. 394.)

In silvis præsertim umbrosis et fagetis *Sueciae* &c.

Obs. Plurimas *Russulae* species iterum conjungendas esse nullum dubium est et optime jam monuerunt *Alb. et Schwein.* l. c. p. 211 sq. Descrip. vero accuratos, ex quibus pateat summa affinitas, superfluas non fore speramus.

Descrip. *Stipes* elongatus 2-3 unc., diam. $\frac{1}{2}$ - $\frac{3}{4}$ uncial. carnosus intus medulla spongiosa fartus basi subangustatus lævis glaber albus. *Pileus* carnosus planiusculus mox depressus concavus albus rectus glaber lævis 3 uncial., margine nudo, absque tuberculis, interdum substriatus. *Lamellæ* albidæ crassiusculæ subdecurentes æquales, nonnullis ramosis.

Variat pileo virescente: *Agar. aruginosus*.
Schum. Sall. p. 246.

84. *Agaricus chioneus*, pileo niveo margine tuberculoso, lamellis stipiteque brevi concoloribus. (*Rusfula chionea*. Fr.)

Ag. niveus. *Perf. Syn. p. 438. Albert. et Schw. l. c. p. 211. (Nomen plurimis commune.)*

In umbrosis cum *priori*.

Ab antecedenti satis forte distincta, sed *fragili, depallenti, exalbicanti* P. maxime affinis. — Species enim enumeratæ ruborem suum sæpe plane omittunt; quod etiam testantur maculæ illæ rubellæ in hac specie sæpe obviæ.

Descr. *Stipes* uncialis glaber lævis æqualis candidus subfirmus, crass. digiti minoris, subcavus l. intus spongiosus non tamen eodem modo ac in præcedenti. (In illo enim substantia spongiosa medullam a stipitis liberam format, in hac vero confluunt.) *Pileus* planiusculus albus 2 unc. glaber lævis carnosus, margine striatus tuberculatus. *Lamellæ* subtenues distinctæ adnatæ albæ. *Sapor* (in nostro) amaricano.

85. *Agaricus leugethejus*, pileo albido margine striato, lamellis stipiteque flavescentibus, (*Rusfula leucoxantha*. Fr.)

In quercetis inter muscos rarior. In *Suecia*.

Descri. Cui affinis sit ignoro et, nostro iudicio, vere est distinctus. *Stipes* carnosus vere solidus, nec medulla spongiosa fartus, uncialis et ultra lævis glaber cylindricus flavidus, basi parum incrassato. *Pileus* depressus subconcauus pallide albidus glaber uncialis et ultra medio subcarnosus margine membranaceus fere striatus subtuberculofus patens. *Lamellæ* flavescentes subtenues æquales basi angustissimæ, ut stipitem vix attingunt.

**** Pileo rubro.**

86. *Agaricus Linnaei*, pileo planiusculo subdepresso margine substriato rubro, lamellis integris albis, stipite albo rubellove. (*Rusfula Linnaei*. Fr.)

Agar. integer. *Linn. et Auctorum! Fl. Svec.* p. 449. *Schaff.* t. 15, 16. fig. plur. &c. *Sowerb.* t. 201.

Ag. pectinaceus. *Bull. Herb.* t. 509. f. plur.

In silvis vulgatissimus.

Qui hanc speciem attente observaverit, omnino videbit, sequentes a me allatas varietates huius esse loci. (Conf. *Albert. et Schwein.* l. c.)

α) *emeticus*, pileo depresso margine fulcato sanguineo-rubro, stipite sæpe rubicundo.

Agar. emeticus. *Persf. Syn.* p. 439.
Swartz l. c. p. 213. *Schæff. t.* 15. *Bull.*
l. c. fig. ~~4. u. 2.~~ *Krapf. Est. Schw. f.* 2, 3.
Schum. Sell. 2. p. 245.

- β) *sanguinarius*, pileo explanato subdepresso
 margine substriato obscure rubro, stipite
 albo.

Agar. *sanguinarius*. *Schum. Sell.* 2.
 p. 244. *Bull. Champ.* ~~R.~~ ^{t.} 42. *Elroib t.* 2.
f. 2. c, d. *Schæff. Fung. t.* 16. *f.* 4, 5, 6.
Fl. Dan. t. 1009 &c.

- γ) *pulcherrimus*, pileo planiusculo rubro ni-
 tente, margine substriato, stipite toto roseo.

In graminosis subhumidis frondosis.

Descr. *Stipes* longus 3-4 uncial.
 solidus fragilis subtenuis lavis glaberrimus.
Pileus 1-2 unc. subtenuis subcarnosus pla-
 niusculus subdepressus, cum stipite latis-
 sime roseo-ruber, margine tenuissime stria-
 tus, interdum inflexus. *Lamellæ* candidæ
 rarissime inæquales et rubromarginatæ.

Sequentibus proximus. Sapor ut in
 omnibus varietatibus nunc dulcis nunc
 acerrimus.

- δ) *fragilis*, pileo depresso rubescente-subopaco,
 margine tuberculoso-striato, stipite albo.

Agar. *fragilis*. *Persf. Syn.* p. 440.

Schum. Sell. 2. p. 245. Albert. et Schwein.

l. c. p. 212.

Obs. Colore a priori et margine profundius striato differt.

- ε) *vagus*, pileo explanato depresso obscure rubro striato, stipite deorsum angustato rubro.

sanguinario proximus.

- ζ) *paradoxus*, pileo disco depresso nigrescenti-rubro, ceterum rubello, lamellis emarginatis!

Ceterum a *sanguinario* non differt. — Sapor dulcis foliaceus. Specimen unicum tantum lectum in consortio *Ag. emetici*.

- η) *depallens*, pileo planiusculo læviusculo fufcescente rubro disco depallescente, stipite albo l. rubro.

Ag. depallens. Pers. Syn. l. c. p. 440.

Ellr. Esb. t. 3. fig. 1, 2. a, b. Schaff.

(t. 112.) 167

Sapor in eodem specimine nunc dulcis nunc acris!

- θ) *expallens*, pileo rubro, margine albo.

- ι) *rotus albidus?*

Cum plurimis aliis, quas omnes enumerare et superfluum et ingratum foret opus. Nulli enim omnino dantur varietatum limites. Et sapor, quod jam *Persoon* l. c. aliique annotaverunt maxime in iisdem formis ludit. *Stipes* al-

bus l. rubellus maxime variat, sed lamellas constanter albas vidi, hinc, quamvis invitus, *alutaceum* separavi. Eriam huc retulissem *Ag. roseum* Pers., vulgarem in Svecia, nisi substantia, auctoritate Cel. *Link in Schrad Journ.* l. c. differet. Contextus in hac vesiculosus et in *R. emitica* flocculosus. Ceterum nulla differentia. Fatentum mihi quoque est, easdem formas eisdem locis quotannis saepe crescere, nihilo tamen minus minime separandæ; hæc enim a loco soloque omnino pendet. Observationes oculatissimorum *de Albertini* et *de Schweiniz* cum propriis omnino conveniunt.

Alia hujus subspecies est:

87. *Agaricus fallax*, pileo, depresso spadiceo margine dilutiori tuberculoso-striato, lamellis stipiteque aquose-pallidis. (*R. fallax* Fr.)

In paludibus subumbrosis inter *Sphagna*. — A priori originem ducit et color, quamvis constans. neminem fallat.

Descr. *Stipes* aquose pallidus albus glaber l. deorsum tenuissime pubescens et parum incrassatus $1\frac{1}{2}$ unc., apice crass. pennæ auferinæ l. parum crassior. *Pileus* carnosus subconcavus depressus glaber, obscure spadiceus margine pallidiori striato-tuberculoso recto l. patente. *Lamellæ* omnes simplicissimæ fragiles candidæ, sed pallidissimæ, distinctæ dentato-connexæ totæ adnatæ.

86. *Agaricus alutaceus*, pileo planiusculo rubro, margine tuberculoso lamellis integris flavis, stipite albo-rubello. (*Rusf. alutacea*. Fr.)

Ag. alutaceus. *Perf. Syn.* p. 441. α , β .
Schum. Sæll. 2. p. 244. *Albert. et Schwein.* l. c. p. 213.

Ag. Rusfala. *Scop. carn.* 2. p. 436.

Rusfala esculenta. *Perf. Obs. Myc.* I. p. 101. β .

In filvis passim.

Eisdem occurrit formis ac *Agar. Linnæi*, quare has enumerare superfluum duxi. Etiam pro hujus varietate habendus:

β . *Xanthopus*, pileo substriato, lamellis flavis, stipite dilutiori flavescente.

Ag. esculentus. *Perf. Syn.* p. 441. *Schæff. Fung.* t. 15. f. 1-3. *Krapf.* l. c. t. 5.

Rusf. esculenta. *Perf. Obs. Myc.* I. p. 101.

Cum priori.

Descri. *Agar alutacei*. *Stipes* crassiusculus, carnosus intus spongiosus sursum incrassatus, diam. $\frac{1}{2}$ unc., rectus hcurvatus 1-2 uncialis lævis glaber albus l. albus rubro variegatus, in β dilute flavescent. *Pileus* planiusculus vertice depressus ruber, rarius expallens glaber subviscosus l. ficans margine striato patente, sæpe pallidiori. *Lamellæ* juniores pallide-flavescentes de-

inde ochraceæ, plus minusve nitidæ, omnes æquales adfixæ. Sapor in hac specie sæpius dulcis solanaceus; sed tamen, licet rarius, emeticus l. stypticus.

Agaricum campanulatum P. non vidi; forte vero ob pileum campanulatum, qualem nunquam observavi, lævemque species est distincta.

Inter reliquos *Agaricos pectinaceos* Bull. etiam hujus exstat icon.

A. nitidus aliique huic forte nimis affines.

*** Pileo fulvo l. flavescente.

89. *Agaricus ochraceus*, pileo convexo vitellino margine striato, lamellis ochraceis confertis, stipite pallescente. (*R. ochracea*. Fr.)

Ag. ochraceus. *Perf. Syn.* p. 443. *Schum. Sell.* 2 p. 245.

In fagetis Sveciæ &c.

Descr. *Stipes* elongatus $1\frac{1}{2}$ - 2 unc., diam. $\frac{3}{4}$ uncial., intus carnosus solidus subæqualis l. sursum subattenuatus lævis glaber candidus in nostro fungo, nec intus medulla spongiosa fartus. *Pileus* carnosus convexus vitellinus subviscosus 2-3 unc. interdum obliquus, margine sæpius reflexo pallidiori fulcato striato. *Lamella* pallidæ ochraceæ l. flavæ æquales confertæ subtenues leviter adnatæ.

Satis diversus videtur ab *Ag. ochraceo* P. —
 Cel. *Swartz in Sv. Vet. Ac. Handl.* 1809. p. 92.
 sub hoc nomine descripsit var. β *fulcatam* *Albert.*
 et *Schwein.*, quæ differt *stipite* solidiusculo intus
 medulla spongiosa farcto, *pileo* margine lævi, nec
 striato.

Agar. ochracei var. *stipite* flavo describunt
Albert. et Schwein. l. c.

90. *Agaricus foetens*, *pileo* convexo-de-
 presso flavo, margine tuberculato-fulcato, lamellis
 subinæqualibus, *stipite*que albidis. (*R. foetens.* P.)

Ag. foetens. *Pers. Syn.* p. 443.

Ag. fastidiosus. *Pers. in Ust. Annal. Bot.*
nov. 9. p. 9.

Ag. piperatus. *Bull. Herb.* t. 292. (exclus.
Syn.)

Rusfula foetens. *Pers. Obs. Myc.* I. p. 102.

In silvis loca graminosa eligit.

Descr. Reliquis *Rusfulis* vix affinis. *Stipes*
 cavus fragilis 2 unc. crass. pollicari subæqualis
 glaber lævis albus. *Pileus* subcarnosus junior
 subconvexus depressus demum et subexplanatus
 flavus viscosus foetidus margine submembrana-
 ceus fornicatus tuberculosis fulcatus patens. *La-*
mella pall. scentes albidæ affixæ, basi venis con-
 nexæ, latissimæ aliæ dimidiatæ aliæ basi furcatæ.

Obs. Hoc de fungo parum dubius sum;

icon *Bulliardii* haud dissimilis; sed descriptio completa desideratur et *Ag. ochroleucus* P., cum quo conjungendum voluerunt *Alb.* et *Schwein.* l. c., nullo modo affinis. Ipse etiam legi varietatem *Ag. ochroleuci* Eorum *fulcatam*, quæ vero in foetentem minime transire posset.

**** Pileo obscuro purpurascente
virescente et subolivaceo.

Ex hac sectione in Svecia crescunt *R. grisea* P., *cynoxantha* Schæff., *livida* P., *nauseosa* P. *furcata* aliæque, de quibus rarioribus in secunda hujus opellæ parte.

VIII. O M P H A L I A.

+ Carnosi.

* Pileo flavescente, rufescente l. spadiceo.

91. *Agaricus cochleatus*, cæspitosus, pileo lobato-contorto tenaci, lamellis ferratis flavescentibus, stipite fulcato basi connato.

Var. *Agar. cochleati*. *Perf. Syn.* p. 450.

Abb. d. Schwämme. 3 H. t. 26. 7 6

Ag. confluens. *Sow. Fung.* t. 188.

In horto ad Femsjö in Svecia lectus.

In omnibus notis essentialibus convenit fungus noster cum Perfooniano, sed colore parum differt. E fig. citatis et descr. *Albert. et Schwein.* l. c. pag. 218 tamen hujus loci esse luculenter apparet.

Descr. *Cæspitosus* fraveolens, substantia tenaci. *Stipes* longus 4-5 unc. solidus, crassitie digiti minimi, subæqualis basi cum pluribus conatus adscendens flexuosus l. contortus, rufescenti-nigricans intus flavus firmus elasticus subfulcatus. *Pileus* carnosus firmus subconvexus umbilicatus glaber lobato-contortus flavescent (rufescenti-fuscus P.) demum rugulosus. *Caro* alba angusta tenax. *Lamellæ* latissimæ ferratæ flavescentes inæquales decurrentes.

92. *Agaricus trichæus*, pileo carnosio alutaceo sparsim squamuloso piloso umbilicato, lamellis decurrentibus stipiteque cavo luteis.

Ag. trichæus. *Perf. Syn.* p. 451.

Ad ligna pinea putrida, in silvis *Sveciæ*, *Germaniæ* &c. solitarie.

Descr. *Stipes* fistulosus uncialis, crass. pennæ gallinæ, subæqualis incurvus glaber, basi radice albido-tomentosa lignum intrante, lavis tenax elasticus. *Pileus* in nostro carnosio-membraceus 1 unc. rarissime 2 unc. sordide flavus squamuloso-pilosus convexus umbilicatus, mar-

gine subreflexo glabro. *Lamella* latiusculæ luteæ distinctæ 2-3 natæ arcuatæ decurrentes carne pilei flavescente latiores.

93. *Agaricus pachophyllus*, pileo subcarnoso cinerascenti-luteo subglabro depresso, lamellis decurrentibus albido-flavescentibus, stipite subsolido luteo.

In silvis, Julio lectus. Pulchra species.

Descr. *Ag. bello*, *trichao* &c. proximus. Solitarius. *Stipes* $1\frac{1}{2}$ unc., crass. pennæ gallinæ, glaber lævis luteus solidus apice subcavus. *Pileus* convexo-planiusculus, vertice depresso, subcarnosus subglaber l. tenuissime flocculosus $1-1\frac{1}{2}$ unc. cinerascenti-luteus, intus *cavus*! margine demum rimosus. *Lamella* crassæ distinctæ horizontaliter decurrentes albido-flavescentes.

De *Ag. rubraformi*, *nivali* &c. vid. alio loco. De speciebus Svecicis *Swartz* l. c.

** Pileo albido.

94. *Agaricus fragrans*, odoratus, pileo convexiusculo nitido lamellisque albidis l. albido-brunneis, stipite tereti concolore. *Swartz* l. c. p. 245 cum syn. et descr. optima.

In pinetis sub juniperos legi varietatem: *Stipite* $2-3\frac{1}{2}$ unc. subflexuoso, pileo toto albo margine reflexo, lamellis viridibus decurrentibus.

95. *Agaricus candicans*, inodorus albonitens, pileo carnoso subdepresso lævi, lamellis subdecurrentibus, stipite fistuloso.

Agar. candicans. Persf. Syn. p. 456. Alb. et Schwein. l. c. p. 219.

Ag. umbilicatus. Bolt. Fung. t. 17.

Inter folia dejecta faginea in *Svecia*, *Germania*.

Descr. Solitarius et raro gregarius. *Stipes* fistulosus uncialis l. parum ultra subcurvatus glaber l. basi tenuissime elongato-radicatus subpubescens, albus lævis. *Pileus* subcarnosus nitens albus haud uncial. glaberrimus lævis, junior convexus dein depressus et fere subumbilicatus. Observavi quoque pileum zonis nitidioribus variegatum! *Lamellæ* candidæ approximatae angustæ 4-natæ subdecurrentes.

Agar. dealbato. Sow. Fung. t. 123 minor, pileus integer nitens, lamellæ nunquam fordidae.

96. *Agaricus chrysoleucus*, cæspitosus, pileo carnoso-membranaceo hemisphærico albido, lamellis subdecurrentibus sulphureis, stipite albo fistuloso.

Ag. chrysoleucus. Persf. Syn. p. 457.

Ad terram nudam fungum *Svecicum* legi.

Descr. Gregarius subcæspitosus. *Stipites* plures læviter connati unciales, crass. pennæ gal-

linæ, æquales glabri albidi l. albido-pallidi, a situ sæpe curvati, juniores solidi dein fistulosi. *Pileus* carnofo-membranaceus, junior convexus deinde hemisphærico-planiusculus subumbilicatus glaber albidus l. pallescens uncialis margine striatulo. Junior interdum etiam dilutissime flavescens. *Caro* tenuissima pallida. *Lamellæ* latiusculæ 4-natæ distinctæ leviter decurrentes sulphurææ l. ochraceæ.

97. *Agaricus phyllophilus*, subcæspitosus candidus, pileo carnofo umbilicato lævi, lamellis decurrentibus, stipite fistuloso basi piloso.

Agar. phyllophilus. Persf. Syn. p. 457. Albert. et Schwein. p. 219.

In filvis abietinis *Sveciæ* legi.

Descr. Stipes cavus 2-3 unc. albidus glaber striatulus l. lævis basi subincurvus radicatus pubescens, radice subfusiformi. *Pileus* carnosus lævis umbilicatus candidus 2 unc. et ultra glaber, demum subinfundibuliformis, margine sæpius deflexo. *Lamellæ* decurrentes 4-6 natæ distinctæ. *Odor* debilis.

*** Pileo nigrescente, cinereo, livido, subviolaceo.

98. *Agaricus adustus*, magnus, pileo carnofo demum nigrescenti-adusto, lamellis albis adnatis, stipite firmo solido.

Ag. adustus. *Perf. Syn.* p. 459. *Schum. Sall.* 2. p. 309.

Ag. elephantinus. *Sowerb. Fung.* t. 36. *Bolz. Fung.* t. 28.

Ag. nigricans. *Perf. Obs. Myc.* 2. p. 50. *Bull. Herb.* t. 370. f. 2. — t. 589. (*non quadrat*)

In silvis et pratis sterilioribus, sat frequens etiam in *Suecia*, sed a Mycologis nostris neglectus.

Descri. dedit *Perfoon* in *Obs. Myc.* l. c. sed parum recedit fungus noster, hujus quamvis certissime loci. *Stipes* breviusculus 1–2 pollic. firmus 1–1½ unc. fere crassus subæqualis solidus glaber lævis albidus mox cinereo-fuligineus; intrus albidus cinerascens dein fuligineo-niger. *Pileus* carnosus 3–6 unc. convexiusculus vertice subdepresso, junior albus l. cinereus lævis mox fuscescens demum niger subtesfulatim rimosus. *Caro* pilei alba cinerascens fuligineo-nigricans compacta nec elastica absque succo rubescente; nec pileus umquam coerulescit l. fit olivaceus. *Lamellæ* albæ pallescentes demum fuligineo nigricantes crassæ fragiles adnatæ carne pilei angustiores.

Descriptiones *Ag. elephantini* melius in nostrum quadrant.

Mycogenus fungus. In hoc nascuntur *Agaricus parasiticus* Buill., *Asterophora agaricoides* Fr. (*Ag. lycoperd.* Buill. *Aster. lycop.* Dittm.) *Peziza*

mycophila Fries et plures *Sclerotii* species cum *Ag. tuberoso* &c.

99. *Agaricus expallens*, pileo carnosomembranaceo umbilicato glabro livido-cinereo margine striato, lamellis decurrentibus dilutioribus, stipite pallido.

Ag. expallens. *Perf. Syn.* p. 461.

Inter *polytricha* &c. jam æstate in *Svecia* legi.

Descr. *Agarico tardo* P. omnino nimis affinis nec ulli dantur limites, quod e descriptione cum optima *Swartzii* l. c. *Agar. tardi* comparata patebit. Solitarius l. gregarius. Stipes solidus uncialis haud ultra, crass. pennæ gallinæ l. anserinæ, elasticus lævis rectus l. basi curvatus glaber, ad radicem subfibrillosus fordide albidus l. cinerascens-pallidus. Pileus planiusculus vertice depresso margine subflexo glaber lævis tenuissime striatulus, uncialem haud attingit latitudinem, tenax carnosomembranaceus livido-cinereus fuscus. Lamellæ dilute cineræ l. fordide albido-cinerascens subdistantes tenues 4-natæ decurrentes.

100. *Agaricus dicolor*, pileo subcarnoso hemisphærico umbilicato cinereo demum albescente, lamellis subdecurrentibus dilutioribus, stipite subcompresso deorsum obscure cinereo.

Ag. dicolor. *Perf. Syn.* p. 462. *Albers.*

et Schwein. l. c. p. 221. cum plur. var., quibus addenda:

♂. *teretiufculus*, pileo subcarnoso, lamellis emarginato-decurrentibus, stipite teretiufculo.

In pinetis *Sveciæ*, gregarie et subcæspitose.

Descr. *Stipes* $1\frac{1}{2}$ –3 unc. villosopruiatus firmus, crass. pennæ anserinæ, deorsum subincrassatus villosus sæpe curvatus teres, rarius subcompressus, cinereus. *Pileus* subcarnosus umbilicatus l. infundibuliformis lævisimus glabriufculus 1–3 unc. nigricante-cinereus subnitens demum expallescent, junior sæpe convexiufculus. *Lamellæ* carne pilei tenui-cinerafcenti latiores subdistinctæ albidæ l. cinereæ 4-natæ emarginato-adnatæ; sed ob formam pilei infundibuliformem decurrentes esse videntur.

101. *Agaricus difformis*, pileo carnosomembranaceo cinereo-livido striato umbilicato, lamellis latiusculis decurrentibus albis, stipite tereti albido.

Ag. difformis. *Perf. Syn.* p. 462.

In graminosis humidis subumbrosis denso agmine. *Septembri*.

Descr. Gregarius nec cæspitosus terrestris. *Stipes* solidus dein fistulosus 2–4 uncial., crassitie pennæ gallinæ subtenuis, sed tenax, subflexuosus albidus glaber lævis æqualis. *Pileus* carnosus

membranaceus planiusculus vertice depresso, interdum excavatus et fere infundibuliformis junior elitus demum cinereo-lividus l. obscure cinerascens $\frac{1}{2}$ - 1 unc. latus glaber subirregularis striatus. *Lamellæ* carne pilei latiores adfixæ plano-decurrentes albidæ demum dilute cinerascens.

102. *Agaricus epichysium*, cinereo-nigricans, pileo submembranaceo infundibuliformi striato, lamellis distinctis decurrentibus dilutioribus, stipite basi subvillosa.

Ag. epichysium. *Perf. Syn.* p. 462. *Ic. pict.* t. 13. f. 1.

β . *icmadophilus*, tenerrimus, pileo obscure cinereo, lamellis plano-decurrentibus cinereis. *Perf.* l. c. p. 463.

In truncis putridis præsertim fagineis in *Svecia*, *Germania* &c. β . in paludibus inter *sphagna*. — Reliquas ab *Albert. et Schwein.* l. c. allatas varietates non apud nos vidi.

Descr. *Stipes* uncialis 1 - $1\frac{1}{3}$ lin. crassus subfirmus æqualis teres basi pubescens, ceterum glaber lævis rectus solidus cinereus. *Pileus* tenuissime carnosus (forte potius dicendus membranaceus) infundibuliformis cinereus l. cinereo-fuligineus, exsiccatus canus, glaber $\frac{1}{2}$ unc., margine reflexo striato. *Lamellæ* decurrentes cinereæ, interdum albidæ, 2-natæ distinctæ subangustata, carne pilei latiores. *Caro* fuscescens.

+ Pileo membranaceo.

* Lamellis adnatis, vix decurrentibus.

103. *Agaricus ocellatus*, pileo plano lævi albo, umbilico obscuriori, lamellis adnatis albis, stipite fistuloso fuscescente.

In pinetis juxta vias locis graminosis. Ad Femsjö in Svecia mihi obuius.

Descr. Stipes fistulosus vix uncialis, crass. culmi graminei minoris, æqualis erectus lævis glaber exalbido-pallide-fuscescens, basi radícula sublonga villosa appendiculatus. Pileus membranaceus planus, vertice depresso obscuriori, glaber tenax $\frac{3}{4}$ - $\frac{1}{2}$ uncial. albidus lævis, margine patente crenulato-striatulo. Lamellæ albæ basi latiores alternæ totæ adnatæ sed non decurrentes et a stipite facile decedunt.

Habitu quodam communi distinguitur; apud recentiores Mycologos non invenio.

104. *Agaricus pictus*, pileo ovali-oblongo nigro-striato fuscescente umbilico discolori, lamellis adnatis albidis, stipite solido spadiceo fusco.

Ag. glandifer, pileo oblongo nigro-fusco striato: margine constricto-albo, stipite lineari nigro-fusco, lamellis albis elongatis. *Batsch. El. fung. Cont. 1. p. 99. t. 18. f. 86. (minime tamen cum Ag. glandifero Pers. confundendus.)*

In silva *Femfjönenski* locis umbrosis supra folia *Vaccin. Myrtilli* 2 l. 3 specimina rarissimi hujus fungi et optime distincti mihi obvia. Unicum tantum vidit Cel. *Bartsch*.

Descr. *Stipes* solidus 2-3 unc. subfiliformis æqualis, vel sursum tenuissime incrassatus strictus lævis glaber nigro-fuscus l. spadiceus apice dilutior basi membranula fusco-strigoso-radiata impositus. Specimen unum vidi stipite valde flexuoso. *Pileus* membranaceus umbilicatus ovali-oblongus subcuculatus $2\frac{1}{2}$ lin. altus 2 lin. latus fusco-spadiceus, junior dilutior, demum nigrescenti-fuscus, umbilico depresso albo-flavescenti (saltem in nostro), striis primum olivaceis mox nigris pictus, margine sæpe constrictus dilutior. *Lamellæ* distinctæ subæquales breves latæ albidæ dein subgriseæ adnato-decurrentes.

Ad hanc sectionem pertinent: *Ag. androsaceus*, *epiphyllus* &c.

** Lamellis longissime decurrentibus.

105. *Agaricus roridus*, pileo hemisphærico-conico sublavi albo, umbilico concolori, lamellis arcuato-decurrentibus, stipiteque copiosissime candido-viscosis.

In pinetis, locis præcipue umbrosis roridis, in ramis *Vaccin. Myrtilli*, furculis *Lycopodii an-*

norini, foliis *fagineis*, *pineis* &c. solitarie, sed minus infrequens in *Suecia*.

Descr. Alia iterum species ab omnibus distinctissima, ut ne ullæ quidem dentur affines. Solitarius. *Stipes* solidus $\frac{1}{2}$ - 1 unc. albus æqualis subtenax adhærens, *visco copiosissimo nigrissimo!* corticatus. *Pileus* junior conicus demum hemisphæricus umbilicatus glaber fuscus sublævis albus 2 - 3 lin. latus. *Lamellæ* latiusculæ candidæ 2-natæ arcuato-decurrentes.

106. *Agaricus stellatus*, pileo convexo striato albo, umbilico concolore, lamellis decurrentibus, stipite sicco membranula convexa strigoso-radiata imposito.

Ag. stellatus, pileo striato, stipite gracili involuero radicali stellata cincto. *Hoffm. Nomencl. fung.* I. n. 306. t. 6. f. 2.?

Ad truncos putridos humidos in *Suecia* rarior.

Descr. *Ag. stylobati*, lacteo β . epiphyllis γ . aliisque ob stipitem basi tuberculatum affinis. Distinctus tamen videtur. Totus albus gregarius. *Stipes* subsolidus vix uncial. æqualis, crass. pennæ passerinæ minoris, subgracilis fragilis curvato-ascendens (a loco), basi membranula convexa concolore 1 - 2 lin. strigoso-radiata. Interdum membranulæ duo confluant. *Pileus* membranaceus convexus umbilicatus diaphanus glaber striatus

sub 4 lin. ficcatus interdum flavescent. *Lamellæ* candidæ tenues integræ l. alternæ distinctæ decurrentes. (In *stylobati subliberæ*!)

An figura citata ad hanc l. reliquos affines pertineat species, dubius sum. Huic tamen proximus.

Ag. stellatus. Retz. in *Vet. Acad. Handl.* 1769 p. 252 est *Ag. corticola*. Pers. Swartz.

Ad hunc etiam pertinent *Ag. corticalis*, Som. t. 243, et *Ag. hyemalis* Osbeck in Retz. *Suppl.* 1. *Fl. Scand.* p. 19.

107. *Agaricus hepaticus*, pileo convexulo profunde umbilicato lævi lamellisque decurrentibus hepaticis, stipite fistuloso deorsum subincrassato subpubescente.

Ag. hepaticus. Barsch. *El. fung.* Cont. 2. t. 38. f. 211.

In terra nuda *Sveciæ*, *Germaniæ*.

Descr. Non *Agar. montani* varietas, ut vult Cel. *Persoon*; sed *Agar. ericetorum* me iudice proximus. *Stipes* 1 - 1½ unc., 2 lin. crass. fistulosus subadscendens lævis dilutior hepatico-rufescens glaber, basi subincrassatus albidus tenuissime pubescens. *Pileus* convexus submembranaceus (parum carnosus) ½ unc. subglaber l. nonnunquam fibrillosus rufo-hepaticus, umbilico profundo concolore, margine subinflexo. *Lamellæ* latiusculæ rectæ 2-natæ albido-hepaticæ arcuato-decurrentes.

Ag. ferrula P. etiam huic affinis videtur, an potius ad carnosas species referendus?

108. *Agaricus ericetorum*, pileo plano umbilicato striato fordide flavescente l. griseo, lamellis decurrentibus, stipite deorsum pubescente subfistuloso.

Ag. ericetorum. *Albert. et Schwein. l. c. p. 225.*

Ag. nothus, albidus cinerascens flavicansve, lamellis decurrentibus, carne alba, stipite gracili farcto. *Gmel. Syst. Linn. 2, p. 1423. (diag. bona, coloris mutationes exprimens.)*

Agar. pseudo-androsaceus. *Bull. Champ. t. 276. (optime.)*

α) *albicans l. flavescens.* *Alb. l. c. p. 225. α . G. Dan. t. 1015. f. 1. (exclus. syn.)*

β) *griseus.* *Alb. et Schwein. l. c. 77.*

Ag. ericetorum. *Pers. Syn. p. 472. Obs. Myc. 1. p. 50. t. 4. f. 12. cum descr.*

In ericetosis humidis α) copiosissime, β) rarissime in *Svecia* legi.

Descr. α) *Stipes* subfistulosus (sed haud raro solidus) $\frac{1}{2}$ –1 unc. subæqualis, crassit. pennæ gallinæ et ultra, teres firmus glaber lævis pallidus albus l. exalbido-flavescens, basi pilis tenuissimis, interdum nullis, obsitus. *Pileus* membranaceus (l. disco carne tenuissima albida) $\frac{1}{3}$ – $\frac{3}{4}$ unc. flavescens pallidus subfordidus planus umbilicatus striatus, margine patente, in speciminibus

luxuriantibus undulato-lobato, radiis obscurioribus picto. *Lamellæ* pallidæ latissimæ basi crassiores (*male, me iudice, dicitur: basi latiores*) 2-natæ pallidæ decurrentes distantes. — In pratis humidis legi formam: *stipite* solido albido glabro $\frac{1}{2}$ — 1 unc. subflavescenti, *pileo* subcarnoso infundibuliformi, turbinato flavescenti-albido sublævigato. *Lamellis* 4-natis.

109. *Agaricus campanella*, subæspitosus, pileo hemisphærico umbilicato ferrugineo, lamellis decurrentibus, stipite fistuloso spadiceo æquali.

Ag. campanella. *Perf. Syn.* p. 469. *Batsch. El.* p. 74.

Ag. fragilis. *Schaff. Fung.* t. 230.

Ad truncos pineos, frequens. *J. Floris Suecicis* neglectus tamen.

Descr. Gregarius subæspitosus. *Stipes* fistulosus 2 unc., crassit. vix pennæ gallinæ, glaber subæqualis elasticus spadiceus l. castaneus sursum dilutior, radice fibrillosa pubescente subcurvata. *Pileus* membranaceus hemisphæricus glaber striatus tenax umbilicatus $\frac{1}{4}$ — $\frac{2}{3}$ unc. flavo-ferrugineus. *Lamellæ* decurrentes dilutiores flavescentes subalternæ basi venis inter se connexæ.

110. *Agaricus tricolor*, gregarius, pileo convexo umbilicato ochraceo-flavo, lamellis decurrentibus basi crassis, stipite solido sursum incrassato.

Agar. tricolor. *Albert. et Schwein. l. c.*
p. 224. t. 9. f. 5.

Habitat in silvis, pascuis &c. Germaniæ,
Sveciæ.

Obs. Fungus Svecicus ad variet. β . *bicolor* propius accedit. Ceterum vix differt nisi lamellis flavescentibus aurantiacis et vix roseis. A priori, cui habitu haud dissimilis, abunde distinguitur: *stipite* solido sursum incrassato, *pileo* minus striato dilutiori et *lamellis* basi crassis. — Non *Ag. tricolor* Bolt.

III. *Agaricus muralis*, subgregarius, pileo convexo umbilicato striatulo rufescente, lamellis decurrentibus, stipite solido fuscescente-rufo.

Ag. muralis. *Swartz in Vet. Acad. Handl.*
1809. p. 94. *Sowerb. t. 322.*

Meum fungum in palustribus legi.

Recedit a descr. *Swarziania*: *stipite* unciali sursum incrassato, *pileo* minus umbilicato et lamellis flavis. Varietas saltem esse videtur.

III. *Agaricus (fibula. P.) γ . hydrinus*, lamellis flavescentibus, *major*.

In palustribus.

Color pilei vulgaris hujus fungi ludit quam maxime, aurantiaco, albido, albo et fulvo variegato, rufescente &c. (vid. *Bull. Champ. t. 550*) sed lamellæ sæpius albæ. In sylva deusta paludosa legimus varietatem nunc describendam, cujus *Stipes* brevis vix uncialis teres junior

solidiusculus deinde fistulosus (hinc in *communi* etiam solidus ab his, fistulosus ab illis describitur) laevis glaber l. deorsum pubescens pileo concolor. *Pileus* subinfundibuliformis membranaceus $\frac{2}{3}$ - $1\frac{1}{4}$ unc. latus! ochraceus l. aquose flavus radiis obscurioribus ramosis striatus, margine deflexo nonnunquam undulato-lobatus. *Lamella* dilutiores flavæ latiusculæ 4-natæ distinctæ decurrentes.

113. *Agaricus Swartzii*, pileo planiusculo umbilicato laevigato albido, disco fusco, lamellis longissime decurrentibus albis, stipite solido.

In fagetis umbrosis roridis, *Svecia* inter muscos, rarior.

Vere distinctus et stipite solido aliisque notis haud difficile a sequenti distinguitur.

Descr. *Stipes* solidus glaber laevis rectus uncialis deorsum subangustatus subfirmus fuscescens albidus. *Pileus* planiusculus junior convexus umbilicatus 2 lin. latus (haud ultra) laevis albidus vertice fusco. *Lamella* longissime decurrentes albæ tenuisculæ in stipite angustiores.

Icones harum alio loco dabo.

114. *Agaricus speireus*, pileo planiusculo umbilicato striato cinerascenti, disco fusco, lamellis longissime decurrentibus albis, stipite fistuloso.

In truncis quercuum dejectis muscosis in *Svecia*, sparsim crescens.

Species illæ *Swartzii*, *speireus* et *ocellatus* tribum naturalem, quem *discoideos* dixi efficiunt.

Descr. Solitarius rarius subgregarius. *Stipes* uncialis et ultra inter muscos sæpe radicans fistulosus rectus lævis glaber gracilis tenax filiformis albidus. *Pileus* membranaceus planiusculus umbilicatus glaber striatus pallide cinereus vertice fusco striisque obscurioribus variegato, 2-3 lin. latus haud ultra. *Lamellæ* candidæ alternæ basi latiusculæ distinctissimæ longissime decurrentes.

115. *Agaricus* (o?) *ditopus*, pileo subcarnoso-coriaceo-membranaceo undulato-lobato cinerascens umbilicato, lamellis adnato-decurrentibus stipiteque cavo concoloribus.

An varietas Agar. buccinalis. *Batsch. El. fung.* Cont. 2. p. 25. t. 39. f. 214.

In filvis abietinis densis *Sveciæ* inter muscos. De tribu hujus fungi dubius sum, *pileus* fere infundibuliformis hujus loci esse svidet, sed excentricus sæpe est, et substantia fere coriacea ad *Pleuropodes* accedit.

Descr. *Stipes* cavus 2-uncial. crassitie pennæ anserinæ crassiorve teres l. compressus cinerascens glaber basi pubescens. *Pileus* difformis carnosus-coriaceus undulato-lobatus subexcentricus subinfundibuliformis cinereus glaber 1-2 unc. et ultra. *Lamellæ* tenues polynatæ cinerascens adnato-decurrentes in lobis pilei divergentes.

Transitum ad sequentem familiam efficit.

IX. P L E U R O P U S.

* Pileo integro, stipite excentrico.

116. *Agaricus corticatus*, pileo carnoso convexo floccoso-villoso cinerascens, margine subinvoluta, lamellis decurrentibus candidis, stipitibus excentricis albo cortinatis.

In truncis fagineis adhuc vivis *Sueciæ*.

Species maxime notabilis. *Fungos* tamen juniores tantummodo vidi, hinc magnam cum *Ag. Lepiota Alb. et Schwein.* l. c. p. 229 suspicor affinitatem. Cortina alba, stipite non verticali aliisque notis differre videtur. An varietas?

Descr. Subsolitarius. *Stipes* solidus 2 unc., $\frac{3}{4}$ unc. crassus subæqualis firmus adscendens fibrilloso-villosus l. g'aber. *Pileus* carnosus convexus cinerascens floccoso-villosus 2 unc. margine subinvoluta, *cortina* alba lamellas nondum explicatas obtegente appendiculatus, excentricus, sed integer. *Lamellæ* candidæ longissime decurrentes inæquales subangustæ basi tenuiores sæpe dichotomæ subdistinctæ.

117. *Agaricus ulmarius*, imbricato-gregarius, pileo exalbido-lutescente laticissimus, lamellis latis subemarginato-adnexus, stipite excentrico nudo.

Ag. ulmarius. *Perf. Syn.* p. 473. *Sowerb. Fung.* t. 67. *Bull. Herb.* t. 510. *Sibth. Ox.* p. 345.

β. epiphegus, subsolitarius, pileo albido-livescente, lamellis truncato-adnexis.

In truncis. *α.* in truncis *Populi nigrae* ad Lundinum. *β.* ad *Fagos* putridas legimus.

Var. *α.* optime descripsit Illustr. *Swartz* l. c. 1808. p. 251 in eademque arbore legit.

Descr. *β.* Minor gregarius, 2-3 basi connati. *Stipes* adscendens curvatus teretiusculus deorsum subincrassatus solidus fere glaber 1-2 unc. elasticus. *Pilei* saepe imbricati, sed pauciores quam in vulgari, integri excentrici horizontales convexi margine deflexo involuto 1-2 unc. albido-cinereascens. *Caro* alba. *Lamellae* albæ 2-4 natæ latæ truncato-adnatæ. — Species non est distincta.

Obs. Distinguendum inter lamellas rotundato-adnexas, truncato-adnexas et emarginato-adnexas, quæ optimos saepe præbent characteres.

118. *Agaricus spodoleucus*, subsolitarius, pileo convexo cinereo glabro, lamellis angustis decurrentibus albis, stipite excentrico nudo.

Ad truncos fagineos *Sveciæ*.

Primo obtuitu *Ag. ulmario β. epiphego* similis, ut pro varietate omnino haberem, nisi lamellis longissime decurrentibus distingueretur.

Descr. Solitarius l. gregarius, vix caespitosus. Odor nullus. *Stipes* brevissimus vix $\frac{1}{2}$ unc. crassus (penna anserina saepius crassior)

solidus elasticus adscendens glaber lævis cinerascens l. pallidus. *Pileus* convexus carnosus uncialis et ultra glaber cinereus l. cinerascenti-fuscus integer; sed excentricus, margine deflexo, initio inflexo. *Lamellæ* albæ l. pallescentes decurrentes angustæ 2-4 natæ carne pilei alba angustiores confertæ eglandulosæ; quamvis *Agar. glanduloso Bull.* et præcipue ejus varietate *β. horizontali Albert.* affinis.

119. *Agaricus torulosus*, carneo-subsuberosus, pileo infundibuliformi margine revolutosubobliquo fulvo-carneo, lamellis longe decurrentibus, stipite brevi griseo-tomentoso. *Pers. Syn.* p. 475. *Albert. et Schwein.* l. c. p. 227.

Ag. griseo-tomentosus. *Batsch. El. fung.* f. 33.

Ab hoc fungo sequentem varietatem separare non ausus sum:

β. tephrotricus, pileo planiusculo cinereo-alutaceo stipiteque griseo-tomentosis.

Ad truncos destructos *Sveciæ.*

Descr. Gregarius. *Stipes* brevis vix unciam attingens crassiusculus deorsum subincrassatus solidus curvatus albidus griseo-tomentosus. *Pileus* planiusculus carnosus molliusculus, integer; sed valde excentricus subobliquus alutaceo-cinereus griseo-tomentosus 3-5 unc. forma sæpius subrotunda l. obovata, margine subinflexo.

Lamellæ candidæ longissime decurrentes subdistinctæ 4-natæ margine integerrimo.

Conf. *Albert. et Schwein. l. c.*

**** Pileo dimidiato, stipite marginali, subnullo.**

120. *Agaricus atro-coeruleus*, mollis subacaulis pileo subvillosa secco atro-coeruleo intus gelatinoso, lamellis albidis.

Ag. molli. *Pers. (sæpe a me lecto) affinis, sed differt colore, odore, pileo villosa &c.*

In trunco fagineo putridissimo loco umbroso ad terram dejecto.

Descr. Subgregarius acaulis, immo caespitosus. *Pilei* horizontales l. parum adscendentes 1-2 uncial. subobovati basi angustati ficei rugulosi villosi, rarius glabri coeruleo-nigricantes molles, margine integro sæpe inflexo. Odor debilis haud ingratus. *Substantia* interna mollis, superne stratum crassum (2-3 lin. et ultra) gelatinosum (ut in volva *Pballi impudici*) fusco-nigricantem ferens, carne pilei albida tenuissima facillime dissepabili crassior. *Lamellæ* ad basin decurrentes 4-8 natæ albidæ demum subflavescentes carne pilei latiores firmæ approximatae margine integerrimo.

Substantia interna gelatinosa omnino paradoxa species. Cum *Perfoonianis*, si a *molli* dis-

cesferis, nihil habet commune, sed affinis videtur
Agaricus. Schæff. Fung. Bav. t. 246.

Legi varietatem odore fortiori gratissimo.

121. *Agaricus flabelliformis*, subsesilis, pileo multiformi cervino subvillosa, lamellis pallidis ferratis, stipite subnullo.

Agar. flabelliformis. Pers. Syn. p. 479.
 Bolt. Fung. t. 157 (minus bona).

In Fago cava ad Femsjö quot annis viget.

Deser. Fungus sesilis cæspitosus imbricatus. Pileus margine stipitiformi elongatus carnosus adscendens l. horizontalis sæpe reniformis 2-4 uncial. difformis subconvexus colore vario, plus minusve rufescens, hepaticus fusco-spadiceus l. castaneo-flavescens adultus subtomentosus, tomento erecto breviusculo concolore denso, et interdum farina alba conspersus, margine glabro dilutiori subcrenato. Lamellæ ad radicem decurrentes divergentes latisimæ (carne pilei 4-plo latiores) albidæ ferrato-dentatæ longitudine inæquales.

An tomentum parasiticum? — *Agarici pleuropodes*, præsertim dimidiatæ, maxime variant pilei forma, colore &c. ex. gr. *Ag. salignus*, *ostreatus*, *mollis*, *nidulans*, *variabilis*, *epigæus* aliique, hinc difficile firmis distinguuntur characteribus.

MERULIUS.

Cantharellus.

122. *Merulius rubiformis*, gregarius sub-
cæspitosus, pileo membranaceo umbilicato ru-
goso-squamuloso, plicis rectis cinerascenti fla-
vis, stipite flavo deorsum incrassato.

M. tubiformis. *Perf. Syn.* p. 489. *Comm.*
Schæff. p. 62. *cum. syn.*

Mer. helvelloides. *Gmel. Syst. Linn.* II,
2. p. 1430.

Mer. villosus, cæspitosus, pileo umbili-
cato squamoso-villoso pallescente-fuligineo,
plicis distantibus cinereo-pallidis, stipite cavo
deorsum incrassato lutescente. *Perf. ic et descr.*
p. 17. t. 6. f. 1. (*me iudice, non differt.*)

Helvella tubæformis. *Bull. Herb.* t. 461.
f. A. C. (*optime*).

Peziza undulata. *Bolt. Fung.* t. 105. f. 1.

β. *epixylus*, pileo turbinato, stipite solido.

In silvis umbrosis subhumidis et fagetis
et pinetis, vulgaris etiam in *Suecia*.

Descr. Maxime variabilis. Gregarius l.
cæspitosus. *Stripes* sæpius cavus, sed etiam, præ-
cipue in β, solidus 2—3 unc. glaber lævis
nunc teres nunc compressus deorsum sæpius
incrassatus rectus l. adscendens subfirmus
elasticus, semper late flavus. *Pileus* membra-

naceus, sed in β subcarnosus, $\frac{1}{4}$ —2 unc. subrugoso-fibrillosus squamulosus, sed etiam glaber, convexus umbilicatus l. totus infundibuliformis demum pervius flavescenti-cinereus l. cinereo-spadiceus, margine subundulato crispo irregulari. *Venæ* flavo-cinerascentes, sæpius dilutius flavescentes ac pileus, sed etiam cinereæ dichotomæ distantes venis connexæ subdecurrentes.

Quibus notis differt *M. fuliginus* P. me fugit.

123. *Merulius cinereus* cæspitosus, pileo subinfundibuliformi squamuloso nigrescente, plicis crassis glauco-cinereis, stipite cavo nigrescente.

M. cinereus. Pers. Syn. p. 450. Ic. et descr. p. 10. t. 3. f. 3.

Helvella hydropipes. Bull. H. t. 465. f. 2.

In silva (*Hallanäsensi*) prope Femsö in Suecia adhuc tantum lectus.

Descr. *Mer. cornucopioidi* maxime affinis, sed hymenio valde plicato aliterque colorato differt.

Reliquas hujus sectionis a me intra Patriæ fines lectas species optime in *Vet. Ac. Hand.* 1809 descripsit Celeb. Prof. *Swartz*.

*** *Stereopa* (pileo dimidiato verticali
 obs. ** *Gomphus*).

124. *Merulius crucibulum*, pileo verticali cupulæformi carnosio alboflavescenti, intus venis luteis ramosis.

In hymenio *Boleti igniari* et ligno putrido.

Descr. Gregarius sessilis carnosus, ubique radiculis byssoides, cupularis (*Cyat. crucibulum* haud male referens) 4—6 lin. et ultra, extus albido-flavescent subglaber, margine inflexo acuto, intus aurantiacus venosus, venis dichotome ramosis crispatis flavis, ejusdem coloris et indolis ac in *M. aurantiaco* Pers.

Obs. Caute distinguendus ab *Merul retirugo*, *bryophilo*, *crispo* &c. quos etiam cupulæformes vidi; sed ab omnibus notis speciebus hujus sectionis colore etc. venarum differt.

125. *Merulius tremellosus*, acaulis imbricatus l. resupinatus, pileo effuso reflexo tomentoso albo, plicis poriformibus acutis rubicundis.

Merulius tremellosus. Pers. Syn. p. 496.

Obs. myc. 2. p. 92. Schrad. spic. p. 139. Schum.

Sall. 2. p. 372. Albert. et Schwein. p. 238.

Merulius. Hall. hist. n. 2329. *betulinus* Gmel.

Agaricus betulinus. Wulf. — Fl. Dan. t. 776.

(omnino hujus loci.)

Ag. cantharellus. *Batsch. El. p. 95. f. 34.*

In truncis *Suecicis* primus, sed frequenter, legi. Apud exteros vulgatissimus videtur.

Tot locis descriptus ut superflua foret descriptio nostra.

**** *Serpula.*

126. *Merulius vastator*, orbicularis aureus planus margine subbyssoideo albo, venis gyrosis crispis.

M. vastator. Pers. syn. p. 497. Tode in Abh. Hall. Naturf. l. p. 351. t. 2. f. 1—4. Schum. Sall. p. 371. Albert et Schwein. l. c. p. 236.

β . *epiphyllus*, tenuissimus glaber dilutior.

α . Ad ligna pinea putrida in silvis. β . in foliis *Syringæ vulgaris* loco umbroso notabilem hanc legi varietatem.

Obs. Odor fungi noster subnullus et tubercula caulescentia rarissima.

Ut Novitiæ Floræ Suecicæ etiam nominandi *M. serpens* Tode et *M. rufus* P., hunc hyeme ad truncos fagineos, illum æstate ad pineos observavi.

127. *Merulius fugax*, resupinatus tenuis candidus, venis prominulis ad tactum evanescentibus.

In ligno pineo putrido.

Descr. B.(P.) *reticulato* affinis, sed venis omnino distinctus. Effusus tenuissimus subglaber, hymenio difformi reticulato l. reticulato-venoso. Adeo tenuis, ut ad tactum evanescat.

***** *Mesenterica* (gelatinosa, venis subdistantibus.)

128. *Obs.* *Mesenterica* genus ad fungos byssoideos s. e. filis compositos referre non potui et cum a *Meruliis* nullo differunt caractere cum his conjungi. *Mer. tremellofi* substantia etiam est tremellofa l. gelatinosa, fugacitate et venis subdistantibus, cum *M. fugaci* aliisque conveniunt. — Vide *Schrad. spic. p. 142*. Sequentes hujus subgeneris species mihi notæ:

- 1) *Merulius (mes.) aureus*; citrinus l. aureus.

Mesent. lutea, *Pers. Syn. p. 706*. *Albertini et Schwein. l. c. p. 374*.

Mes. tremelloides, *Tod. fung. Meckl. 1. p. 7. t. 2. f. 12*.

- 2) *Merulius coeruleus*, *coeruleus*.

Mesent. coerulea, *Pers. l. c.* — *Tod. l. c. β*.

- 3) *Merulius erysibe*, æruginosus, venis reticulatis.

In trunco fagineo.

- 4) *Merulius griseus*, griseus, venis ramosis-
sissimis. — *Albert. l. c.*

Eodem ac antecedens loco meum fungum legi.

- 5) *Merulius sanguinolentus*, lacteus, ad tactum sanguinolentus.

Mes. sanguinolenta. *Alb. l. c.*

- 6) *Merulius argenteus*, albus immutabilis.
— *Pers. Syn. p. 706. Vaill. par. t. 8. f. 1.*

Ex his *M. coeruleum* et *sanguinolentum* non vidi. — Quam maxime tamen dubito, an species sint distinctæ. Si igitur quis has sub nomine *M. mesenterici* conjungeret, non multum forte erraret.

SCHIZOPHYLLUS. FRIES.

Sporangium pileiforme sessile subtus lamellis longitudinaliter bifididis, margine revolutis, thecis superficialibus, spora subeffusa (*in rima longitudinali*).

129. *Schizophyllum communis*. Agaricus alneus. Linn. et omn. fere Auct. — Sow. t. 183. Bull. ch. t. 346. 581.

Ag. multifidus. Batsh El. fung. f. 126.

Merulius alneus. Gmel. Syst. Linn. II. 2. p. 1431. Schum. Sall. p. 370.

In truncis fagineis, alneis, tiliaceis, betulinis et populneis legi. Singularis hic fungus, nunc ad *Agaricos* nunc ad *Merulios* relatus, et ab omnibus Mycologis male intellectus, distincti esse generis. nullus dubito. Lamellæ fissæ, quæ *Valvellæ* dici possent, margine revolutæ et villosæ illum a veris *Agaricis* et *Meruliis* abunde distinguunt; sed fructificationis processus etiam tam longe recedit, ut ab his non possit non separari. Hujus iconem ad aliud tempus reservare cogor.

Obs. Ut *Sistotrema* ab *Hydno*, *Schizophyllum* ab *Agarico* differt.

DÆDALEA. PERS.

130. *Dædalea quercina*. Pers. Boll. t. 63.
hujus legimus varietatem.

γ. *nigricantem*, extus nigram, intus atrofuscentem.

Agaricus dædaleis finibus excavatus nigricans. Batt. fung. arim. p. 72.

In truncis quercinis. — Specie non differt ut in Synopsi suspicatus Cel. Persoon.

131. *Dædalea betulina*, coriacea zonata tomentosa pallescens, sinulis rectiusculis, laminis ramosis.

D. betulina. Rehent. fl. neom. p. 371.

Agaricus betulinus. Linn. Fl. Suec. p. 1244.
Sow. fung. t. 182. Batt. t. 30. f. D.

A. coriaceus. Pers. Syn. p. 486. Bolt. F. t. 158. Bull. t. 394.

A. versicolor. Plan. Erf. p. 4.

In truncis betulinis et quercinis.

Descr. Gregarius, raro imbricatus. Pileus durus firmus tuberoso-coriaceus dilatato reniformis subcinerascens planus l. convexus zonatus tomentosus, zonis magis elevatis villosis aliis depressis glabriusculis. Sinuli recti, laminis lignoso-coriaceis sæpius ramofoconnatis difformibus interdum rectis simpli-

cibus 4 natis, albide-pallidis. Ætate fufcescit et nomumquam virescit ut *D. gibbosa*.

Synonymon *Linnaei* ad hanc speciem sine dubio pertinet.

132. *Dædalea sepiaria*. Swartz l. c. 1810. p. 2. cum syn. et descr.

Synonymis addendum.

Agaricus acaulis utrinque planiusculus. *Linn. Fl. Suec. n. 1243.* (Errant igitur qui hunc pro *A. betulino* L. habent).

Ag. squamosus. *Liliebl. Sw. Fl. 2.*

Merulius squamosus. *Schrad. in Gmel. Syst. et in spic. p. 140.* (ubi et apud *Wulf. l. c.* plurima synonyma).

M. sepiarius, Wahlenb. Fl. lapp.

133. *Dædalea cinerea*, imbricato-connata sububerofa tomentosa cinerea, sinulis variis flexuosissimo-sinuatis l. periformibus, laminis lignosis.

Agaricus squamosus dædaleis sinibus cinereus. *Batt. Fung. t. 38. f. G. optime.*

In truncis fagineis muscosis; ad Femsiö in Suecia.

Descr. Pilei acaules cæspitofi imbricato-connati numerosissimi juniores interdum fere resupinati demum subpulvinati uberofi duri; sed non crassi, subelastici flexibiles 1½—3unci-

ales subundulati plus minusve cinerascens, etiam ferruginescentes et virescentes, villosi zonati, zonis latis elevatis paucioribus, margine patente concolore l. spadiceo-cinereo, subtus planus demum convexus. *Sinuli* valde diffformes, in junioribus pauci poriformes subrotundi, demum longissimi maxime divaricati flexuosi angustissimi albidii. l. pallescens, detriti fuscescentes; etiam cretaceo-albivisi. *Lamini* non distincti crassi lignosi minime flexibiles, ut in priori.

Optimam hujus generis ideam præbet, an satis differt *D. confragosa*, Pers.?

134. *Dædalea gibbosa*, gregaria suberosa subtomentosa albida, basi gibboso-porrecto, sinulis rectiusculis.

Dædalea gibbosa. Pers. *syn.* p. 501. *Schum.* *Sall.* p. 373. *Rebentisch Fl. neom.* p. 372.

Merulius gibbosus. Pers. *obs. myc.* 1. p. 21.

Boletus suberosus. Linn. *Suec.* p. 453.

Bol. sinuosus. Sowerb. *f. t.* 194. (*sec Rebent.*)

In truncis fagineis meum fungum frequenter legi.

Descr. Gregarius vix cæspitoso-imbricatus. *Pileus* suberosus albidus, demum pulvere viridi conspersus, pulvinatus elasticus subtomentosus interdum zonatus magnitudine varia

(2—4 unc. latús) basi sæpe gibboso-porrecto, margine obtuso sæpius obscuriori cinereo; pileus demum cinerascit. Caro alba tenacissima. Sinuli oblongi recti, alii subrotundi albi. Synonymon *Linnaei* huc certe pertinet.

135. *Dædalea mollis*, gregaria carnosa rugosa incarnata, sinulis variis flexuoso-elongatis et subrotundis albidioribus.

Boletus mollis. *Perf. syn.* p. 526. *Obs. myc.* 1. p. 22. *Albert et Schwein.* l. c. p. 247. In truncis pineis *Sueciæ*, Germaniæ.

Descr. Gregarius l. subcæspitose-imbriatus. Pileus carnosus mollis dilatatus (1—6 Lin.) crassiusculus triqueter incarnatus rugosus l. subvillosus-strigosus, etiam glaber, margine acuto, basi effuso-dilatato. In adultis zonam erecto-strigosam nonnumquam observavi. Sinuli perangusti alii poriformes pluri demum longissimi perangusti flexuosi albidii l. pallide incarnati a pilei substantia.

136. *Dædalea albilla*, gregaria suberosa subglabra alba, sinulis angulatis subelongatis concoloribus.

Agaricus dædaleis sinibus minoribus. *Batt. fung. ar.* t. 38. F. E. F.

Ad truncos betulinos *Sueciæ* legit *Forsander*.

Descr. Acaulis gregaria rariùs imbricata. *Pileus* dilatatus subreniformis convexus albus glaber subrugulosus, junior albida sericea hirsutiae obtectus, suberoso-coriaceus azonus l. zonis elevatis concoloribus, subtus concaviusculus. *Caro* tenuissima suberosa concolor. *Simili* juniores periformes angulati, dein elongati suberosi firmi recti, marginem versus poriformes, albi, e pilei substantia formati.

137. *Dædalea heteromorpha*, resupinata, sinulis irregularibus pallidis flexuosis, margine byssino albido.

Ad ligna pinea putrida *Sueciæ*.

Mera omnino *Dædaleæ* species, nec *Poriæ*, quamvis habitus diversus.

Descr. Resupinatus effusus latisimus, subiculo subobsoleto, margine irregulari byssino albicante. *Simili* suboliqui longissimi flexuosi connati nonnulli poriformes maxime irregulares, non vero dentato-laceri, ut *Sistotrema* affinis esset, molliusculi pallefcetes, tacti flavescenti-cinerascentes l. subrubiginosi, *tubis* obliquis breviusculis persistentes.

BOLETUS. Fries.

Suillus. Mich. Gen. p. 126. t. 6. f. G.

O. Pori integri toti e substantia propria, a pilei diversa. et libera, formati.

Habitus: *Stripitati carnosì horizontales, tubis elongatis a pileo facile disjunctis.*

Obs. Genus hoc, a *Michelio*, alio licet sub nomine, constitutum, maxime naturale, et omnia Genera naturalia sunt optima et servanda, præcipue cum characteribus optimis distingvitur tantique momenti, quanti apud Lichenes, inter quos ad Classes (*Ach. syn.*), l. si malles familias, distinguendas valent.

* Stipite annulato.

138. *Bolerus luteus*, annulatus, pileo pulvinato viscoso subcastaneo-livido l. flavescente, poris subrotundis luteis, stipite albo crassiusculo.

B. luteus. Linn. Fl. Suec. p. 452. Schrad. spic. p. 142.

β. *virgatus*, pileo rufō virgato, de cetero nullo modo differt.

B. annulatus. Pers. Syn. p. 503. Comm. Schæff. p. 45. Schæff. fung. t. 114. Bull. champ. t. 332.

γ. *sericeus* pileo subcastaneo sericeo.

B. luteus Schæff. t. 84.

In filvis frequens.

Var. β . præter notas allatas omnino convenit, in Europa meridionali frequentior esse videtur; rara autem in Suecia ubi a vulgatissima.

Obs. Verus est *Bolet. - luteus* Linn. contra sententiam summi Mycologi *Persoonis*.

139. *Boletus flavidus*, annulatus, pileo pulvinato subglabro cinereo-flavescente, poris angulatis flavescentibus, stipite dilutiori tenui.

In filvis humidis Sueciae.

A priori omnino diversus.

Descr. *Stipes* annulatus 2—3 unc. albidus l. flavescent angustus, diametr. pennæ anserinæ l. parum crassior, æqualis et basi ne minimum incrassatus glaber lævis tenax rectiusculus l. subflexuosus. Variat stipite striato. *Pileus* convexus pulvinatus pallide flavescent l. cinereo-flavus interdum umbonatus glaber fuscus l. coelo pluvioso viscosus 2—3 unc., margine nunc regulari patente nunc irregulari repando, subtus planus. *Caro* flavescent mollis immutabilis, interdum vero cinerascens. *Tubuli* $\frac{1}{4}$ — $\frac{1}{3}$ unc. flavi. — *Pori* angulati magni difformes subcompositi in stipitem fixi et sæpe decurrentes.

Affinitate quadam naturali cum *Bolbo* conjunctus.

Differt; 1.) ab *B. cortinato* P. (species mihi dubia) stipite tenui æquali, carne flavescente nec crispa, pileo porisque aliter coloratis.

2.) a *B. decurrenti* Schum. et 3.) *elegante* Sch. poris angulatis maximis, colore &c.

4.) a *B. cinereo* P. (floccopode *Fl. Dan. t.* 1252) pileo glabro pluribusque characteribus.

** Stipite squamuloso.

140. *Boletus niveus*, pileo pulvinato glabro niveo, poris subrotundis stipite verrucisque concoloribus.

In fagetis umbrosis humidis et pinetis.

Descr. Subgregarius. *Stipes* solidus elongatus 3—5 unc. candidus sursum sensim angustatus e punctis concoloribus reticulato-leaber squamulosus, tactus cinerascit, basi albotomentosus nec bulbosus. *Pileus* pulvinatus convexus carnosus mollis lævis glaber albus, demum virescens, 2—unc., margine patente lævis, subtus planus. *Caro* mollissima alba digitis tacta cinerascit. — *Tubuli* albidii subcinerascetes carne pilei longiores. *Pori* candidi minuti rotundi.

B. scabro P. (*viscido* vero *Linnaei*) affinis. — Hujus varietas lævis est *B. rufus*, *Perf. Obs.* 2. p. 13, nec specificè differre videntur *B. aurantius*, *leucopodius* et noster *niveus*.

Ad hanc sectionem pertinent species nominatæ et *B. circinans* P. e Suecicis.

*** Stipite lævi l. reticulato.

141. *Boletus bovinus*, pileo pulvinato glabro flavescente. sordido, poris compositis acutis flavo-griseis, porulis angulatis brevioribus, stipite pallescente.

B. bovinus, *Linn. Fl. Suec.* 1246. (*exclus. plur. syn.*)

In silvis Sueciæ, ubi vulgaris; sed vix alibi.

Obs. De distinctissima hac specie dubii sunt omnes Botanici apud exteros. Et cum horum neminem illum descripsisse l. depinxisse viderim, extra Sueciam vix crescere suspicor. *B. bovinus* *Gmel.* ad *edulem*, *Schum.* ad *subtomensum* etc. pertinent.

Descr. *Stipes* 2—3unc. subæqualis, crassit. $\frac{1}{2}$ —1 unc., sæpe curvatus elasticus nudus lævis l. e. poris decurrentibus reticulatus, glaber firmus flavesens, numquam rubicundus. *Pileus* carnosus convexus pulvinatus demum planiusculus etiam vertice depressus glaber

lævis 3 uncial. et ultra, subficcus, sed æstiate pluviola viscosus, qua nota a *subtomentoso* etiam differt, sordide flavescent l. cinereo-flavus, margine patente. Caro semper immutabilis pallida spongiosa. Tubuli nunquam $\frac{1}{2}$ unc. attingentes carne pilei breviores. Porii flavogrisei omnium maximi, e pluribus (3—4) brevioribus conformibus compositi, angulati, interdum flexuosi.

A. Gregarius *Fl. Dan.* huic proximus.
142. *Boletus subtomentosus*, pileo pulvinato subtomentoso subflavo-cinerascenti, poris subsimplicibus majusculis læte flavis, stipite subrufescente.

B. subtomentosus. *Linn. Suec.* 1251. *Persf. Syn. p.* 506. *EjUSD. Obs. myc.* 2. p. 9—11. (ubi varietates, synonyma etc. optime determinantur.) *Albert. et Schwein. l. c.* p. 239. B. chryseuteras. *Schrad. spic.* p. 144.

B. bovinus. *Schum. Sall.* 2. p. 376.

Icones. Schæf. f. t. 112. 133. *Fl. Dan. t.* 1074. *Bull. h. t.* 393, 495, 499. f. 2. *Bolt. t.* 87. Vulgaris. 2 et 3.

Descr. Conf. Persf. l. c. non tamen superflua fungi Suecici, ut magis differentia ab antecedenti, cui tamen minime affinis, eluceat.

Stipes 1—3 unc. solidus, statu senescit interdum cavus, crass. digiti minioris majorisve subæqualis glaber carnosus flavescens l. subrubicundus, plus minusve reticulatus, etiam lævis. *Pileus* carnosus pulvinatus $1\frac{1}{2}$ —5 unc. olivaceo-flavescens, alutaceo cinerascens & ferriceo-tomentosus l. glaber, nunquam viscidus, margine obtuso, subtus initio concavus mox planiusculus. *Tubuli* carne pilei subæquales flavi. *Pori* flavi (numquam in priori) majusculi angulato-subrotundi inæquales. *Caro* immutabilis, sed interdum coerulescit.

143. *Boletus piperatus*. P. Sow. t. 34. Swartz l. c. 1810. p. 9. (cum descr.)

β. *squamosus*, pileo pulvinato, squamulis dilute flavis obsito, stipite brevi.

Color pilei flavescens. Loca humidiora ac vulgaris eligit. De cetero minor.

144. *Boletus felleus*, pileo molli dilute castaneo-fulvo glabro, poris albido-roseis subrotundis carneque demum roseis, stipite reticulato.

B. felleus. Bull. champ. t. 379. (optime) Albert, et Schwein. p. 240.

In pinetis ad Femsiö Sueciæ nonnulla legi specimina.

Descr. Inter pulchras species, solitarius. *Stipes* nudus carnosus subrectus 2—4 unc.

glaber reticulatus subæqualis l. basi incrassatus $\frac{1}{3}$ —1 unc. crassus, in nostro olivaceo-fulgineus, intus albus, marginem versus sæpe virescens, vulneratus dilutissime roseus. *Pileus* carnosus junior convexus, dein planiusculus dilute castaneus l. fulvus glaberrimus lævis $1\frac{1}{2}$ —4 unc., carne alba demum subrosea mollissima spongiosa, margine obtuso. *Tubuli* longissimi molles recti sæpius albido-rosei. *Pori* subrotundi regulares albido-rosei, tacti sæpius obscuriores.

145. *Boletus alutarius*, pileo molli alutaceo glabro, carne subimmutabili, poris albis subroseis, stipite elongato læviusculo.

In pinetis, numquam vero in prioris consortio, rarissime. Medius inter *felleum* et *cyaneum*.

Descr. Priori affinis: at ut ejus subspecies non censenda. Characteres constanter servat, et majore jure, ac *aurantiacus* P. a *scabro* distinguitur, eodem modo ut *rubeol.* a *lurido*. *Stipes* elongatus 4—5 unc. sursum subangustatus glaber lævisculus l. tenue reticulatus albidus carnosus, carne immutabili. *Pileus* convexus pulvinatus carnosus molliusculus pallide alutaceus, sed sordidus, glaber lævis 3—4 unc. *Caro* alba immutabilis l. dilutissime

rosea. *Tubuli* mediocres albi, sæpius dilute roscæcentes. *Pori* l. difformes l. subrotundi albido-rosei.

146. *Boletus cyaneſcens*, pileo pulvinato planiusculo flaveſcente l. ſubfuligineo, poris albis carneque mox cyaneſcentibus, ſtipite lævi.

B. cyaneſcens Bull. h. t. 369. *Perſ. ſyn.* p. 508 (*ſub conſtricto*).

In ſilvis Galliæ, Angliæ, Sueciæ.

Quoad colorem fungus noſter recedit, ſed omnia dubia, quin hujus eſſet loci, *Icon Bulliardiana* optima, ſuſtulit.

Descr. Species notabilis et a reliquis carne coeruleſcente præditis quam maxime differt. *Diagnosis* noſtra in fungum et gallicum et ſuecicum quadrat; ſed deſcriptiones tamtumodo fungi a me lecti dabo.

Stipes, ut in reliquis, ſolidus; ſed in hac ſpecie medulla molliori ſubſpongioſa (nec elatiſtica) cum ſtipite contigua factus, nudus $1\frac{1}{2}$ — $2\frac{1}{2}$ unc. deorſum incræſſatus ventricofus ſemper *lævis* glaber l. villoſo-pulverulentus albus, interdum flaveſcens, tactus l. aëri expoſitus olivaceus l. fuligineo-cinereus, intus albus coeruleſcens. *Pileus* junior ſemiglobofus, mox convexus ſubtus planus, demum planiusculus inferne convexus, carnoſus haud elatiſticus 2—5 uncial. flaveſcens, nunc læte

nunc pallide, nunc etiam subcinereus, villosopulverulentus, e pilis quasi adpresfis brevibus, raro glaber, margine patente acuto. *Caro* albida, rupta mox coerulescit, compressa succum coeruleum sæpius copiosum fundens. *Tubuli* breves albi coerulescentes. *Pori* minuti subrotundi albi, a stipite liberi, etiam colorem coeruleo-cyaneum brevi assumentes.

147. *Boletus variegatus*, pileo pulvinato fordide flavo fasciculato-piloso, pilis adglutinat, poris viridi-luteis, stipite lævi flavo. *Swartz in Vet. Ac. Handl.* 1810. p. 8.

In pinetis Suecicis vulgaris.

Descr. *Stipes* nudus 1—3 unc. crassit. $\frac{1}{2}$ —1 unc. subæqualis glaber omnino lævis cylindricus firmus unicolor flavus, raro rubescens. *Caro* stipitis pallide flava subimmutabilis elastica nec spongiosa. *Pileus* plano-convexus nunc piloso-squamulosus nunc glaber ficcus l. tempestate pluviosa subviscosus flavus, margine acute lævi, subtus junior concaviusculus dein planus. *Diam.* 3—5 unc. *Caro* pallide flavescens mollis nec elastica crassiuscula subimmutabilis rarius coerulescens. *Tubuli* carne pilei breviores cinereo-flavescentes hymenio læte flavo villosiusculo adfixæ. *Pori* obscure flavi, juniores fere spadicei dein

cinereo. l. viridi-flavescentes æquales minuti subrotundi obtuse angulati.

Stipitem nonnisi rarissime rubicundum vidi.

148. *Boletus pachypus*, pileo pulvinato alutaceo l. olivaceo-fuscescente, poris demum flavis (*minime rubris*) carneque subcoeruleis, stipite reticulato flavo rubroque vario, basi bulbofo.

B. olivaceus. Schæf. fung. t. 105.

B. vitellinus. Persf. Obs. myc. 2. p. 11. *hujus, ni fallor, loci. Descr. bona.*

In fagetis umbrosis Germaniæ, Sueciæ, ad Almåhs.

Sequenti affinis, sed vere distinctus, et qua ratione e Synopsi a *Persoonio* exclusus nescio. A *Lurido* P. *rubeolario* P. (Bull. t. 100) et *erythrop* P., qui familiam l. tribum naturalem efficiunt, abunde distinctus: *poris numquam rubris*, eademque nota ab his differt A. *calopus*. P.

Descr. *Stipes* brevis 1—2 uncialis ovatus crassus apice angustatus 1 unc. crass. et ultra carnosus firmus evidenter reticulatus deorsum medioque ruber apice flavus (et hunc colorem constantem vidi, qualem etiam descripsit *Persoon*,) glaber l. deorsum radicalis paucis tomentiformibus, carne margine flave-

sciente intus albo mox coerulescente. *Pileus* convexus fuscescenti-cinereus olivaceus l. alutaceus 2—8 unc. latus compactus subtomentosus demum glaber, margine subinvolutumox patente obtuso. *Caro* albida, parum coerulescens et nonnumquam subimmutabilis, qualem etiam fungum suum optime descripsit *Persoon*. *Pori* minuti subrotundi læte flavi, juniores albidii, tacti coerulescentes, *tubulis* concoloribus breviusculis subvirescentes.

149. *Boletus calopus*, pileo pulvinato subolivaceo, poris demum flavis carneque coerulescentibus, stipite reticulato rubro equali.

B. calopus. *Pers. syn. p.* 513.

B. terreus. *Schæff. fung. t.* 315.

In sylvaticis, ericetis & pasim.

Descr. *Stipes* carnosus æqualis 3—6 unc. glaber reticulatus ruber unicolor l. deorsum nonnumquam olivaceus, numquam apice flavus, 1 unc. crass. firmus, intus flavescens mox coerulescens, substantia fibrosa. *Pileus* convexus demum plano-pulvinatus olivaceus l. pallide flavescenti-cinereus, junior subvillosus, dein glaber læviusculus mollis 4—6 unc. *Caro* albida mollis demum spongiosa mox coerulescens. *Tubuli* sublongi læte flavi, fracti mox coerulei. *Pori* æquales in omni

ætate læte flavi juniores tenuissimī, adulti æquales subrotundi.

150. *Boletus? sistotremoides*, pileo utrinque plano castaneo-rufo, stipite rufescenti pallido tenui, hymenio gyroso-poroso venoso denticulato, gyris a pileo seperabilibus.

Paradoxi hujus fungi specimina plurima gregarie in silva *Femfjönensi*, mense *Septembri*, legi.

Descr. Gregarius. *Stipes* solidus nudus 2—4 unc. subæqualis, crassit. digiti minoris erectus l. parum curvatus glaber, basi subvillosus, subrufescenti-pallidus l. parum flavescens. *Pileus* carnosus utrinque planus castaneo rufus l. vaccinus l. fusco-brunneus 2—3 unc. submollis, aqua atmosphærica bibulus, glaber l. subsericeus, margine patente obtusiusculo. *Caro* mollis aquosa albida immutabilis. *Hymenium* gyroso-porosum venosum flavescens, demum subfulvum; margine interdum poris angulatis integris; tubulis brevissimis vix 1 lin. a substantia propria formatis et a pileo liberis, et medio in dentes difformes compressos obtusos lacerum.

Sistotremati quoad characteres, Boletis quoad substantiam proximus. An vero monstrosa progenies? Nulli vero alio fungo a me viso similis.

POLYPORUS. MICH. restit. FRIES.

Mich. Gen. p. 129. t. 70. f. A. B.

Pori integri toti e substantia pilei formati, cum qua et inter se connati. *Habitus: Stipitati, dimidiati et resupinati; carnosi, coriacei ut plurimum suberosi, tubis brevibus pileo arcte adnatis nec disjungendis.*

Obs. 1. Inter *Polyporos* et *Porias* nulli dantur limites et plurimi *Boleti* acaules etiam resupinati variant. Ceterum, præter formam, characteres habituales solum præbentem, nullæ dantur notæ,

Obs. 2. Quamvis omnes hujus Generis species notas in *Mscr.* disposuerim, has hoc loco enumerare superfluum duxi, cum non solum nomina trivialia *Persoonii*; sed sæpe etiam specifica retinuerim. Hinc rariores nonnullas modo describam species.

I. Mesopus Albert. et Schwein. pileo integro stipitato; sed sæpe excentrico. (*Pelteia* Fries.)

151. *Polyporus subsquamosus*, pileo carnosio albido subsquamoso, poris oblongis flexuosis niveis, stipite brevi centrali. *Disp. Boletoid. Mscr.*

Boletus subsquamosus. Linn. Fl. Suec. 1250.

In pinetis Sueciæ, rarior.

Descr. Stipes brevis $\frac{1}{2}$ unc. l. parum ultra crassit. fere unciali cylindricus acqualis carnosus glaber lævis (reticulatum non vidi) sordide albus l. cinerascens. *Pileus* carnosus compactus firmus forma varia, nunc convexus, sæpe planiusculus, nunc subdepressus, 2—5 unc. latus ficcus glaber sed in squamulas discedens (*Hydno imbricato* subfimilis) ficcus pallidus sordide albus l. subflavescens, margine junioris inflexo demum evoluto reflexo subundulato glaberrimo acuto. *Caro* dura alba crassiuscula immutabilis. *Tubuli* vix ulli, nec ad *Boletum* pertinet, ut suspicatus est Illustr. Swartz (in *Vet. Ac. Handl. l. c.*) *Squamulae* diffformes apice subpatentes. *Pori* nivei minuti elongati flexuosi diffformes,

Species optime distincta. *Bolet. caranthiacus* Pers. *syn.* p. 514. (*Wulf. in Jacq. Coll.* 1. p. 342.) si non idem, saltem varietas.

152. *Polyporus ciliatus*, pileo spadiceo-rufo suberoso coriaceo margine ciliato, poris minutissimis subrotundis pallidis, stipite tenui centrali. *Disp. Bol. msc.*

In truncis Sueciæ, rarior.

An distinctus ab *B. ciliato* Fl Dan., qui ad hunc l. *P. circularium* pertinet.

Descr. *Pol. circulario* maxime affinis, sed colore obscuriori et præcipue poris minutissimis æqualibus rotundis abunde diversus. Ceterum *Stipes* uncial 1, fere et subradicatus, sæpe flexuosus et *Pileus* ciliis marginalibus non rigidis oblitus.

153. *Polyporus nummularius*. *Disp. Bol. Mscr.* (*Bol. num. P.*)

γ. minimus, minutissimus, pileo plano subtus convexo, stipite centrali deorsum nigro.

In stipitibus *filicum*, *urticæ* etc. locis umbrosis humidis dejectis.

Descr. *Stipes* 1—4 lin. æqualis durus lignosus glaber subflexuosus læviusculus, superne flavesens inferne niger. *Pileus* $\frac{1}{2}$ — $1\frac{1}{2}$ lin. nec ultra suberosus crassiusculus glaber lævis subnitens junior albidus demum flavesens, su-

perne planus, inferne convexus, *poris* minutis rotundis æqualibus pallidis.

II. *Labellum*, stipite laterali, manubrii instar elongatus (unde nomen: λαβη.)

154. *Obs.* Familia hæc cum priori sæpe confluit, ex. gr. *P. badius*, *varius* &c. qui nunc integri nunc dimidiati reperiuntur. Ad hanc sectionem de cetero pertinent: *Pol. frondosus*, *giganteus*, *platyporus*, *lucidus* &c.

III. *Acorus*, Dimidiati sessiles, stipite nullo.

155. *Polyporus croceus*, pileo dimidiato effuso-sessili subcarnoso subtomentoso lutescente, poris croceis.

Boletus croceus. *Pers. syn. p. 525. Obs. myc. I. p. 87.*

In Quercu cava in *Suecia* legi.

Obs. Optime demonstrat, quanta inter *Polyporos* et *Porias* affinitas; fungus enim meus Quercum nominatam lateri interiori totus obtegens, totus fere erat resupinatus et ad margines tantum reflexus, qualem *Persoon* descripsit.

156. *Polyporus chioneus*, pileo dimidiato sessili carnosio glabro porisque candidis.

In truncis putridis.

Descr. Sessilis subsolitarius. *Pileus* basi resupinato dilatatus utrinque subconvexus carnosus mollis albus 1—2 unc. junior subpubescens, dein glaberrimus, margine acuto. *Substantia* carnosa mollis aquosa candida. *Tubuli* 2 lin. sublongi *Porique* rotundi glabri candidi minuti tenues. Subodorus. *P. styptico* proximus; sed margo pilei concolor, nec cæspitosus et gibbosus &c. Magis affinis *P. destructori*, sed poris glabris &c. diversus.

157. *Polyporus amorphus*, acaulis subimbricatus difformis terrestris carnosus albus, poris difformibus flavescentibus.

In foliis dejectis *Pini Abietis* rarissime.

Descr. Cum *P. heteroclito* comparandus. Polymorphus. *Pileus* carnosus subresupinato-reflexus subcæspitosus albus subpubescens odor subcupulæformis, margine acuto. *Tubuli* brevissimi flavescentes, poris subrotundis difformibus flavescentibus. Forma *Mer. tremelloso* similis.

158. *Polyporus dichrous*, pileo sessili subimbricato molli coriaceo villoso albo, poris minutis subrotundis fusco-cinnamomeis.

Boletus pelleporus. *Bull. champ.* t. 501.
f. 2. (*non differt.*)

Ad truncos fagineos, *Sueciæ*, *Galliæ* &c.

Descr. Thelephoræ obiter inspectus superne haud dissimilis. Cæspitosus, sæpe imbricatus basi effuso resupinato plurima specimina conjungente. *Pileus* coriaceus flexilis molliusculus (et hinc ab *P. conchato* Pers., cui affinis videtur, distinctus,) 1—2 unc. longus candidus villosus, hirsutiae alba adpressa, zonis elevatis subobsoletis concoloribus, margine acuto patente concolore. *Caro* tenax nivea. *Pori* minuti subrotundi æquales integri fulco-cinnamomei.

159. *Polyporus pubescens*, pileo sessili subimbricato carnofo-suberoso pubescenti albo, poris minutis subrotundis concoloribus.

B. pubescens. *Schum. Sæll.* 2. p. 384.

In truncis betulinis *Sueciæ*. *Daniæ*.

Descr. Gregarius subcæspitosus imbricatus. *Pileus* junior carnofo-suberosus dein suberosus pulvinatus 1—2½ unc. latus, crass. varia, albus subreniformis, zonis elevatis pubescentibus latiusculis variegatus, ceterum fere glaber, margine patente acute concolore l. lutescente (in meo numquam ferrugineo), subtus planus. *Pori* subrotundi albi, demum subflavescentes, minuti glabri.

160. Plures in parte secunda. Ceterum sequentes, ut Novitiæ Fl. Suecicæ, nominandi:

Polyp. betulinus. Bol. *betulinus*. P. Fl. Dan. t. 1254.

In truncis *betulinis* frequens.

Pol. fraxineus. B. *fraxineus*. P. syn. Bull. champ. t. 433. f. 1.

Pol. ochraceus. B. *ochraceus*. Pers. syn. Schaff. t. 269.

Hic in truncis *populneis* frequens; ille semel in trunco *fraxineo* visus ad Bökeberg in par, Femsiö.

Pol. crispus. Bol. *crispus*, Pers. Obs. myc. 2. p. 8. Albert. et Schwein. l. c. p. 249.

In truncis *fagineis*.

Pol. ribis. Bol. *ribi*. Schum. Sall. 2. p. 386. (affinis *fomentario*, ut recte annotavit Re-bentisch.)

Suo loco frequens ad Lundinum.

Pol. angulatus. Bol. *angulatus*. Schum. Sall. 2. p. 388.

In truncis *fagineis*.

Et nonnullæ aliæ species e Schumacherianis, alio loco describendis, Ceterum conf. Swartz l. 6.

IV. PORIA. Resupinati.

161. *Polyporus laticrimus*, effusus crassus suberosus undulatus subpallidus, poris demum prominulis inæqualibus, tubulis brevissimis.

In truncis fagineis dejectis muscosis. Difficile describitur distinctissima species. *Substantia* suberosa crassa albida inter zonata. *Pileus* resupinatus latissime effusus ($\frac{1}{2}$ —3 pedalis) firmus pallidus, tactus subobscurior; margine angustissimo albo sterili, sed non byssino, hinc inde libero, sæpius adnato. *Superficie* junior sæpe apora, dein poris hinc inde prominulis subliberis rotundis minutis punctiformibus, absque tubulis; demum tota porosa, poris variis, aliis elongatis angustis flexuosis, aliis rotundis, tubulis brevissimis cum pilei carne confluentibus.

162. *Polyporus xanthus*, effusus elongatus luteus, margine subnullo glabro, poris subrotundis obliquis, tubulis mediocribus.

In affulis semiputridis, serie elongata. Colore lætissime luteo ab omnibus differt. *Subiculum* subnullum. *Pori* ejusdem formæ ac *P. obliqui*, cui proximus, sed crebrius prominuli et subrotundi — 2—3 unc. longus,

2 lin. crassus, duriusculus et ficcatus omnino immutabilis.

163. Speciebus hujus familiæ ab illustr. *Swartzio l. c.* descriptis, ut indigenæ, addendæ:

Pol. salicinus. — *Bol. salicinus*. *Perf. syn.* p. 543.

Suo loco in Scania, Smolandia.

P. destructor. — *B. destructor*. *Schrad. spic.* p. 166.

In domibus. Illustr. Prof. et Eques *Acharius*. et in truncis pineis.

P. contiguus. — *B. contiguus*. *P. syn.*

In truncis fagineis.

P. molluscus. — *B. molluscus*. *P. syn.*

Ad ligna pinea.

P. epiphyllus. — *B. epiphyllus*. *P. Obs. myc.* 2. p. 15.

In foliis dejectis; a priori sine dubio distinctus, sed variis aliis nimis affinis.

P. byssinus. — *Bol. byssinus*. *Schrad. spic.* p. 172. t. 3. f. 1.

In truncis pineis.

P. subtilis. — *B. subtilis*. *Schrad. l. c.* p. 173. t. 3. f. 2.

In trunco populi nigra.

P. sanguinolentus. — *B. sanguinol.* *Albert. et Schwein. l. c.* p. 287.

Ad truncos pineos.

P. reticulatus. — *B. reticul.* *Pers.* *Mucilago retic.* *Hoffm. Germ.* 2. t. 12.

Cum priori. &c.

Obs. *Fistulina (hepatica)*, me iudice, distinctum efficit genus.

SISTOTREMA. *Pers.*

*** *Xylo don.*

164. *Sistotrema fallax*, interrupte effusum tenue, dentibus minutis acutis flavescenti-ochraceis, margine byssino albo. *Fries Nov. Fl. Snec.* p. 39.

In hymenio *Polyporis suberosi* parasiticum.

Obs. *Polyp. suberosi* hymenium adeo incrustans, ut pro *Hydri* species, nec *Polypori*, haberi posset, *Subiculum* membranaceum tenue, aculeis flavescenti-ochraceis minutis confertis acutis incisilis, margine sterili byssino albido.

L. c. etiam descripta sunt *S. fagineum* *P. quercinum*, *P. et obliquum* *Albert.*

HYDNUM. *Linne Pers.*

† *Stipitata* (Harum specierum monographia.)

* *Suberosa*.

165. *Hydnum aurantiacum*, pileo azono aurantiaco l. albo-variegato, intus zonato stipiteque aurantiaco.

H. aurantiacum. Swartz in *Vet. At. Hand.* 1810. p. 247. — *Albert et Schwein. l. c.* p. 265. — *Fl. Dan. t.* 1439. (*A tribus Auctoribus eodem fere tempore eodemque nomine, ut distincta et nova species [descriptam!]*)

Hydn. suberosum. β. aurantiacum. Batsch. El. fung. Cont. 2. f. 222.

Hydn. floriforme. Schaff. t. 146. *f.* 4. &c.

Hydn. hybridum. Pers. comm. Schaff. p. 59. (*exclus. nonnullis syn.*)

In silvis Femflönenfibus, ubi hoc et sequentia crescunt, optimæ mihi fuerunt occasiones has species observandi et quamvis, Botanicorum sententiæ ut placerem, hoc loco distinxerim, nullos inter hoc et sequens præcipue dari limites, mihi persuasissimum habeo. — Ceterum, ut in *Synopsi Lichenum* p. 103 optime monuit Cel. Prof. *Acharius*, non omnia, quæ confluere possunt, commiscere licet; ut nullus etiam dubito, quin magis erret, qui separet, quam qui conjungat.

Synonyma a plurimis Auctoribus nimis neglecta.

Descr. Swartziano l. c. optima (quare etiam ejus Synonymon primum attuli) sed in omnes status non quadrat. *Pileus* totus sæpe albus. — Ceterum Substantiæ internæ color certissimos semper mihi præbuit characteres.

166. *Hydnum compactum*, pileo azono flexuoso rugoso olivaceo-ferrugineo tomentoso, intus fusco et coeruleo variegato, stipite brevissimo.

Hydnum compactum. *Pers. syn.* p. 556. *Swartz in Vet. Ac. Handl.* 1810. p. 245. *Wahl. Fl. lapp.* p. 532.

Hydnum floriforme. *Schæff. t.* 146. f. 35. 147. f. 7. &c.

H. suberosum a. *spongiosum.* *Batsch. l.* c. f. 221. a.

In filvis ericetosis muscosis.

Descr. *Varium flegarium subconfluens.* *Stipes* brevissimus durus crassus subtuberosus rigidus difformis subangulosus, sæpe subnulus. *Pileus* olivaceus l. ferrugineo-spadiceus, tomento concolore l. albido obtectus; aut turbinatus superne planiusculus subregularis læviusculus 1—2 unc. latus, aut dilatato-confluens terram crustæ instar obtegens conve-

xiusculus undulato - tuberculatus 3—6 une. margine inæquali, coelo pluvioso aqua repletus. Caro suberosa tenax coeruleo et fusco-variegata. *Tubuli* numerosissimi longiusculi sæpe obliqui spadicei fusci l. castanei.

167. *Hydnum ferrugineum*, pileo azono ferrugineo-rubro molli-suberoso, intus stipite-que elongato concolore.

Hydnum hybridum. *Bull. herb. t. 453. f. 2.*

β. *glabratum*, pileo glabro.

In pinetis.

Descr. Gregarium nec confluens. *Stipes* duriusculus 1½—3 unc. subdifformis ferrugineus, intus concolor, rectus, basi vix incrassatus. *Pileus* molliusculus planus l. depressus obconicus l. sæpius dilatatus subconnatus villosus ferrugineus 1—4 unc. *Substantia* mollior ferruginea unicolor. *Sibuli* recti firmi tenaces longiusculi fusco-ferruginei.

Varietatem *Hydn. suberosi* ferrugineam etiam commemorat *Cel. Barsch.* Magis a præcedentibus differt; saltem ab *aurantiaco* distinctissimum.

168. *Hydnum pullum*, svaveolens, pileo subzonato sericeo tomentoso, intus albo-coeruleo-variegato, stipite inæquali coerulescente.

Hydnum pullum. *Swartz l. c. p. 248.*

H, suberosum γ . cinereum, *Batsch. l. c.*
f. 223.

In filvis, cum prioribus.

Species a præcedentibus optime distincta; sed cum H. *svaveolenti* magnam suspicor affinitatem, saltim bona descriptio ab *Albertini* et *Schweinitz l. c. p. 264* vix recedit.

Descr. Gregarium subconcreescens suberosum obconicum svaveolens. *Stipes* brevis, unciam haud superans, plus minusve albido-coeruleus rigidus subtomentosus, basi sæpe bulbosus. *Pileus* turbinatus aut simplex aut cum vicinis coalitus difformis varie undulato-lobatus, junior sericeo-pubescent deinde glabriusculus subrugulosus albus l. sordide cinereus, rarissime in meis speciminibus fusco-olivaceus, fuscus (nec viscosus, ut H. *svaveol.* describitur) planus l. subdepressus marginem versus sæpe obsolete zonatus. *Caro* pilei alba, zonis coeruleis variegata, stipitis unicolor coerulea. *Subuli* cæcio-violacei subcinerei tenues tenaces in stipite decurrentes.

169. *Hydnum nigrum*, inodorum, pileo azono coeruleo-nigro, intus unicolori stipiteque nigro.

In filvis abietinis densis *Suecia*. Optime distincta species.

Descr. Cæspitosum confluens difforme. *Stipes* suberosus uncialis radicans rigidus inæqualis, ex eadem basi plures et sæpe confluentes, niger, inferne tomentosus superne glaber, intus unicolor. *Pileus* suberosus, junior clavæformis dein turbinatus demum convexus planiusculus cum vicinis semper fere coalitus, hinc difformis rugulosus tuberculatus, coeruleo-niger, tomento concolore l. albedo vestitus, margine acuto concolori l. albedo. *Caro* crassiuscula dura unicolor nigra nec zonata, quo caractere a reliquis distinguitur. *Subuli* albidii tactu cinerascetes tenaces breviusculi.

Obs. Minimè H. *tomentoso* affinis species, ut e descriptione pilei forte quis suspicaret.

170. *Hydnum infundibulum*, pileo carnosio suberoso infundibuliformi spadiceo lævi, margine flexuoso-plicato, subulis albis demum badiis, stipite toruloso inæquali. *Swartz l. c. p. 244.*

In pinetis Sueciæ, rarius.

A prioribus abunde differt substantia, stipite 2—3 unc. deorsum angustato, pileo infundibuliformi &c. Ad sequentem sectionem transitum efficit. Semper *coriaceos* suberosis maxime affines esse credidi, sed aliud natura docuit. *Descr.* hujus dedit *Swartz l. c.*

** Carnosa.

171. *Hydnum coeruleum*, carnosum obconicum, pileo convexo coeruleo centro al-bido, subulis albidis, stipite basi fusco. *Fl. Dan. t. 1374.*

In pinetis Norvegiæ. *Vahl.*

Obs. Speciebus dubiis adnumerare non debui, cum e figura distinctissimam esse speciem patet. Gregarium. *Stipes* brevissimus vix $\frac{1}{2}$ unc., crassus $\frac{1}{3}$ —1 unc. glaber lævis colore chocolatæ sursum in *pileum* 1—3 unc. violaceo-coeruleum convexum expallescentem crassum dilatatus. Aculei albi in stipite decurrunt. (*Quoad icon.*)

β . *violascens*, carnosum-fibrosum sordide violaceum, pileo rugoso flexuoso sublobato tomentoso, margine albicante, subulis tenuibus acutis albis, stipitibus basi subtuberosis connatis longiusculis. *H. violascens. Albert. et Schwein. l. c. p. 265.*

In pinetis Germaniæ. *Albertini et Schweiniz.*

Obs. Prioris videtur varietas. Color stipitis (omnino), pilei ex albo-coeruleo variegati et subularum, consistentia, subuli &c. omnino conveniunt. Differt præcipue stipite elongato subtuberoso.

172. *Hydnum imbricatum*, pileo planiusculo umbrino squamoso, subulis albido cinereis, stipite abbreviato.

Hydn. imbricatum. *Linn. Suec. n. 1257* (cum descr. ad n. 1256) et *Auct.* — *Perf. syn. p. 554. Schæff. fung. t. 140. Sow. fung. t. 73. Fl. Dan. t. 176 & t. 1500.*

Hydn. cervinum. *Perf. Obs. myc. l. p. 74. Hall. helv. 2324. β.*

In pinetis autumnno frequens.

Obs. Illustr. a *Linné* non descriptiones proprias solum confudit, sed etiam synonyma ex. gr. *Erinaceus esculentus albus crassus Mich. Gen. t. 72. f. 2.* ad hunc fungum dubie pertinet; sed hujus synonymon e *Mic helio* est: *Erinaceus esculentus*, pileo superne veluti squamoso et fuliginoso, inferne cinereo, pediculo cum superna pilei parte ejusdem coloris. *Mich. Gen. p. 133*; nec in hunc quadrant synonyma *Halleri*, *Bauhini* et *Raji*; nisi, ut monuit *Haller l. c.* hujus speciei varietas sit.

Descr. Solitarium l. gregarium, nunquam cæspitosum. *Stripes* brevis uncialis crassus deorsum subangustatus glaber lævis firmus solidus cinerascienti-albidus, *Pileus* carnosus, junior convexus dein planiusculus, latissimus palmaris et ultra subfirmus umbrinus l. fusco-subcinereus squamosus, squamis apice

subreflexis imbricatis, centro erectis obscurioribus; versus marginem acutum pallidior. Caro albida. Subuli teretes acuti numerosissimi subtenaces recti l. rarius obliquialbido-cinerascentes.

Hydn. subsquamosum *Batsch* (*El. fung.* p. III. f. 43. *Hydn. squamosum*. *Bull.* p. 310. t. 409) huic nimis affine videtur; distinctam tamen esse speciem urgent *Albert.* et *Schwein.* l. c. p. 264. Ipse vivum non vidi et nec ex *Iconibus* neque e *descriptionibus* sufficientes inveni characteres, quare ex hac monographia *Hydnorum stipitatorum*, a me rite cognitorum, exclusi.

173. *Hydnum repandum*, pileo carnoso subrepando flavo l. rufescente glabro, subulis subteretibus pallidis stipiteque difformi.

Hydn. repandum. *Pers. syn.* p. 555. *Obf. myc.* p. 90. *Schum. Sæll.* 2. p. 293.

a. *Denudatum*, pileo repando glabro lævi.

Hydn. repandum, pileo convexo lævi flexuoso, *Linn. Suec.* 1258. cum *syn. Mich. Dillen. et Hall.*; sed descr. ad 1257 et *Auct.*—*Sow. fung.* t. 176. *Bull. h. t.* 172. *Fl. Dan.* t. 310.

a.) *Flavidum*, pallide luteum.

Hydn. flavidum. *Schæff. fung.* t. 318. *Erinaceus* esculentus pallide luteus. *Mich. Gen.* p. 132. t. 72. f. 3.

b. *Rufidum*, rufescens.

Hydn. rufescens. *Schaff. fung. t. 141.*
fig. plur.

H. carnosum. *Batsch. El. fung. t. 26. f. 136.*

Echinus petiolatus subrufus, petiolo
lævi. *Hall. helv. 2325.*

c. *Albidum*, pallidum.

H. medium. *Perf. Obs. myc. 2. p. 97.*
hujus sine dubio loci.

β. *Squamosum*, pileo repando subsquamoso.

Hydn. squamosum. *Schaff. f. t. 273.*

H. imbricatum. *Bolt. f. t. 87. (excl. syn.)*

γ. *Intermedium*, pileo orbiculari regulari
lævi.

Boletus pileolo pulvinato integerrimo
&c. *Gled. meth. fung. p. 74. n. 6.*

Vulgare in fagetis et pinetis.

Descr. Fungus gregarius l. subcæspitosus, colore forma et superficie valde variabilis. *Stipes* irregularis flavescens glaber fragilis deorsum interdum incrassatus, longit. et crass. varia. *Pileus* difformis, in γ. regularis, flexuosus l. lobatus flavus, sed variat nunc ruber nunc albidus, lævis l. rugulosus, in β. subsquamosus, fragilis margine inæquali concolore; interdum dimidiatus. *Subuli* crassiusculi fragillimi flavescences pallidi, nunc teretes integri, nunc incisi. *Substantia* carnosa pallide flava.

Plures ad hoc adhuc referri debent species ab Auctoribus descriptæ. Nec differt *H. carnosum* pileo convexoglabro rubello, aculeis griseis stipiteque subconcoloribus. *Gmel. Syst. Linn.* 11. 2. p. 1438.

174. *Hydnum rufescens*, pileo orbiculari rufescenti-subcarneo subtomentoso, subulis subcompressis incarnato-ochraceis, stipite cylindrico.

H. rufescens. Pers. syn. p. 555. Obs. myc. 2. p. 95.

H. repandum. Bolt. fung. t. 88. (sec. Pers.) Meum fungum in fageto legi.

Descr. Priori tantum affine, ut specie vix distinguendum; nec notæ, quibus distinguitur, sunt constantes. Differt: *Stipite* subæquali tenuiori albido, *pileo* regulari tenuissime tomentoso marginem versus interdum obsolete zonato, subulis sæpius incisfis aliter coloratis.

175. *Hydnum lævigatum*, pileo carnosio convexo castaneo lævi, aculeis mollibus albido-cinereis, stipite inæquali elongato. *Swartz l. c. p. 243. — Mich. Gen. p. 132. t. 72. f. 1.*

β *Coriaceum*, pileo coriaceo planiusculo lævi, aculeis tenacibus albidis, stipite æquali elongato.

γ . *Brachypus*, pileo carnosio-coriaceo plano subdepresso lævi, aculeis tenacibus albido-cinereis, stipite brevi crasso.

In pinetis Suecicis α) legit Cel. *Schwartz*, β . γ . a me lecta sunt.

Varietatem α . descripsit *Schwartz* l. c. et ad hanc fungum meum pertinere vix credi. diffem, nisi Ipse illustr. vir specimina mea pro suo fungo agnovisset. Var. β . *Descr.* *Stipes* elongatus crass. pennæ anserinæ l. digiti minimi subcarnosus erectus 2—3 unc. glaberrimus æqualis solidus dilute cinereus. *Pileus* carnosus-coriaceus tenax flexibilis planiusculus glaber lævis azonus cinereus, margine sæpius flexuoso vario concolori. *Caro* tenuis alba tenax. *Subuli* numerosi tenues flexiles tenaces albi, ficcati vero rufescentes.

Ab amic. *Forsandro* etiam observatum.

Var. γ . *Descr.* A priori differt: *stipite* brevi l. unc. longo et crasso subdifformi et medio interdum incrassato demum cavo (a vermibus forte perforato) *pileo* 3 unc. plano vertice subdepresso ruguloso, (a foliis pineis scil. obtectus) subulis albidis l. dilute cinereis.

*** Pileo coriaceo.

176. *Hydnum melaleucum*, pileo convexo-plano subzonato glabro vertice scrobiculato umbilicato violaceo-atro, margine aculeisque albis.

Hydn. melaleucum, *Schwartz* in litteris.

Hydn. n. sp. Swartz in *Vet. Ac. Handl.*
l. c. post pulli descr.

H. pullum Schæff. fung. t. 272.

H. zonatum. Gmel. syst. Linn. 11. 2. p. 1438.

H. tomentosum. γ. atroalbum. Albert.
et Schwein. l. c. p. 266. (descr. bona.)

H. cyathif. β. Schrad. Spic. p. 175. exclus.
Syn. Willd. Batsch. &c.

In silvis abietinis muscolis rarissime.

Descr. Suberoso-coriaceum gregarium,
rarius simplex, sæpissime difforme, stipitibus
pileisque confluentibus. *Stripes* uncialis crass.
pennæ gallinæ, vix anserinæ, tenuis elasticus
glaber deorsum sæpe angustatus, sæpissime
monstrosus e pluribus connatis ramosus com-
pressus &c. ater. *Pileus* planus umbilicatus,
sed non infundibuliformis, coriaceus zonatus
coeruleo-niger l. violaceo-ater glaber vertice
scrobiculato, margine albicante l. concolore,
sæpius autem, quam maxime irregularis, plu-
res connati l. imbricati lobati &c. etiam fere
proliferi. *Subuli* albi, tacti subcinerascens,
breves subtenaces, in stipite non decurrentes.

Egregia ab Auctoribus confusa species!
et distinctissima tamen. In synonymis eru-
endis summam posui operam et allata vera
esse spero.

177. *Hydnum tomentosum*, pileo plano-infundibuliformi zonato subcinereo vertice tomentoso, margine aculeisque albis.

Hydn. tomentosum, Linn. Suec. p. 455 (cum descr.) et Florist. Suec. Pers. syn. p. 556. α. β.

Hydn. cyathiforme, Fl. Dan. t. 1020. f. 2. (excl. syn. Mich. et Hall. Schaff. fung. t. 139.

Descr. Plurima specimina confertim sæpius crescunt, pileis stipitibusque subconcretis. *Stipes* prioris subbrevior vix unc. superans crass. fere pennæ anserinæ subglaber cinereus inæqualis basi subbulbosus l. firato subjacente subterraneo adnatus; cum plures connati sint, quam maxime irregularis. *Pileus* planus l. subinfundibuliformis subuncialis flaccidus nunc dilutius nunc obscurius sed semper constanter cinereus, centro obscuriori fusco-tomentoso, tomento brevi, fasciis discoloribus subzonatus, margine acuto patente albido. *Substantia* pilei et præsertim stipitis suberoso-coriacea subzonata. *Aculei* albi firmi breviusculi nec in stipite decurrentes.

Obs. *Hydn. tomentosum*, Fl. Dan. 534. ad hanc speciem non pertinet; sed forte potius varietas monstrosa *Hydni strigosi* Swartz.

178. *Hydnum scrobiculatum*, pileo infundibuliformi azono ferrugineo subpubescente, vertice colliculoso squamoso, subulis fuscorufescentibus, stipite glabro subæquali.

H. cyathiforme, exiguum coriaceum ferrugineum, stipite brevissimo, pileo primum suborbiculari, omnino echinato, demum infundibuliforme. Bull. Champ. p. 308. t. 156. (hujus videtur loci).

H. cyathiforme, gregarium pileo infundibuliformi fasciato fibroso elastico, stipite brevi *Gmel. syst. Linn. 11. 2. p. 1439. (excl. var. β. cinerea)* *Schrad. spic. p. 174. (excl. var. β. zonata)*

Erinaceus alpinus minor discoides fulvo-ferrugineus. Mich. Gen. p. 132. t. 72. f. 7.

In pinetis non infrequens.

Nomina Auctorum specifica, quia illustrant fungum attuli.

Descr. *Stipes* brevis $\frac{1}{2}$ unc. circiter, teres suberosus tenax elasticus subæqualis glaber fusco-rufescens, radice elongata radicans. *Pileus* valde infundibuliformis coriaceo-suberosus (sequentis multo minor sed crassior fere) junior subplaniusculus valde lobato-difformis vix concreescens subtilissime sericeus, vertice tuberculato-scrobiculatus squamoso-fibrosus, valde irregularis semper azonus. *Substantia* interna subsuberosa azona. *Subuli* tenues fragiles densi fusco-rufescentes nec in stipite decurrentes.

Magna cum sequenti, præcipue quoad colorem, affinitas, ut dubius hæserim, utrum conjungerem l. separarem. Separandum tamen est; characteres constanter servat et a sequenti magis differt, quam *rufescens* a *repando* et *aurantiacum* a *compacto*. Ceterum incredibilem in hoc Genere confusionem, Botanicorum conjungendi studio ortam, distinguendo tollenda, nec confundendo augenda; nec illustrat rem, qui varia ob similitudinem commiscet.

179. *Hydnum conrescens*, pileo plano-infundibuliformi ferrugineo-cinereo l. spadiceo-zonato glabro striato, subulis fusco-nitidis, stipite tomentoso inæquali.

H. conrescens. *Perf. syn.* p. 556. *Obs. myc.* 1. p. 74 (*exclus. syn.*) Swartz l. c. p. 246. *Alb. & Schwein.* l. c. p. 266.

H. zonatum. *Willd. her.* 1242. *Batsch El. Fung. Cont.* 2. p. 111. f. 224.

Erinaceus caespitosus coriaceus ferrugineus superne circularibus striis excavatus &c. *Mich. Gen.* p. 132. 6.

Erinaceus infundibulum imitans coriaceum colore ex fulvo-ferrugineo, pileolo desuper veluti fericeo(?) & striis circularibus excavato. *Mich.* l. c. 4. t. 72. f. 4. *vix differt.*

In pinetis satis frequens.

Deser. vide Sw. l. c.

A. priori differt: Stipite longiori bulboso unciali inæquali s. bventricoso ferrugineo-tomentoso, rarissime deorsum glabro; pileo juniore infundibuliformi, demum excavato-planiusculo coriaceo glabro, vertice non colliculoso, sed radiato striato colore e zonis obscurioribus variegato, subulis cinereis, dein fuscescentibus subtenacibus &c.

180. *Hydnum auriscalpium*, pileo coriaceo dimidiato spadiceo, stipite tomentoso.

H. auriscalpium. Linn *Suec.* n. 1260. cum syn. Pers. *Syn.* p. 557. *Comm. in Schæff.* p. 55. Schæff. f. t. 143 Bull. h. t. 481 f. 3. Curt. Lond. t. 190. Fl. Dan. t. 1020. f. 1. Bolt. f. t. 90.

Erinaceus parvus hirsutus ex fusco-fulvus pileolo semiorbiculari, pediculo tenuiori. Mich. Gen. t. 72. f. 8. Hall. *holo.* n. 2321.

Fungus erinaceus parvus in conis abietis nascens. Buxb. cent. 1. t. 57. f. 1. Buxb. Hall. 129. c. fig.

β *bicolor*, pileo dilutiori. Alb. l. c. p. 267.

In silvis passim in conis pineis.

β in abiegnis tantum legi.

Deser. Subsolitarium. *Stipes* 2--3 unc. firmus flexibilis sursum attenuatus crassit. pennæ gallinæ erectus tomentosus spadiceo-niger, basi cortice tomentoso-spongioso Pileus dimidiatus laterali; sed suborbicularis, excisus, lobis haud raro circumscriptis, coriaceus tenax flexibilis fusco-spadiceus l. badius, junior sæpe ferrugineus & pallidus, convexiusculus tomentosus $\frac{1}{2}$ — $\frac{3}{4}$ unc., in margine patente acuto interdum

flexuoso, lateribus sæpe hymenioque dilutioribus. *Subuli* spadicei longiusculi fimi.

**** Pileo gelatinoso.

181. *Hydnum gelatinosum*, albido-glaucum, stipite laterali.

H. gelatinosum. *Perf Syn* p. 560 *Swarz* l. c. p. 249. *Scop. carn.* 2. p. 472. *Jacq. austr.* t. 239. *Alb. & Schweinl* c. p. 263. *a. & β. dichroum*, nec differt *Schæff.* f. t. 144 145, quod

H. clandestinum. *Gmel. Syst. nat.* II. 2 p. 1439.

H. crystallinum. *Fl. Dan.* t. 717.

Ad truncos putrescentes pineos, præsertim abiegnos.

Deser. Gregarium gelatinosum (suum carnosum dicit *Schæffer*.) *Stipes* lateralis brevis aquoso-pallidus mollis teretiusculus subæqualis glaber l. tenuissime papilloso pubescens. *Pileus* junior sublinguiformis l. obovatus deinceps cordato-reniformis aliasve formæ adscendens l. horizontalis $\frac{1}{2}$ —2 unc. subdiaphanus plus minusve planiusculus cinereo-glaucus l. aquose albidus, superficie tenuissime ruguloso-papillosa. *Subuli* dilutiores albidii subdistincti pyramidati molles nec fragiles.

Variat pileo difformi subfessili.

182. *Hydnorum stipitatorum species mihi dubie:*

- 1) *H. suaveolens*, pileo rugoso viscoso subto-
mentoso, aculeis violaceis, stipite brevi crasso.
*Scop. carn. 2. p. 472. Alb. & Schwein 1.
c. p. 267.*

An varietas *pulli* Sw.?

Obs. Pileus dicitur coriaceus; sed habi-
tus *Hydni compacti*.

- 2) *H. cinereum. Bull. champ. t. 419.*

Ad *H. scrobiculatum* retulit *Gmelin* sed vix
recte quoad iconem. Forte distinctum l. to-
mentoso proximum.

- 3) *H. — Erinaceus non vescus ramosus niger.*
Mich. gen. p. 70. c. Desideratur descriptio
aculeorum &c., forte ad *H. nigrum* l. *mela-*
leucum referendum, illo vero, secundum *Gle-*
ditichium l. c. propius accedere videtur.

- 4) *H. album — Erinaceus esculentus albus*
crassus. *Mich. Gen. 1. c. f. 2.* secundum
Hallerum varietas est *H. imbricati*.

De *H. subsquamoso* *Batsch* ad *H. imbrica-*
tum, pro cuius subspecie habeo, meam attuli
sententiam.

183. *Species paradoxæ l. falso ad Hydnum stipitata relata:*

1) *H. lactescens* HaH. *Helv.* 2323.

Hydnum species vix est.

2) *H. spadiceum*. Pers. — Swartz l. c. p. 232. &

3) *H. sublamellosum*. Bull. t. 453. f. 1.

Sistotrema species sunt.

4) *H. tomentosum*. Fl. Dan. t. 534. Ceterum, ut species falsæ, plurimæ nominari possent, quæ l. ex aliis compositæ sunt l. quorum differentia in solis nominibus consistit. Has ad species ad quas pertinent, annotavi.

††† Odontia.

184. *Hydnum pinastri*, resupinatum flavescens membranaceum, margine byssino albedo, aculeis obliquis subadnatis. *Sistotrema pinastri*. Fries *Nov. Fl. Suec.* 2. p. 38.

In ligno pineo putrido.

Descr. Resupinatum interrupte effusum membranaceum leviter adnatum, junius villosum, dein reticulatum, demum aculeatum glabrum, margine byssino. *Aculei* acuti subinte-

gri flavescentes longiusuli, sed subiculo adpressi & fere adnati.

185. *Hydnum fusco-atrum*, resupinatum late effusum ferrugineo-fuscum glabrum, subulis distinctis nigris. Fr. l. c. p. 39.

In ligno dejecto fagineo putrido.

Obs. Hujus, sequentis, *crustosi* &c. descr. l. c. dedi.

186. *Hydnum Agardhii*, effusum subcarnosum albidum glaberrimum intus viridescens, aculeis minutis obtusis, apice rubis. Fr. l. c. p. 39.

In truncis abiegnis ad terram dejectis.

THELEPHORA P.

187. *Thelephora* (Cort.) *maculeformis* adpressa pallide rosea, junior subrotunda, pseudo-papillo unico, mox confluens, margine glaberrimo.

In cortice ramulorum.

Obs. A *Theleph. fallaci* omnino distincta & *Th. laevi* & *roseæ* affinior; sed margine glaberrimo distincta. *Pileus* effusus crustiformis tenuissimus membranaceus glaberrimus dilate & eleganter roseus, statu simplici vix lin., subrotundus mox confluens difformis 2-4

lin. *Papilla* unica centralis a nodulis corticis orta, in speciminibus confluentibus plures, ceterum laevissima.

188. *Thelephora odorata*, fraveolena suberoso-membranacea glabra laeviuscula confluens pallida.

Ad truncos abiegnos destructos putridissimos.

Desc. Resupinata effusa tota arcte adnata suberoso-membranacea crassiuscula glaberrima (etiam in margine) albida l. exalbido-alutacea l. subrufescens 1—2 unc. confluens 4—5 unc. laeviuscula, papillis rarissimis. Odor gratus, fere *B. odorati*, cum quo affinitatem habet, licet abunde distincta.

189. *Thelephora amphibolia*, effusa tenuis brevis fuscescens subalutacea, papillis sparsis majusculis.

In hymenio *Boleti suberosi* tantum, sed saepe legi.

Obs. Inodora & tota arcte adnata color, pallescens. A *Sistotr. fallaci* omnino distincta.

190. *Thelephora ochracea*, effusa latissima subtenuis ochracea, papillis sparsis irregularibus majusculis.

In ligno abieigno putrido.

Obs. A *Th. granulosa* P. vere distincta. Tota aretissime adnata, longissime & atissime effusa pedalis & ultra. Margo nunc glaber nunc hyssinus villosus. Papillæ ab asperitatibus ligni ortæ videntur.

191. *Thelephora gigantea*, maxima difformis subibera glabra lactea, margine sublibero glaberrimo.

Th. lactea. Schrad. *Spic.* p. 186 (quoad diagnos brevissimam hujus quidem loci; sed *Th. lactea* P. longe alia est species.)

Supra corticem *Pini sylvestris*.

Descr. Membrana 5—9 unc. longa lactaque maxime difformis truncos leviter adhærens flexibilis undulata, sine vestigio papillarum, contigua, utrinque margineque glaberrima lactea.

Obs. Species maxime singularis. An distincti Generis?

192. *Thelephora confluens*, subrotunda arete adnata lævissima contigua glabra, margine adnato.

α. alba.

β. lævissime flava.

Th. lucida. Achar. *Sec. spec.*

Ad ramos betulinos, raro.

De fer. Membrana lævissima contigua vix umquam uncialis ramos arctissime adhærens subrotunda ætate confluens irregularis utrinque margineque glaberrima. *Color* nunc albus nunc late flavus, detritus immutabilis, nec fuscescens l. inquinans.

193 *Thelephora cretacea*, effusa humida farinoso-cretacea alba subinquinans.

Theleph. calcea γ . *Pers. Syn* p 581.

Corticium Sambuci. *EjUSD. disp. meth.*

p 31.

Ad truncos *Sambucinos* frequens, in *Pineos* rarior.

Obs. vere distincta species.

194. Varietatas novæ notabiliores specierum descriptarum.

Theleph. frustulata β . *tessulata*, crassissima tessulata cinnamomea, pulvere concolori oblecta.

In ligno ad terram dejecto.

Thel. hirsuta. δ . *decipiens*, effusa resupinata glaberrima lævigata aurantiaca.

In truncis horizontaliter dejectis.

Obs. Facillime pro distincta haberi posset specie.

Th. laciniata. β . *permutata*, effusa resupinata, papillis marginalibus sparsis.

195. *Thelephora corrugata*, effusa erectissime adnata tenuis rimosissimo-corrugata lurida, papillis nullis.

In ligno decorticato.

Obs. Facile rugis suis, a rimis ortis, distincta species, sed color difficiliter determinatur. Proxime tamen accedit *luridum*, quale in *K. Vec. Ac. Handl.* 1813 t. 4. f. 11. Cl. *Billberg* delineavit. Irregulariter effusa, 2—3 unc. glabra.

196. *Thelephora Pini*, effusa duriuscula subtenuis rimoso-partita glabra ferugineo-fusca, papillis irregularibus.

Theleph. Pini. Schleicher.

Ad truncos *Pini sylvestris* in Helvetia, Suecia.

Obs. Specimen, quod ex Helvetia possideo, a Suecico non differt. Ceterum frustulis tenuibus, glabritie, colore &c. a reliquis affinibus facile distinguitur.

197. *Thelephora evolvens*, junior subrotunda clausa, dein evolvens tubcupulæformis, extus margineque pallida tomentosa, disco glabro obscuriori. Tab. 10 f. 1 a. b.

In ramis & truncis betulinis rarissimam & distinctissimam speciem invenit mecumque liberalissime communicavit ill. Prof. & Eques *Acharius*.

Descr. Vera omnino *Thelephora* species; sed more crescendi a reliquis quam maxime differt, nec alii quidam mihi notæ affinis. Primo tuberculum subrotundum convexum villosum pallidum sistit, dein magis magisque evolvitur & capularis fit $\frac{1}{2}$ —1 unc. disco dilatato glabro subincarnato livido, primo laevi, dein subrugoso; extus margineque involuto pallide tomentosa. Papillas veras non vidi.

STEREUM.

LINK.

in *Berl. Mag.* 1809 p. 39. f. 32.

198. *Stereum rubiginosum*, imbricatum rubiginoso-spadiceum subglabrum, papillis majusculis.

Thelephora rubiginosa. *Pers. et Auct.*

Singularem hujus legimus varietatem quæ, cel. *Achario* per litteras monente, forte est *Thei. fuliginosa* Schleich. & diximus.

ß. corriciatum, effusum resupinatum papillosum confluens concentrice zonatum.

Ad truncos quercinos ad terram horizontaliter dejectos.

MERISMA.

192. *Merisma clavulare*, erectum coriaceum rufo fuscum inodorum pulverulentum, basi subtuberosum, sursum dilatato-ramosum, ramulis subconcoloribus. Tab. 1. f. 1.

In silvis abiegnis Sueciæ. Ad Åker Amic. *Hesflin*. Ad Femsiö Ipse.

M. anthocephalo Swartz in *K. Vet. Ac. Handl.* 1811. p. 84 affine; sed diversum videtur. (Conf. *Ic. Bulliardi* cum nostra) $1\frac{1}{2}$ — $2\frac{1}{2}$ uncial. fusco-ferrugineum, basi subtuberosum. *Stipes* deorsum angustatus subteres sensim compressus digito-ramosus. *Rami* pulvere concolori conspersi complanati inæquales, apice sublaciniati concolores.

CLAVARIA. LINN.

200 *Clavaria suecica*, ramosissima erecta ochraceo-incarnata opaca, caule crassiusculo deorsum albido-tomentoso, ramis multifidis acutis.

In pinetis abiegnis muscosis Sueciæ, sæpe frequens.

Descr. *Gregaria ochracea* L. ochraceo-rufescens. *Caulis* crassiusculus (non tamen ad crassicaules pertinet) solidus mollis brevi sculus æqualis, basi tomento fibrilloso a) b) do radicans, 3—4 lin. crassus erectus numquam decumbens. *Rami* 1—1½ unc. laxiusculi flexibiles suberecti glaberrimi solidi, intus aquosi, multifidi acuminati.

Obs. — *Cl. gracili* P. proxima; sed vere distincta.

201. *Clavaria byssifeda*, varia ramosa albida, radiculis byssoideis concoloribus insidens. *Cl. byssifeda*. Pers. Syn. p. 593. — Obs. myc. l. p. 39. cum. descr. breviss.

Ramaria fimbriata. Holmsk. ot. p. 98. cum ic.

In ligno dejecto quercino.

Descr. Species valde polymorpha & habitu singulari. Junior *Himantia alba* similis; sed ramulis furcatis spinulosis differt. *Truncus* non quidem crassus; sed ramis simul sumtis sæpe crassior, albus subglaber, basi radiculis albis byssoideis pubescentibus insidens. *Rami* stricti recti albi breves subsimplices acuti.

Meum fungum numquam rufescentem vidi.

202. *Clavaria rugosa*, ramosa albida, caule tenui, ramis simplicibus rugosis difformibus incrassatis.

Clav. rugosa. *Pers. Syn. p. 594. Comm. p. 57 cum syn.*

In pinetis Femsiönensibus; satis frequens in Suecia.

Descr. Majuscula albida gregaria variabilis, junior subsimplex. *Caulis* brevis angustus (ramo singulo subangustior.) glaber solidus sublævis. *Rami* pauci difformes sæpe connati incrassati glabri rugosissimi, juniores læviores, solidi 1—2 unciales subobtus. Substantia mollis.

De numerosis hujus varietatibus conf. II. ec. Hujus alia varietas lævior est *Cl. rugosa* Fl. Dan. t. 1301. quæ in nostris silvis cum *δ. hercynica* P. crescit.

Optimam Clavariarum Suecicarum monographiam dedit Cel. *Swartz in Vet. Ak. Handl. 1811.* Desiderantur species descriptæ, *Cl. helveola*, quam legi in ericeto ad Jonstorp in Hallandia, *Cl. palmata* &c.

GEOGLOSSUM P.

203. *Geoglossum difforme*, nigrum viscosum, clavula nitente compressa demum curvata difformi.

In pinetis locis graminosis. Sept. Octbr.

De fr. Gregarium interdum subcaespitosum 2—unciale totum nigrum viscosum, nec villosum l. squamulosum.

Stipes $\frac{1}{2}$ unc. subcompressus, non elasticus.

Clavula nitens complanata obtusi subcurvata difformis stipiti longior laevigata, intus cinereo-nigricans, a stipite subdiscreta.

Clavula junior subcanaliculata.

A *Geogl. glutinoso* & viscoso in Nov. Fl. Succ. l. c. descriptis, abunde differt.

Non in omnibus *Geoglossi* speciebus stipes a pileo discretus, ut apud *Link* l. c. p. 40; ex. gr. in *G. viscoso*. &c.

MITRULA PERS.

disp. meth.

Pileus stipitatus (*Leotia Link.*) ovatus l. cylindricus undique thecigerus, stipitem distinctum aretissime ambiens.

204. *Mitrula Todei*, stipite elongato capillari albido glabro, pileo sublineari flavo.

Chordostylum *Clavaria*, erectum tenacissimum simplex, stipite capillari glabro flexuoso, capitulo pyriformi coriaceo. *Tode Fung. Meckl.* l. p. 37. t. 6. f. 55.

In stipibus locis humentibus dejectis, præcipue in Filicibus & frondibus *Pteridis aquilinae*. Aug. Septemb.

Obs. Vera omnino me judice, *Mitrulae* species, quamvis habitu peram recedat. Fungis mycoroideis nullo modo adnumerari potest; sed de synonymo *Todei* vix dubius sum.

Vix hujus loci reliquæ *Chordostyli Todei* species.

Reliquæ species Suecanæ sunt *M. Hederi*, *Ludvigii*, *Dicksonii* & *Bulliardi*, quæ vero tres ultimæ nimis affines.

205. Characteres Generum affinium: —

1. *Mitrula Pers.* disp. meth. Fr. (*Leotia Link.* *Leotia* sect. 1ma *Pers. Syn.*)

Vide supra.

2. *Leotia Pers.* — Fr. (*Leot.* sect. 2da *Pers. Syn.* *Helotii* species *Link, Swartz.*)

Pileus stipitatus obiculis plano-convexus margine revolutus, interne thecigerus.

Ab Helotiis abunde differt fructificatione inferna, a qua pagina sporulæ fumi instar avolantes nudo oculo videri possunt.

Species Suecane sunt: *L. circinans* *P. lubrica* *P. β. tremula, marcida* *P. & truncorum* *Alb. & Schwein.*

- 3) *Verpa Swartz* (Leot. sect. 3tiæ *Pers.*)
Pileus stipitatus lavis æqualis conicus inferne thecigerus.

Inferne etiam fructificat, sec. observationes
Ipsius *Swartzii*.

Spec. Suec. *V. candida* *Sw. & conica*,
(*Phallus conicus* *Fl. Dan.*)

RHIZINA. FR.

Pileus sessilis bullatus convexus, supra thecigerus, subtus concavus fibrillis pluribus radicatus (*unde nomen.*)

206. *Rhizina undulata*, undulata fusca subtus pallida.

Helvella acaulis. Pers. Syn. p. 614. Obs. myc. 2 p. 20. Alb. & Schwein l. c. p. 298.

H. inflata. Schæff. fung. t. 153

In silvis sabulosis & deustis in Smolandia ad Femsiö &c.

Caro fragilis albida. Subtus concava albida læviuscula glabra, fibris pluribus crassiusculis simplicibus albis radicata.

207. *Rhizina lævigata*, convexa læviuscula, subtus pallida.

Octospora Rhizophora. *Hedw. Descr. musc. frond.* 11 p. 15 t. 5 f. a.

Peziza rhizophora. *Willd. Ber. n.* 1170. *Phallus acaulis Batsch.*

In silvaticis; etiam in *Suecia*.

Obs. Priori valde affinis. Utriculi (theæ) in mea non ita prominent, ut in *icone Hedw.*

Obs. Medium Genus inter *Helvellam* et *Pezizam*; his vero propius accedit, ut jam monuit *Cel. Link*; sed distinctum genus.

PEZIZA. LINN.

I. *Burcardia*. Gelatinosæ s. *Tremellosæ*.

Obs. Proprie ad hanc sectionem tantummodo referre vellem *P. inquinantem* *P.* (*Tr. Agaricoid.* *Retz.*) *P. Burcardiam* *P.* (*Lycoperd. truncatum* *Linn. sec. Hall in Vet. Ac. Handl. 1812.*) *P. puram* *P.* (*Fries Nov. Fl. Suec. p 21*) & *clavum* *Albert.*; reliquas tamen ut heteroclitas & intermedias heic militantes reliqui. *Pez.*

tamen *Gelatinosa* P. (Tr. *sagarum* Retz) & *sarcoidis* P. ad Tremellas venosas forte potius reducendæ. Plurimas novas species possideo; sed alio loco cum iconibus illustranda.

208. *Peziza viridi-nigra*, majuscula carnofo-gelatinosa virescenti-nigra subglobosa, disco cupulari concolori.

P. atrovirens. *Summa Veg. Scand.* p. 63. (nec. Persf.).

In ligno pineo putrido autumnno pluvioso, rara.

a *P. atrovirente* differt: Substantia firmiori, magnitudine, colore magis opaco subnigro, forma globosa urceolato-cupulæformi: intus concolor est.

209. *Peziza rubella*, minuta carnofo-tremelloso-rubra subconvexa, margine demum flexuoso-ferrulato.

P. rubella. *Persf. Syn.* p. 635.

In ligno putrido fagineo legi. Rarior Fungus.

Descr. Gregaria sessilis tremelloso-minuta tota glabra pallide incarnato-rubra, subtus planiuscula nuda lævis, supra convexa, margine primum integro dein ferrulato flexuoso.

II. *Aleuria*. Extus farinosa s. *Helvelloidei*.

210. *Peziza badia*, magna sessilis subflexuosa intus obscure brunnea, margine extusque dilutior, basi farinosa.

P. badia. *Perf. Obs. myc.* 2 p. 78. *sine descr.* *Syn.* p. 639.

P. cochleata. *Bolt. Fung.* t. 99. (bona).

In silvaticis locis graminosis *Suecia*, *Agliæ* *Germaniæ* &c.

Descr. Pulchra species; omnino digna, quæ describatur. 1—2 unc. lata, $\frac{1}{2}$ —1 unc. alta, cupularis difformis subflexuosa intus glabra lævis obscure brunnea l. cinnamomeo-rufa, margine inæquali in junioribus subinflexo, extus dilutior subglabra & basi tantum farinosa subtilissime villosa (sub lente).

P. fascicularis Alb. & Schwein an ad sect. V referenda? crescit in *Suecia*.

III. *Lachnum*. Extus villosa.

Pezizæ hujus *Familiæ* hoc caractere facile distinguuntur.

V. *Dermia*. Coriaceæ glabræ.

211. *Peziza mycophila*, gregaria minuta planiuscula pallide virescens. *Nov. Fl. Suec.* p. 21.

In Agaricis putrescentibus ex. gr. *Ag. adu-*
sto &c.

Descr. Coriacea sessilis minuta glaberri-
ma, junior concava viridis margine æquali inte-
gro; dein planiuscula virefcenti-albida subcon-
fluens difformis.

212. *Peziza leucostigma*, gregaria mi-
nuta alba disco subcinerascenti.

In ligno nigricante fagineo, in quo ut
granula alba apparet.

Descr. Sessilis coriacea minuta grega-
ria concava junior punctiformis, dein subappla-
nata, alba disco subcinerascente, exsiccata sub-
flavescit, margine orbiculari integro $\frac{3}{4}$ — $1\frac{1}{4}$ lin.
lata.

213. *Peziza abietis*, caespitosa rugosa
opaca sordide nigra rotunda aut subcompressa.
oris inflexis. *Perf. Syn.* p. 671.

Quam in ramis exsiccatis *pineis* legi, pa-
rum differt, hinc describenda:

Cæspitosa coriacea sessilis subglabra opaca
nigra. *Cupula* subcompressa rugosa difformes
subconfluentes, oris integris conniventibus. Cæ-
spit. 2—3 lin.

Obs. Numerosissimas ceterum Floræ Sue-
cicæ Novitias ex hoc Genere legi, in *Vet. Ac.*

Handl. 1812 non allatas. Ex. erint *Peziza brunnea* Alb. & Schw., *P. pura*, *P. gelatinosa* *P. sarcoides* *P. turbinata* Vahl, *carnosa* V. *aurea* *P. applanata* Hedw. *macropus* *P. carbonaria* Alb. & Schw. *fascicularis* Eorund. *villosa* *P. clandestina* *P. Rosa* *P. marginata* Schum. *Phiala* Vahl, *ciborium* Vahl, *tuba* Bolt. *faginea* *P. confluens* *P. umbonata* *P. atrata* *P. compressa* *P. hystereum* *P. populnea* *P. prunastri* *P. truncatula* *P. Verrucaria* Alb. & Schw. *subhirsuta* Sch. *flava* Schum. &c.

214 *Peziza lecideola*, minuta gregaria subconfluens concava nigra, crustae cinerascenti insidens. Tab. IV. f. 2. a. b.

In Helvetia ad terram supra folia pinea dejecta.

Descr. *Cupulae* minutae punctiformes glabrae concavae margine subobtusō, in junioribus æquales, dein subconfluentes irregulares, omnino pezizoides; sed crustae effusae cinerascenti tenui, a solo, cui innascitur, inæquali insidentes & hinc *Lecideis*, quam maxime affinis; sed ad has non referendam esse, docuit summus Lichenologus Cel. *Acharius*.

215. *Peziza xanthostigma*, minuta

subgregaria orbicularis concava flavescenti-aurea coriacea glabra.

In ligno *Pini Abietis* putrido in Polonia.

Cupulis tenuissimis coriaceis minoribus margine orbiculari integerrimo a *Pez. aurea*, cui proxima, & colore cupulisque magis concavis a *P. leuco-stigmate* distincta.

† PHACIDIUM. FR.

Pileus innato-fessilis subplaniusculus epidermide (*nigra*) demum rupta involucratus, thecis oblongis subprominentibus compositus.

216. *Phacidium coronatum*, margine epidermidis ruptae crenato, disco pallide flavescenti.

Ascobolus coronatus *Schum. Sell.* 2 p. 437.

Xyloma Pezizoides. *Pers. Syn.* p. 105.
Is. pict. t. ¹⁸~~16~~ f. 1.

Peziza comitalis. *Sov. Fung.* t. 118.

P. viridis. *Rolt. Fung.* t. 119. f. 1.

Sclerotium quercinum. *Fl. Dan.* t. 1380.

In foliis, dejectis *Quercinis*, *Fagineis*, *Betulinis*, *Alneis* &c.

Fungus distincti Generis, hinc nunc ad hoc, nunc ad illud est relatus. A *Xylomatis* propriis (vide *Link in Berl. mag.* 1. c. p. 35)

omnino differt & cum thecæ in superficie demum sunt collocatæ & sporidia in his seriatim disposita ad fungos proprios (*Link l. c.*) referri debet. Ex his, ut optime jam observavit Cel *Schumacher* Ascobolis proxime accedit; sed pluribus characteribus & toto habitu differt. Ad Scieromycos tamen transitum efficit.

217. *Phacidium integerrimum*, margine epidermidis ruptae integerrimo, disco palide lutescenti.

In caule plantæ cujusdam aridæ semel legi.

Deser. Priori affine; sed minus, margine epidermidis integerrimo cinctum & hinc quasi patellatum. Color præced.

Hujus Generis etiam forte aliæ Xylomatis species Auct. ex gr. *X. salicinum* P. disco concolori, epidermide rupta, sporidiis fumi instar avolantibus &c.

SPHÆRIA. P.

II. *Peripharica*.

218. *Sphæria repanda*, subcupulæformis nigra, subtus corrugata, margine repando, stromate albo, sphaerulis papillatis. *Tab. I. f. 5. a b. c.*

In fruncis sorbeis.

De ser. Nulli alii speciei affinis. Gregaria; sed non confluens, subcupulæformis oblonga $\frac{1}{2}$ —1 unc., extus nigra, subtus quam maxime corrugata, supra laevis; sed papillis sphaerularum tuberculosa. *Sphaerulae* periphericae (in cupulam supernam) oblongo-lineares, stromato albo suffultæ.

219. *Sphæria connata*, turbinato-substipitata cinerascens mox nigra, supra planiuscula sphaerulifera, dein latissime confluens.

In fago cava copiose legi.

De ser. Suberosa gregaria, junior subcinera, mox nigra, 2—2 $\frac{1}{2}$ lin. alta, supra plana 2—4 lin. lata, turbinato-substipitata, stipite crasso brevi, dein vero plurima specimina plane confluunt & stipes fere evanescit, ad marginem tamen Sphaeritæ hujus latæ effusæ videtur. *Sphaerulae* majusculæ prominentes paucae in disco superiori positæ.

Obs. Hæc & antec. species a reliquis hujus sectionis differunt: sphaerulis tantummodo supernis; sed hæc a modo crescendi pender.

220. *Sphæria multiformis*, subeffusa varia opaca fusco-rubiginosa intus obscure nigra, cortice non disseparabili, sphaerulis prominentibus.

Sphæria argillacea. Pers. ic. pict. t. 3.
f. 1—3 (quoad ic.)

Polymorpha; sed numquam globosa, ut
Sphæra fragiformis.

Varietates præcipuæ sunt:

α. *Sph. coarctata*, erumpens gregaria convexa elongata utrinque attenuata rufo-fusca, sphærulis sublatentibus. Tab. I. f. 2. a. b. c.

Ex epidermide *Berule albe* transversim erumpit, forma lanceolata distincta, locis vero decorticatis fit.

β. *dilatata*, superficialis effusa confluens planiuscula, sphærulis prominentibus. Tab. I. f. 3.

In truncis betulinis.

Ab his magis recedit:

γ. *alnea*, erumpens crassa varia subnigrescens, sphærulis prominentibus papillatis.

In truncis alneis cortice crasso tectis.

Forma in omnibus varietatibus a loco, cui innascitur, pendet.

Diagnosis *Sphæriæ fragiformis*, subglobosa confluens miniato-rubiginosa intus atro-nitens, cortice disseparabili (*decidua*), sphærulis papillatis. — *Sph. rubiginosa*, in truncis fagineis Sueciæ vulgaris, huic vix affinis. —

221. *Sphæria argillacea*, subglobosa, argillacea l. lutescenti-cinerea, intus fuscescenti-nigra, sphærulis prominentibus papillatis. *Tab. 11. f. 5. a. b.*

Sph. argillacea. Pers. syn. p. 10. Alb. & Schwein. l. c. p. 4.

In truncis betulinis legi.

Obf. *Sph. fragiformi* & *multiformi* maxime affinis & colore præcipue differt. Sphærulæ ut in *multiformi*. Cum vero Icon Persooniana (*l. pict. t. 3. f. 1—3*) colore a nostra omnino recedat, colore & magnitudine naturali depictam sistit *Tab. 1. f. 4. a. auctam. f. 4. b.*

222. *Sphæria rhodogramma*, erumpens subseriato-confluens convexiuscula rosea, sphærulis latentibus. *Tab. 1. f. 4. a. b.*

In truncis dejectis fagineis.

Descr. Pulcherrima species, cum *Sph. rhodogramma* P. comparanda, cui tamen vix affinis. Erumpens suberosa rosea subpulverulenta intus nigricans, junior subrotunda 2—3 lin. longa, dein seriato-confluens $\frac{1}{2}$ —1 unc. subirregularis magis explanata convexiuscula. Sphærulæ periphæricæ, ostiolis latentibus.

III. COMPOSITÆ.

223. *Sphæria Mezerei*, erumpens orbicularis planiuscula intus concolor, ostioliis papillatis prominentibus.

Sph. Mezerei. Schleich. sec. spec. Helv.

In ramis emortuis *Daphnidis Mezerei*, in Helvetia, Suecia (*Forsander.*)

Obs. *Sphæriæ disciformi* nimis affinis, in qua ostiola, negante licet *Persoonio*, etiam sæpe prominent; sed hæc minor est, ostioliis confertioribus & substantia interna nigra.

224. *Sphæria rubra*, epiphylla orbicularis rubra, margine glabro, ostioliis subimmersis obscurioribus.

Xyloma rubrum. *Pers. syn. p. 105. Obs. Myc. 2. p. 101. Schum. Sæll. 2. p. 179. Alb. & Schw. p. 64.*

In foliis *Pruni spinosæ & domesticæ* æstate & autumnno frequens.

Vera omnino *Sphæriæ* species & *Xylomatis* l. *Æcidiis* nullo modo affinis, ut omnes voluerunt Auctores.

225. *Sphæria Xantha*, epiphylla subirregularis ochraceo-flava, margine glabro, ostioliis immersis obscurioribus.

Sph. ochracea. *Wahlenb. Fl. Lapp.* p. 518
sec. spec. Lapp. (excl. syn. *Perf.*)

Xyloma aurantiacum. *Schleich. sec. spec. Helv.*

In foliis *Pruni Fadi* frequenter.

Priori nimis affinis; differt tamen forma minus orbiculari, colore & ostioliis rarioribus magis immersis.

Ne cum *Sph. ochracea* P. l. *aurantia* Ejusd. confundatur, nomina Auctorum trivialia mutare coactus sum.

226. *Sphæria Ulmi*, epiphylla convexa nigra, ostioliis numerosissimis papillatis.

Sphæria Ulmi. *Schleich. exsicc.*

In foliis *Ulmi campestris* dejectis putridis, in *Suecia*, *Helvetia*.

Obs. Vix alii affinis. Sub epidermide folii crescere incipit & hac tecta cinerascens apparet. Habitus fere periphæricarum; sed intus cava & stromate caret. Hinc ad sect. 4tem pertineret. 2—3 lin. lata.

Specimina Suecica, a. Cel. *Swartzio* liberalissime missa, ab Helveticis nullo modo differunt.

IV. MONOSTICHÆ.

227. *Sphæria atro-purpurea*, effusa applanata atro-purpurea, sphærulis ovatis, ostiolis papillatis.

Sph. atropurpurea. Tod. Fung. Meckl. 2. p. 32. t. 13. f. 106. a—d. Rehent. fl. Neom. p. 324.

In ligno fagineo putrido in Suecia meum legi fungum.

Distinctissima quidem species, quam *Sph. rubiginosa* varietatem esse suspicatus est Cel. *Persoon* & *Sph. stigmatis* Cel. *Schumacher*. Neutri tamen affinis, vixque Cl. *Tode* Ipse fungum perfectum vidit. Ad latitudinem 3—5 unc. effusam legi, colore plus minusve atro-purpureo, sphærulis adultis prominulis, apice planiusculis papillatis.

Sph. Maculae Tod., in ligno fagineo Sueciæ etiam crescenti, proxima quidem, sed omnino distincta; nec hæc cum *Sph. stigmati* P. conjungenda, ut voluit Cel. *Schumacher*.

228. *Sphæria Ostruthii*, epiphylla minuta difformis, sphærulis crustæ cinerascenti immerfis subconfluentibus.

In averla parte foliorum *Imperatoria Ostruthii*.

Sphæria Anethi vere affinis & forte non differt *Sph. Podagrariæ* Roth. Nudo oculo maculam nigricantem & irregularem sistit; sub lente vero crusta tenuis cinerascens, cui sphaerulae nigricantes subimmerse, apparet.

V. TUBERCULATÆ.

229. *Sphæria extensa*, subrotunda nigra, stomati late effuso cinereo-nigro innata, ostiolis oblongis opacis depresso perforatis. Tab. 3. f. 2. a h.

Sph. extensa. Achar.

In ramis arborum, rarior. (in *Sorbo*).

Species optime distincta.

VII. CÆSPITOSÆ.

230. *Sphæria elongata*, caespitibus inæqualibus seriatim undique confluentibus, sphaerulis nigris disco planiusculis annulatis papillatis, intus concoloribus faretis.

In ramis dejectis fagineis, primo vere.

Descr. Pulchra species. *Cæspites* maxime polymorphi. rarissime subrotundi æquales distincti, plurimi seriatim erumpentes longissime undique confluentes, & ramulos sæpe an-

bientes. *Stroma* (receptaculum P.) tenue nigrum.

Sphærule globosæ nigræ solidæ læves, apice depresso. planiusculæ linea depressa annulata subpapillatæ.

Leaturn. 231. *Sphæria rufo-fusca*, cæspitibus subrotundis æqualibus, sphæruleis rufo-fuscis intus albidis cavis depresso-umbilicatis. Tab. IV f. 3 a—d.

Sphæria berberides. *Schleich. sec. spec. a cel. Achario miss.*

In ramis exsiccatis, *Berberidis* forte, in Helvetia.

Descr. *Sphær.* appendiculatæ proxima & *Sph. berberidis* vix affinis. Cæspites subrotundi distincti haud magni convexi 2—3 lin. *Stroma* tenue nec appendiculatum glabrum. *Sphærule* subrotundæ læves cavæ umbilicatæ rufescenti-fuscae intus albidæ. — Variat cæspit. confluentibus.

232. *Sphæria conglobata*, cæspitibus subglobosis minutis, sphæruleis solidis nigris ovatis astomis subrugulosis.

In ramis betulinis, rarior.

Descr. Inter minores hujus sectionis & tuberculatis sat similis; sed sphæræ liberæ

sunt. *Cespites* $1\frac{1}{2}$ —3 lin., e sphaerulis valde prominentibus, subglobosi nigri aequales distincti. *Stroma* subconvexum tenue. *Sphaerulae* minutae astomae ovatae l. globosae opacae tenuissime corrugatae.

VIII. SIMPLICES.

I. MACROSTOMÆ.

Sph. platystomas P. sub *Lophio* retuli.

233. *Sphaeria excipulaeformis*, sphaerulis nudis ovatis nigris laevibus ostiola cylindrica æquantibus. *Tab.* IV. f. 5. a—c.

In cortice arborum frondosarum *Aceris*, *Fraxini* & Æstate.

Obi. Forma ovata, ostiolo brevi crasso (sphaerulis etiam nonnumquam breviori) & loco natali a reliquis hujus sectionis facile distincta. Proxima *Sph. biformis* β, in qua tamen sphaerulae globosae sparso pilosae & ostiola conica, capsula breviora.

234. *Sphaeria capillata*, sphaerulis subnudis minutis ovatis nigris, ostioli rectis setaceo-capillaribus.

In fungis suberosis, ex. gr. *Boletis pini cola*, ad terram dejectis.

Defer. *Sphærule* nunc nuda, nunc subimmerfa ovata nigra nitida duriuscula lævis, bafi omnino glabra, minuta vix 1 lin. lata, *Oftiolo* longiffimo fetaceo concolori fubrecto; fed minime ftricto.

235. *Sphæria rostellata*, fphærule congestis nigris fubrotundis fupra depreffis umbonatis, oftiolis cylindricis obtufis hifpidis. *Tab.* III. f. 3. a—e.

In caulibus dejectis. *Acharius*, qui nomen & iconem dedit.

Species polymorpha. *Sphærule* primum hemifphæricæ oftiole papilliformi (f. b), dein vero elongantur (f. d.) Variat etiam oftiolis pyramidatis (f. c) & collapsis (f. e.)

2. BRACHYSTOMÆ (♂ *astomæ*).

236. *Sphæria rhodomelas*, fphærule confertis fubrotundis patellatis nigris, cruftæ tomentofæ rubræ infidentibus.

In trunco *Pyri Mali*.

Sph. rostellæ Alb. & Schw., quam femel tantum in fuecia vidi, valde affinis. Crufta eadem; fed *Sphærule* maxime differunt & in *S. rhodomel.* funt patelliformes, margine integerrimo, nigræ &, ni fallor, *astomæ*.

237. *Sphæria nigrella*, spherulis ovato-globosis lævibus nigris, macula elongata concolori innatis, ostiolis distinctis perforatis. *Tab. IV. f. 4. a—c.*

In caulibus plantarum ex. gr. *Galeopsidis* &c. rarior.

De scr. Macula elongata $\frac{1}{2}$ —1 unc. tenuissima glabra lævis. *Spherule* oblongæ basi applanatæ, (hinc ovato-globosæ) nigrae nitidæ læves glabræ ostiolis pertusis l. umbilicatis. (f. c. dissecta.)

Cum *Sph. herbarum* P. aliisque in macula heterogena parasiticis non confundenda.

238. *Sphæria malina*, spherulis sparsis minutis rubris astomis, maculae aurantiacæ immerfis. *Tab. II. f. 3. a. b.*

In pomo *Pyri Mali* vivo semel tantum vidi.

Macula $\frac{1}{4}$ — $\frac{1}{2}$ unc. lata tenuissima, sensim dilatata. *Spherule* immerse & parum tantum prominent, oblongo ovatæ roseo rubrae.

239. *Sphæria palmicola*, gregaria nigra nitida glabra convexa, demum umbilicato-perforata.

In epidermide fructus *Coccoes nucifera* ex India Orientali allati.

Descr. Inter minores, subimmersa. Color præcipue sub lente nitidus. Irregularis est & hinc diagnofin e speciminibus perfectis defumfi. Eisdem locis maculæ fordidæ nigricantes opacæ observantur, quæ caute deftinguendæ.

240. *Sphæria refina*, sphæruis subimmersis globosis glabris lævibus aurantiaco-rubris astomis.

In refina Pini Abietis. *Swartz.*

Obf. Minuta punctiformis subfolitaria, humectata gelatina albida farta. Non confundenda, cum:

Lecidea refina, crusta tenuiffima membranacea contigua cinerascenti, apotheciis sparsis sessilibus concavis flavo rubris.

Quam in eodem loco ad Femfö legi.

241. *Sphæria? crispa*, gregaria subconfluens, sphæruis convexo-applanatis crispato tuberculosis astomis nigris.

Sph. crispa. *Achar. cui spec. debeo.*

Ad corticem arborum in Lusatia. *Mosig.*

Obf. De loco dubius sum. Nifi Lichenis species, quod negat Lichenologus summus; sphæriis proxima. 1—2 lin. & plures demum confluunt.

242. *Sphæria umbonata*, gregaria subconfluens, sphærulis convexiusculo-applanatis lævibus nigris papillatis.

In ligno fagineo putrido.

Descr. Tota nigra (etiam intus) orbicularis ligno adplanata disco convexiusculo papillato. Papilla minuta punctiformis.

243. *Sphæria applanata*, sparsa, sphærulis concaviusculo-applanatis nigris intus concoloribus astomis, disco demum subrugoso.

In lignis emortuis.

Descr. Primo obtusa. Pezizæ cuidam planæ similis. Astoma intus & extus nigra junior tenuissime concava submarginata lævis; dein vero convexiuscula immarginata atra, margine orbiculari lævi, disco subruguloso.

244. *Sphæria sepincola*, gregaria, sphærulis subrugosis nigris intus subalbidis cavis, ostioli minutis impresso-punctatis, sæpe nullis.

In caulibus *Rosarum* emortuis. Swartz.

Ipsæ.

Obs. *Sphær. herbarum* forte proxima; sed sphærulis cavis, ostiolo &c. differt. Sub epidermide crescit & hinc junior testæ. In nonnullis observavi ostiola subperforata, in aliis nulla, quod cum multis aliis commune, quam

ob causam Astomas cum Brachystomis conjunxi; etiam forte paucissima l. nulla vere astoma sunt, quamvis ostiola nondum detecta. Hæc disponendi methodus species maxime affines, ex gr. *Sph. rosellam* & *rhodomel.*, nimis etiam removet.

245. *Sphæria Lonicerae*, aggregata, sphærulibus globosis superficialibus nigris intus albicantibus, disco depresso pezizoideo.

In caulibus aridis *Lonicerae Peryclymeni*.

Deser. Affinis *Sph. patella*; sed abunde diversa: forma subglobosa, disco pezizoideo profunde depresso, margine obtusissimo orbiculari subconnivente, disco laevi astoma, strato interno tenuissimo albicaute, magnitudine seminis Papaveris.

246. *Sphæria mycophila*, subconfluens, sphærulibus convexis superficialibus fragilibus nigris subnitidis papillatis, ostiolis oblongis dein deciduis perforatis.

In fungis suberosis putridis.

A *Sph. pulveracea*, *pertusa* aliisque firmo caractere ægre distinguitur. His tamen nullo modo affinis. *Perithecium* nigrum nitidum læve glabrum convexum tenuissimum fragillimum intus cavum.

247. *Sphæria pithya*, gregaria varia; sphærulis subrotundis nigris intus concoloribus solidis lævibus superficialibus astomis.

In cortice *Pini sylvestris*, vulgaris.

β. *Strobilina*, polymorpha junior subapplanata, rugulosa.

In fibribus dejectis *Pini sylvestris*, frequens. Forma irregulari & loco natali præcipue distincta species. Inter minimas.

248. *Sphæria Hederae*, epiphylla sparsa, sphærulis superficialibus convexis nitidis nigris, ostiolo albo distincto pertuso. *Tab. IV. f. 6 a—c.*

In foliis *Hederae Helicis*. Vere.

Descr. Pulcherrima. *Sphærule* innatae orbiculares convexae glabrae laeves nigrae nitidae. *Ostiolum* distinctum album pertusum, subfarinulosum; sed in junioribus subobsoletum. —

Sphærule demum magis applanatae, ostioli umbilicatis nudis subnigris.

In foliis *Hederae* etiam vigent: *Sphæria Dematium* β. P. *Mucor Dematium* Fr. *Sphæria punctif.* β. P. &c. (In horto Lundensi)

249. *Sphæria Taxi*, gregaria minuta, sphærulis convexis subimmersis nigricantibus, ostiolo truncato albo.

In ramulis & foliis *Taxi baccatae*, per totum annum; sed specimina ostioli instructa vere tantum vidi.

Affinis priori; sed omnino distincta. Multo minor & color nudo oculo cinerascens.

Non confundendum cum *Xylomate Taxi*. Fr. a quo ostiolo truncato albo, nec pertuso ut in præc., differt.

Sph. Hederae, Taxi, atrovirens Alb. & Schw., quæ omnes in horto Lundensi adfunt, tribum naturam efficiunt.

250. *Sphæria parella*. Tod.

Hujus, in caulibus herbarum vulgaris, varietatem vidimus, quæ:

β *Xylaria*, minor, disco lævi, non plicato, papillato.

In lignis exsiccatis duris, rarior.

251. *Sphæria ocellata*, sparsa, sphæru-
lulis plano-concavis opacis nigris, margine orbiculari albo cinctis.

In ramis *Betulae nanae* emortuis exsiccatis in Smolandia.

Epidermidem subelevat & in ambitu albam reddit, qua nota facile distinguitur hæc species. Margo vero albus ab incauto pro *Sphæria* & *sphærule* nigra pro ostiolio haberi posset

quo facto vix differret *Sph. melastoma*. Re-
bent. — Sæpius astoma, & papillam mini-
mam in disco observare semel tantum mihi
contigit.

252. *Sphæria speirea*, sparsa, sphæ-
rulis planis immerfis opacis nigris orbiculari-
bus immarginatis astomis, in foliis vivis para-
fiticis.

In foliis vivis *Ægopodii Podagrariæ*.

Cum *Sph. Ægopodii* P. & *Podagrariæ* Roth.
loco convenit; neutri vero affinis. Macula
enim nulla & sphærulæ sparsæ planæ, qua nota
etiam a *Sph. maculiformi* differt. Differunt
quoque:

1. *Sph. punctiformis*, sphærulis subniti-
dis, disco subumbilicatis, loco in foliis emor-
tuis &c.

2. *Sph. applanata*, sphærulis 3 — plo &
ultra majoribus superficialibus submarginatis,
disco demum ruguloso, loco &c.

3. *Sph. Glechomæ* Schleich. (sec. spec.
Helv) sphærulis majoribus rotundis superficia-
libus, minime nigricantibus.

253. *Sphæria alnea*, epiphylla sparsa,
sphærulis minimis convexis nigris opacis, osti-
olo albo, demum collapsis concavis.

In foliis putridis *Betulae Alni*.

Folia, quibus hæ sphaerulae minimae innascuntur, subalbicant. — Sub Microscopio determinanda & nihilo tamen minus characteres difficillime ex his atomis eruuntur.

Ostiola tantum in adultis adsunt. Magis a *Sph. Hederae* & *Taxi* differt, quam e descriptione crederes.

254. *Sphaeria macularis*, epiphylla, sphaerulis minutissimis subrotundis atomis nigris, maculae cinerascenti tenuissimae insidentibus.

α populnea, macula cinerascenti. *Tab. II.*
f. 4. a. b. c.

In foliis populneis dejectis.

Sphaerulae nudo oculo inconspicuae fere.

β grisea, macula grisea subradiata.

In foliis betulinis.

Affinem speciem, in foliis *Allii* crescentem, nuper misit Cel. *Swartz*, Fautor summus; sed magnitudine maxime diversa. (*Sph. allicina*)

NÆMASPORA. P.

255. *Næmaspora hyalosperma*, tuberculo tenui, cirrhis hyalinis.

N. hyalosperma. Achr.

Ex America boreali, ad ramos crescens.

Obs. Inter minores species, tuberculo tenui epidermide corticis, parum bullatim elevati, rupta cincto. Cirrhi in specimine meo nulli adfuerunt.

256. *Næmaspora melasperma*, tuberculo tenui, cirrhis solitariis nigris crassiusculis.

N. nigra. Rehbent. fl. Neom. p. 340

In ramis betulinis emortuis fungum meum sat frequenter legi.

Obs. Tuberculum, licet tenue, adest.

A. *N. juglandis* & *faginea* Schum. cirrhorum forma & a reliquis colore differt.

N. grisea a *N. leucosperma* parum differt.

SPÆRONEMA.

Fr. Tab. II. f 6—8 a—k.

(*Sphæriæ globiferæ* P. Tod. Calicii spec. Ach.)

Receptaculum nullum (in nonnullis tamen hypothallus heterogenus adest). Perithecium receptaculiforme varium, sæpius stipitiforme cy-

lindricum. *Gelatina* in globulos demum fati-
scentes erumpens.

257. *Sphaeronema cladonisca*, peri-
thecio cylindrico simplici l. ramoso, globulo
minuto punctiformi albo. Tab. II. f. 7. c—f
magn. auct.

Calicium cladoniscum. *Achar. syn. Lich.*
p. 61. *Lich. univ.* p. 241.

In lignis putrescentibus Helvetiæ, Lusa-
tiæ, *Sueciæ*.

Obs. Hypothallus tenuissimus albidus
sæpe deest. *Observ. in Lichen. Univ. Add.* p.
673 forte ad *Sphaeron. aliam* pertinet. Hanc
listit. Tab. II. f. 6. a. b.

258. *Sphaeronema ventricosa*, peri-
thecio simplici ventricoso compresso nigro, glo-
bulo minuto punctiformi cinereo. Tab. II.
f. 8. m. a.

Calicium ventricosum. *Achar. l. l. c.*

Ad corticem arborum in Lusatia.

Specimina cum iconè Cel. *Achario*, Fau-
tori summo, debeo.

Obs. Perithecium vix lin. attingens.
Hypothallus cinerascens in hac specie semper
fere adest. Colore & perithecii & hypothalli

globalisque, perith. cylindricis &c. sequens differt species.

259. Reliquæ hujus Generis species:

3. *Sph. acrospermum*.

Sphæria acrosp. *Tod. Fung. Meckl.* 2.

P. 47.

α. nigra. *Tod. l. c. t. 15. f. 119. Alb.*

& *Schwein.* p. 50.

β. tricolor. *Tod. l. c. f. 120.*

4. *Sph. cylindrica.* —

Sphæria cylindrica. *Tod. l. c. p. 42. t.*

15. t. 114.

Etiam in Suecia frequens. Primus legit

Cel. Prof. *Agardh.*

5. *Sph. acicularis.*

Sphæria dubia. *Tod. l. c. p. 45. Alb. &*

Schwein. l. c. p. 50.

6. *Sph. subulata.*

Sphæria subulata. *Perf. syn. p. 94. Tod.*

l. c. p. 44. t. 15. f. 117.

Vulgaris in Suecia.

7. *Sph. conica.*

Sphæria conica. *Tod. l. c. p. 43. t. 15.*

f. 116. *Alb. & Schwein.* p. 51.

In Suecia etiam viva.

8. *Sph. parabolica.*

Sphæria parab. *Tod.* l. c. p. 43. t. 15.
f. 115.

9 *Sph. hemisphærica.*

Sphæria hemisphærica. *Alb. & Schwein.* l.
c. p. 51. t. 8. f. 8.

In Suecia.

10. *Sph. versiformis.*

Sphæria versiformis. *Alb. & Schwein.* l.
c. p. 52. t. 9. f. 3.

Genus habitu & firmis characteribus optime distinctum. Affinitatem quandam cum *Nemasporis* habet. *Sphæria penetrans* *Tod.* (*S. patella* *P.*) *S. peziza* *T.* & *gelatinosa* *T.* (*pallida* *P.*), *Perfoonio* jam monente, hujus Generis non sunt.

ACTIDIUM. Fr.

Tab. III. f. 1. e, f.

Receptaculum nullum *Perithecium* sphæricum solidum, rimis pluribus radiatis. *Gelatina* nulla.

260. *Actidium Acharii.* Tab III.
f. 1. a—f. *Sphæria stellulata.* *Achar, see. spec.*

In ligno emortuo Ostrogothiæ. *Acharius.*

In ramis *Rhamni frangulae* in Hallandia.

Ipse.

Nudo oculo *Sphaeria* cuidam minutæ similis. Substantia interna *Hysteriorum*, quibus propius accedit, quam *Sphaeriis*.

LOPHIUM. Fr.

(*Sphaeria Macrostomæ*. P.)

Receptaculum nullum. *Perithecium* compressum l. subglobosum solidiusculum, *Ostiole* sublatissimo rimæformi cristato. *Gelatina* subindurata.

261. *Lophium fallax*, perithecio basi semiglobofo farcto, sursum cylindrico nigro, ostiolo lato aperto.

Sphaeria fallax. *Pers. syn.* p. 90. *Ic. & descr.* 2. t. 10. f. 11. 12. *Wahlenb. fl. Lapp.* p. 522.

In ramis & lignis exsiccatis.

Obs. Ceterum ad hoc Genus referendæ omnes *Sphaeria macrostomæ* *Pers.* — Inter *Sphaerias* & *Hysterias* medium est; his forma & characteribus externis, illis substantia accedit. In nonnullis tamen hæc etiam cum *Hysteriis* fere convenit. Omnes *Sphaeriae* nostræ ostiolo

æquali instructa sunt, & hinc hoc genus a me non temere esse distinctum, spero.

Hysterium mytilinum P. & *pedicellarum* Schum. etiam affinia sunt.

HYSTERIUM. TOD.

262. *Hysterium varium*, superficiale nigrum subelongatum varium, labiis subobsoletis discoque concoloribus.

In ligno exsiccato quercino durissimo, rarissimum.

Descr. Habitus a reliquorum recedit. Magnum; sed nudo oculo tamen *Sphæriæ* cui-dam immaturæ similis. *Forma* nunc rotunda, nunc oblonga, sæpius elongata. *Labia* sub lente distincta apparent, rugosa opaca demum confluentia & discum convexulum difformem formant.

263. *Hysterium melaleucum*, superficiale nigrum oblongum læve, labiis subconniventibus albis. Tab. II. f. I. a. b. c. d.

In foliis *Vaccinii Vitis idææ* emortuis copiose.

Descr. Pulchra & optime distincta species. *Perithecium* nigrum læve oblongum sub-

triquetrum linea transversali alba notatum, quæ sub lente *Labia* alba distincta lævissima.

264. *Hystერიum pulchellum*, superficiale nigrum ovale læve, labiis distinctis virescenti-flavis. Tab. II. f. 2 a—d.

In foliis *Vaccinii Vuis idææ* putridis, rarissime.

A priori omnino distinctum, forma ovali (nec triquetra) juniore concava, labiis distinctis aliter coloratis &c.

Huic proximum *H. versicolor* Wahlenb.; sed sec. spec. Lappon. omnino diversum.

265. *Hystერიum elongatum*, superficiale elongato-lineare opacum nigrum, labiis convexis læviusculis.

H. elongatum, oblongo-lineare acutiusculum tumidulum striatum. *Wahlenb. Fl. Lapp.* p. 523. *excl. syn. Pers.*

In ramis salicinis — *Rhamni Frangulæ* in Hollandia.

H. pulicari P. et *fraxineo* P. valde affine, ab utroque tamen diversum videtur. Ab hoc enim situ superficiali, forma magis angustata et ab illo forma, labiis læviusculis (etiam in spec. mea Lapponica) &c. differt.

266. *Hysterium graphicum*, superficiale radiato ramosum curvato - flexuosum difforme nigrum, radiis demum distantibus concoloribus.

In cortice *Pini sylvestris*, semel (sed copiose) legi.

Descr. Quoad habitum *Opegraphis* simillimum; sed crusta nulla et perithecia omnino *Hysterii*, quamvis ramosa radiata, qua nota a reliquis differt haec species. *Labia* juniora distincta subconniventia, dein distantia, disco dilatato opaco, demum tenuissima. Majusculum.

267. *Hysterium sticticum*, immersoerumpens minutum punctiforme oblongum acutum, labiis tenuissimis sub-convexis.

In ligno putrido in Helvetia.

Obs. *Stictibus*-affine; sed hujus sine dubio loci. Minutiae et forma a reliquis immeris differt. Sub lente determinanda; sub nudo enim oculo ut puncta cinereo-nigra apparet. *Labia* convexuscula laevia. *Perithecia* subnullum, hypothallo tenuissimo albicanti immersum.

268. *Hysterium ellipticum*, immer-
sum nigrum minutum ellipticum, labiis te-
nuissimis planiusculis.

In cortice crassiuscula laevigata *Pini sylve-*
stris. *Acharius*.

Priori affine; sed diversum videtur. Hy-
pothallus plane nullus. Sub nudo oculo caesio-
nigrum, sub lente nigrum perithecio tenui sub-
nullo, labiis planiusculis laevibus.

269. *Hysterium viride*, erumpens
elongatum subpulverulentum viride, perithecio
labiisque subnullis.

In ligno dejecto quercino putrido passim.

Dubia species, ob formam huc relata *Pe-*
rithecium proprium non vidi nec labia; ligno
enim prominente cinctum, *Discus* viridis, ju-
nior subpulverulentus; dein dilatatus. Ob sum-
mam cum *H. pezizoidi* Wahlenb. affinitatem
huc retuli. Species illa dubia *H. viride* Alb.
et Schw. hujus forte status imperfectus.

H. pezizoides W. et *Sorbi* Ejusd. in Sue-
cia australi etiam sunt lecta.

H. tuberculiforme Schum. non est vegeta-
bile. Varias adhuc species, *H. Berberidis*, *Co-*
tini, *Aucuparia* &, ex Helvetia favore Cel-
Acharii possideo; sed parum differunt duo.

ultima ab *H. petiolari* Alb. et Schw. aliisque
et *H. Berberidis* ad *Xylomata* nimis accedit.

LEPTOSTROMA.

Fr. Tab. I. f. 6 b.

Receptaculum nullum *Perithecium* tenuissimum effusum maculæforme disco subumbonatum. *Ostiolum* gelatinaque nulla.

270. *Leptostroma sphaeroides*, orbiculare dilatato-tenuissimum, disco convexulo lævi.

In caulibus Herbarum majorum ex. gr. *Cherophylli bulbosi* &c.

1—1½ lin. Ob discum latius umbonatum *Sparia* sub simile; sed defectu gelatinæ ostioli et fere perithecii differt.

Obs. Medium Genus inter *Hysterium* et *Xyloma*; ab utroque nimis differt perithecio subnullo. Affinia sunt *Hysteria* sine perithecio vero et hujus forte Generis.

271. *Leptostroma hysterioides*, oblongum nigrum varium, disco substriato carnuloso.

In caulibus variarum plantarum ex. gr. *Euphorbia palustris*, *Pæonia* &c.

Forma ovali oblonga distinctum.

272. *Leptostroma xylomoides*, subrotundum varium nigrum, disco ruguloso. Tab. I. f. 6 a. b.

In caulibus Herbarum, præcipue *Scirpi lacustris*.

Xyloma macula olim dixi.

273. *Leptostroma filicinum*, elongatum nigrum difforme, disco læviusculo tenui. In stipitibus *Osmundæ regalis* &.

XYLOMA. P.

274. *Xyloma Leptostromæ*, epixylum tenuissimum nigrum subrotundum, dein irregulariter ruptum.

In Caulibus *Vaccinii Vitis idææ* emortuis. Bene distincta species; sed habitum &c. *Leptostromatis* propius accedit. Cortex vero exterior reliquorum Xylomatum epixylorum instar rumpitur, ut de genere nullum sit dubium — 1 lin. latum.

275. *Xyloma Montiniæ*, epiphyllum orbiculare clausum planiusculum nigrum, intus fuscescens.

In Capite Bonæ spei in foliis *Montiniæ acris*. Dedit Cel. *Acharius* sub nomine *Sphæriæ cassæ*.

Inter majores species, 2 l. latum orbiculare, nudo oculo laeve, armato ruguloso. Colore interno optime distinctum.

276. — *Xyloma Lonicerae (rubiforme)*, epiphyllum compositum orbiculare sparsum convexum nigrum tuberculosum, tuberculis prominentibus. Tab IV. f. 7. a—c.

In Helvetia ad folia *Lonicerae*. Sub hoc nomine liberalissime communicavit Cel. *Acharius*.

Descr. Pulcherrima species. *Tubercula* astoma nitenti-nigra laevia in acervos orbiculares æquales convexos 1—1½ lin. lato sparsos conglomerata. An cum sequenti ad *Sphaerias* referendum?

X. rubiforme potius dicendum.

277. *Xyloma betulinum*, epiphyllum compositum angulato difforme confluens planiusculum nigrum tuberculosum, tuberculis punctiformibus.

In foliis *Berulae albae* semivivis autumnno, frequens in Suecia.

Descr. Priori quidem affine; sed certissime distinctum. *Forma* juniorum, punctiformis, in adultis confluentibus maxime inæqualis. *Perithecia* 1—2 lin. lati; dein vero

confluentia, planiuscula nigra superne punctato-tuberculosa.

278. *Xyloma Solidaginis*, epiphyllum maculaeformae orbiculare cinereo-nigrum, peritheciis simplicibus subglobosis aggregato-confluentibus.

In pagina inferiori foliorum vivorum *Solidaginis Virgaureae* in monte inter *Kungibacka* et *Tölo* in Hallandia boreali tantum l-gi. Aestate.

Descr. Cum *Xylomate punctato* P., cui tamen sec. specimina mea e Germania vix affine, comparandum. In pagina foliorum superiori observantur maculae steriles irregulares flavescentes et in inferiori orbiculares 3—4 lin. latae l. confluentes planae tuberculosae, *Uredini* cuidam familia, cinereo nigra margine flavescenti. Sub microscopia observantur *Perithecia* numerosissima globosa densissime aggregato-confluentia opaca cinerascenti-nigra.

Etiam ab *X. concentrico* distinctissimum.

279. *Xyloma Linnaeae*, petiolis immersum nervisequium lanceolatum laeve nigrum, intus album.

In petiolis *Linnaeae borealis*.

Xylomati nervali, Alb. et Schwein proximum; sed diversum esse ex ipsa diagnosi patet. 2—3 lin.

In foliis *Linnae* crescit *Sph. setacea* P.

280. *Xyloma Pteridis*, epiphyllum superficiale subnervisequium elongato-lineare cinereo-nigrum subcorrugatum, intus subconcolor.

Sphaeria Pteridis. *Rebent. Fl. neon. t. I.*
f. 3. p. 344.

In foliis subemortuis *Pteridis aquilinae* frequens.

Hujus, ni fallor, generis et *X. nervali* proxima species. Semel ruptum inveni et pulvere nigricante fartum.

281. *Xyloma Trientalis*, epiphyllum subrotundum l oblongum tenue rugulosum nigrum intus concolor.

In foliis emortuis *Trientalis europae*.

Ab affinibus (*X. Andromedae*) tenuitate et colore interno differt.

Obs. Xylomata haec majora valde inter se differunt et ad distincta forte genera referenda. Saltim diversas efficiunt sectiones, ex, gr.

A) *Epixyla*. Alb. et Schw.

B) *Tuberculata*: Fr. *Xyloma betulinum*,
Lonicerae.

C) *Macularia*: Fr. *A. punctatum* P. *Solidaginis* Fr.

D) *Nervisequia*: X. *nervale* Alb. *Linnaeae*, *Pteridis*.

E) *Contigua*: X. *Andromedae*, *Trientalis*, *Montinae*, *acerinum* &.

F) *Punctata*: X. *rubrum* P. *aurantiacum* Schl. ad *Sphaerias retuli*.

282. *Xyloma Taxi*, minus applanatum nigricanti-cinereum opacum laxè et irregulariter ruptum, nucleo tenui nigrescente.

In foliis *Taxi* in Horto Lundensi, raro obvium.

Descr. Subgregarium. A. [*Sph. Taxi*, cum qua non confundendum, colore opaco cinereo, forma applanata 3plo latiori, epidermide irregulariter rupta, laciniis albicantibus &c. differt.

283. *Xyloma Ilicis*, minus convexum nigrum transversim ruptum, nucleo tenui nigrescente.

X. *Ilicis*. Ach. *sec. spec.*

In foliis *Ilicis Aquifolii* in Europa australi.

Priori affine; sed multo minus, colore, forma et rupendi modo differt. Laciniæ albicantes,

Tribum naturalem haec duo species et nonnullæ aliæ efficiunt.

284. *Xyloma erumpens*, minutum erumpens nigrum, junius sphaericum, dein umbilicato-depressum, demum cupulare.

In foliis emortuis *Vaccinii Vitis idæa*, rarius.

Nullam aliam speciem laciniis epidermidis foliorum cinctam erumpere vidi. Clausum de cetero remanet, quod sub nomine X. *Vaccinii uliginosi* Wahleub. dedit Cel. Swartz.

X. *erumpenti* nostro non affine; sed ad sequens potius accedit.

285. *Xyloma geographicum*, minutum concaviusculum nigrum, macula tenuissimæ flavescenti insidens.

X. *geographicum*. Pers. sec. spec. ab. Achar. liberalissime misa.

Lichen castanearius. La Mark.

In foliis *Fagi Castaneæ* in Gallia.

Nudo oculo punctiforme; lente concaviusculum subpezizoideum, laeve opacum parum difforme.

Cum *Sph. Castanea* Schum., etiam apud nos vulgatissima, confundi nequit. Etiam sc. ipsec. ex Helvetia omnino differt *Xyloma Castanea* Schleich.

Xyl. Convallaria, Schleich. ad hoc genus vix pertinet.

Plurimas alias hujus Generis species legi, ex quibus *Xyloma Sorbi*. Mihi (compositum) in foliis *Sorbi aucupariae* semivivis autumnis vulgatissimum.

SCLEROTIUM. T O D.

Genus priori maxime affine. Sic ex gr. huc referri possent *X. nervale* Alb., *Linnaea* Fr. et ad *Xylomata*.

286. *Sclerotium dipsaceum*, nervisequium nigrum oblongum applanatum, ipso subconcolor. In Calycibus exsiccatis *Dipsaci*, hieme et vere.

Deser. *Scl. duro*, ut videtur, proximum - Calycibus totum aëctissime adnatum et cum horum substantia quasi connatum. *Forma* varia, in calyce communi ovato-acuminata, in reliquis oblonga lanceolata l. linearis, semper nervisequia. *Substantia* carnosae tenuis extus

nigra, intus cinereo-nigricans. *Superficies* lævis glabra; sed e ciliis calycis pseudohispida. β . *Stramonii*, rotundo-oblongum conglomeratum.

In capsulis *Daturæ Stramonii*, vere.

287. *Sclerotium rectum*, magnum subtectum difforme applanatum læve nigrum, intus albidum.

In caulibus exsiccatis Herbarum majorum, primo vere.

Sub epidermide crescere incipit, hinc subtectum. $\frac{1}{4}$ — $\frac{1}{2}$ unc. longum.

288. *Sclerotium vulgatum*, liberum globosum flavum læve glabrum, intus albidum.

A. *vernale*, opacum, interdum difforme. In stipitibus dejectis putrescentibus, vere.

μ . *Filicinum*, læte flavum globosum liberum.

In frondibus dejectis *Pteridis Aquiline* æstate.

γ . *Fimetarium*, flavum globosum subimmersum.

In fimo locis sylvaticis, rarius. *Autumno*.

289. *Sclerotium rugulosum*, oblongum nigrum semiaplanatum glabrum subrugulosum.

In *Agarico* putrido exsiccato.

Uno latere planum, altero convexum 2. lin. læve, sub lente rugulosum.

290. *Sclerotium cornutum*, gregarium confertum rufum corniforme læve glabrum, intus albidum.

In *Agaricis* putridis.

Descr. 3—5 lin. longum, cornu caprino exacte sæpe simile, læve glabrum rufum intus albidum, basi rotundatum obtusum, apicem versus subcurvatum attenuatum acutum.

In hac specie, et in nonnullis aliis *Sclerotiiis*, parasitat *Ag. tuberosus* P. — Ab omnibus, tamen Auctoribus pro basi stipitis hujus fungi habetur; at perperam. Conf. *A. tuberosus* p. 39.

291. *Sclerotium foliaceum*, foliaceum sublobatum nigrum.

In stipite *Agarici parasitici* P. parasiticum, autumn.

Nulli alii affinis et distincti forte Genèris fungus. Folium sublobatum exacte refert, erectum tenue nigrum opacum glabrum 3—4 lin. planum læve l. rugosum.

292. *Sclerotium tomentosum*, subrotundum album villosum.

Scl. villosum, sphaericum villosum aggregatum. *Tod. Fung. Meckl.* l. p. 6, t. I, f. 10? (sub *Peziza villosa* P.)

In tuberibus putrescentibus *Solani tuberosi* *asfare*.

2—3 lin. Etiam distincti, forte generis species.

ERYSIPHE. DE CAND.

293. *Erysiphe varium*, receptaculo albo, capitulis flavescentibus fuscisque.

Sclerotium *Erysiphe*. *Pers. syn.* p. 124. *Obs. Myc.* p. 13. et *Auct. recent.*

Scl. *Heraclei*.

Mucor *Erysiphe*. *Linn. Suec.* p. 462. et *Auct. veter.*

Species maxime vulgaris et polymorpha, in foliis plantarum copiosissimum vitium.

β. *suffultum*, disco impresso villoso.

Sclerotium *Erysiphe*. β. *Corylae*. *Pers. syn.* p. 124.

Scl. *suffultum*, epiphyllum, granulis globosis fusco-nigricantibus, in basi pilis rigidiusculis suffultis, disco impresso villoso. *Reb. Fl. Neom.* p. 300 t. III. f. 14 a. b.

Variant *Capitula* suffulta, pedicellata et sessilia; flava et fusca; globosa, plana et depressa & sæpe in eodem specimine.

294. *Erysiphe radiosum*, receptaculo fusco fibroso-radiato, capitulis nigris.

Sclerotium radiosum. *Achar.*

In foliis *Rosæ*. *Acharius.*

In foliis observantur maculæ tenues fuscescentes 3—5 lin., sub lente fibrosæ stellato-radiatæ, capitulis nigris l. nullis.

295. *Erysiphe album*, receptaculo tenuissimo subnullo, capitulis nigris.

Er. album. *Achar.*

In foliis *Iuglandis* in Helvetia.

Nudo oculo ut puncta alba apparet.

TUBERCULARIA. P.

296. *Tubercularia Acaciæ*, minuta applanato-fessilis convexa dura rubra, intus concolor, humectata flavescens.

In ramis emortuis *Robinia Pseudoacaciæ*, vere.

Species bene distincta, licet reliquæ hujus generis species nimis confluunt. Substantia compacta firma nec humectata gelatinosa, nec ficca farinosa. *T. salignæ* (vid *Alb. et Sw.* p. 69), in *Scania* semel lectæ, affinis quidem est, sed species vere diversa.

297. *Tubercularia farmentorum*, minuta sessilis convexa oblonga extus farinosa rubra, intus gelatinosa obscurior.

In ramis exsiccatis *Vitis viniferae*, *Hedera Helicis*, *Panmaceris 5 folia*, *Clematis* &c.

Sequenti affinis; sed magnitudine, colore, forma et substantia differt. Intermedia est:

298. *Tuberc. Menispermii*, minuta sessilis subglobosa dura obscure rubra, intus concolor.

In ramis *Menispermii canadensis*. Media inter *T. Acaciae*, *farmentorum* et *herbarum*.

299. *Tubercularia herbarum*, majuscula sessilis subglobosa pallide rubra, intus concolor, humida gelatinosa, sicca farinosa.

In caulibus exsiccatis herbarum ex. gr. *Cucubali tartarici* &c. Statu humido *Agyrio Urricae* Fr. (*Tremella* P) fimillima; sed colore dilutiori, forma subglobosa et exsiccatione differt. — *T. Artemisiae* Schum. stipitata, minor &c.

300. *Tubercularia Liceoides*, gregaria sessilis globosa, cortice exteriori fusco-spadiceo rufo deciduo, pulvere argillaceo-roseo nucleum tegente.

In ramis deciduis vere.

Alii vix affinis; sed veram *Tubercularia* speciem esse nullus dubito. Gregaria saepe

confluens laevis granuliformis, granulis magnitudine seminum *Viciae*, rarius *Pisi*; cortice exteriori rufo-fusco l- fusco-spadiceo demum deciduo. Dein farina subcompacta copiosa argillacea rosea includit fungum, quasi nucleum intus duriusculum pallidum, zona obscuriori fusca circumdatum.

STILBUM. P.

301. *Stilbum Lanugo*, capitulo sub-turbinato, stipiteque nigro.

In caulibus et foliis emortuis putridis *Paeoniarum* vere.

Nudooculo villum tenuissimum sistit, submicrosc. comp. stipes distinctus saepe flexuosus, deorsum subincrassatus. Capitulum nigrum nunc turbinatum nunc oblongum saepe deciduum.

MUCOR. P.

302. *Mucor albo-virens*, caespitosus ex albido-virens, peridiis globosis adultis albidis.

In *Hydno aurantiaco* exsiccato vidi. Iunior viridis est stipite brevi: dein stipites elon-

gati et peridia albicantes. Sporidia minuta numerosissima.

303. *Mucor Dematium*, cæspitosus, peridiis albis, stiptibus tuberculo radiato-infertis.

In foliis *Hedera Helicis*, rarior.

Minuta species, tuberculo suo facile distincta. Fugax est; præcedens vero subpersistens.

MONILIA. LINK.

304. *Monilia hyalina*, filis aggregatis simplicibus erectis sursum crassioribus hyalinis, articulis subglobofo-ovatis. *Achar.* — Ic. Tab. III. f. 4. a—d.

In culmis gramineis.

Media videtur inter *Torulas* et *Monilias* *Link*; sed ob fila erecta subrigida et articulos non exacte globosos hujus loci.

HIMANTIA. P.

305. *Himantia ochracea*, densissime intertexta ochracea, fibris tenuissimis.

In pinetis ad strobilos *Pini Abietis* putridos &c.

Xylostromatis affinis; nam densissime est intertexta. Pulchra species et ab *H. sulphurea* vere distincta.

306. *Himantia Rosa*, radiato-ramosa rosea, fibris crassiusculis, extimis byssaceis.

In cortice *Iuniperi*, *Pini* &c. dejecto.

Colore a reliquis differt; margine subalbicat. Nullius fungi initium.

307. *Himantia grisea*, densissime contexta villoso-pulverulenta grisea, fibris tenuissimis.

In *Hypno prolifero* locis silvaticis, rarior.

Nudo oculo villoso-pulverulenta et fibri sub microsp. compos. detegendi.

308. *Himantia violascens*, densissime contexta violaceo-alba, fibris arachnoideis.

Ad terram nudam in silvis.

Fibri etiam hujus sub microsp. compos. determinandi.

Obs. Omnes hujus generis species Poriarum, Thelephorarum aliorumque fungorum primordia esse statuit Cel. *Link*; sed haec sententia in Observationes meas non quadrat *).

*) Rhizomorphas tantummodo Merulii esse primordia pessime statuit Cl. *Palliot*. Conf. hujus Generis monograph. in Vet.

Hyphasma funicola, nigra et nonnullæ aliæ species Rebent. in Suecia crescunt et ad hoc Genus referendæ.

A c. Hand. 1814. p. 201. f. IX. 1—15. Auct. Achario. Distinctissimum esse hoc Genus et ad Lichenes referendum l. c. luculentissime demonstravit. Species nonnullas a me detectas et l. c. descriptas, novo uno alioque synonyme addito, enumerare placet.

1. *Rhizomorpha Alectorioides*, thallo plano compresso gracili nitido elevato-substriato atro, intus griseo, ramosissimo, ramis capillaceis parentibus ramulosis confertis complicatis. Achar l. c. p. 214. t. 9. f. 3. a.

Rh. alectorioides. Fries.

In trunco putrido coryleo, ramis Vaccinii Vitis idææ, Myrtilli &c.

β. subsimplex, thallo subsimplici tortuoso striato. Ach. l. c. f. 3. b.

Rh. alect. subsimplex. Fr.

In ramulis et foliis dejectis.

Distinquenda ab Rh. fetiformi β. scandente Fr. — Achar. l. c. p. 213 in Pinis destruccis raro observata.

2 *Rh. Racodioides*, thallo teretiusculo molli tenerrimo atro opaco, intus concolori ramosissimo; ramis creberrimis flexuosis attenuatis densissime intricatis, ramulis

Obs. *Ozonium auricomum* Link in Suecia crescit cum variis aliis novis hujus Generis speciebus; cum autem Cel. *Link* Ipse se plu-

capillaceis multifido-cirrhosis. Achar. l. c. p. 215. t. 9. f. 15. a. b. Rh. Racodioides. Fr.

In Rhizomorpha angulosa emortua ad Femsiö. Conf. Perf. disp. meth. Fung. p. 42.

3. Rh. cirrhosa, thallo tereti gracili glabro dilute castaneo intus albedo, superne ramoso, fibrillis lateralibus terminalibusque capillaceis simplicibus cirrhosis. Ach. l. c. p. 217 f. 11. Rh. cirrhosa. Fr.

In trunco fagineo putridissimo ad Femsiö in Smolandia.

Pulcherrima species.

4. Rh. corrugata. Achar. syn. Lich. p. 391. Ver. Ac. Handl. p. 221. t. 9. f. 13.

In truncis putridis in palude ad Femsiö Iecra.

5. Rh. angulosa Ach. syn. Lich. p. 228. Ver. Ac. Handl. p. 226. f. 12.

In Scania, Smolandia.

6. Rh. Friesii, thallo teretiusculo renui strigoso fragili nigro, intus concolori, dichotome ramoso, ramis ramulisque erectis strictis acuminatis. Achar. Ver. Ac. Handl. p. 227. t. 9. f. 4. a. b.

res novas species vidisse dixerit, harum determinationem expectare satius duxi.

Rh. stricta. Fr.

Sub cortice Pini sylvestris ad Fem-
flō, rarissime.

7. *Rh. Achariana*, thallo tereti fili-
formi laeviusculo fragili nigro-fusco opaco,
intus rufescente, simpliciusculo pinnato-ra-
moso; ramis lateralibus approximatis hori-
zontalibus distichis rectiusculis attenuatis api-
ce subdivisis rufis. Achar. l. c. p. 229. t.
9. f. 8 a. b.

Rh. Achariana. Fr.

In ligno dejecto abietino, frequens in Smo-
landia. Superficie, substantia &c ab Him. um-
brina maxime differt.

8. *Rh. muscicola*, thallo compres-
siusculo tenui glabro elevato-striato obscure
fusco, intus griseo, ramoso, ramis sparsis
capillaceis planis adpressis raris elongatis cur-
vatis subsimplicibus. Ach. l. c. p. 229 t.
9. f. 6.

Rh. muscicola. Fr.

β epiphylla. Fr.

Inter muscos locis humidis umbrosis. *β*
in foliis dejectis.

9. *Rh. corticata.* Ach. syn. p. 391.
Ver. Ac. H. p. 232. t. 9. f. 5.

RACODIUM. P. LINK.

309. *Racodium vulgare*, expansum molle laxè contextum nigrum, filis æqualibus.

Rhiz. filiformis. Fr.

Agaricum luteum, radicibus arborum fibrosis simile inter humum et ligna l. lapides innascens. Mich. Gen. p. 125.

In ferragine ad Femio.

Sub nomine: *Elvella monstrosa*, lutea et alba corpore fibroso, inæquali, ramosissimo et expanso, hanc cum sequenti conjunxit *Gleditsch* in libro paradoxo: Meth. fung. p. 52. XII. a.

10. Rh. *Xylostroma*. Fr. Achar. l. c. p. 233. t. 9. f. 7. Rh. cancellata. Ach. syn. Lich. p. 391.

Agaricum candidum, radicibus arborum fibrosis simile &c. Mich. gen. p. 125. n. 22.

In truncis cavis et ad terram.

11. Rh. fibrillosa, thallo teretiusculo capillari molli lævi candido ramosissimo fibrilloso, axillis compressis, ramis strictis divaricatis subanastomosantibus sursum fibrillosis. Ach. l. c. p. 228. t. 9. f. 9.

Rh? fibrillosa. Fr.

Elvella nivea corpore reticulari, fibroso, ramis compressis, in apice cristato-lacinatis.

Rac. nigrum. Schum. Saell. II. p. 447.

Dematium pinastrum. Schleich. sec. spec.

Helw. (Non differt).

In ligno et ramis *Betulae*, *Fagi*, *Pini*, *Abietis* &c. locis umbrosis, muscis et foliis tertis.

Purimæ *Racodii* species nigrae sunt. A. R. cellari P. difficile distinguitur, nisi colore magis nigro, contextu laxo, loco &c.

310 *Racodium resinae*, expansum molliusculum dense contextum nigrum, filis inaequalibus.

In resina *Pini Abietis* in silvis Sueciae passim.

Habitu et loco natali distinctum. Fila divaricato-ramosa; aia rigidula apice capitulifera, sub microsc. *Coremio* Link. similia. *Demat. villosum* Schleich huic simile; sed sub microsc. fila maxime differunt.

Gled. l. c. b. *Himantia albida*. Pers. disp. Feth. Fung. p. 42.

Obs. Ultimos tres, ut etiam R h. capilaris Roth ad fungos potius pertinent, et 9-10 distinctum forte efficiunt genus.

TAPHRIA. FR.

Erinei spec. P.

Thallus e floccis cæspitosis globosis granulæformibus non septatis foliis cupulaforme impressus.

311. *Taphria populina*, aurea.

Erineum aureum. Ach. — *Perf. syn.* p. 700 *Alb. et Schw.* p. 371. *Schum. Saell.* II. p. 446.

In foliis *Populi dilatata*, *nigræ*, *tremulæ* &c. æstate.

Genus maxime distinctum nec sequentibus affine. An flocci sic dicti in hoc sporidia? Ulterius examinandum, specimina enim siccatandum vidi.

PHYLLERIUM. FR.

Erineum. Link.

Thallus e floccis cæspitosis simplicibus varie tortis subcompressis non septatis, apicibus subattenuatis.

Obs. Ad *Rubiginem* Link, nomen alii subgeneri jam sacratum, nomen *Erinei* reservo; in hoc enim quadrant descr. *Perfoonii* et plurimæ species.

312. *Phyllerium sorbeum*, floccis dense intricatis compressis latiusculis albis, exficcalis flavescentibus.

In foliis *Sorbi aucupatia*, frequens. Aestate.

In utraque foliorum pagina viget. Nudo oculo lanam refert. Flocci sub microscopio composito simplices intricati varie torti compressi, *Ulae* subsimiles.

313. *Phyllerium acerinum*, floccis densissime intricatis tenuissimis filiformibus albedo cinereis.

In foliis *Aceris platanoidis*, vere.

In utraque pagina foliorum crescit. Nudo oculo præcedenti simillimum, hoc vero magis villosum, illud grumosum apparet. Sub microsc. composito flocci quam maxime differunt, ut e diagnosi patet.

314. *Phyllerium juglandinum*, planum floccis dense intricatis brevibus filiformibus roseo - cinereis.

Erineum juglandis. *Schleich.*

In foliis *Juglandis* in Helvetia, unde mea specimina habeo.

Paginam folii inferiorem occupat, sequenti affine; sed forma, focus &c. differt.

315. *Phyllerium tiliaceum*, convexum pulvinatum, floccis densissime intricatis variis pallide-roseis.

An *Erineum tiliaceum*. Pers. syn. p. 700.

Obs. Myc. I. p. 25. et Auct.?

In foliis *Tiliae Europae* vulgare. Aestate.

De synonymis dubius cum cinereo-pallidum omnes describunt Auctores, nostrum vero pallide roseum. Utramque foliorum paginam amat.

316. *Phyllerium viteum*, effusum planiusculum, floccis intricatis tortis compressis, dein rufis.

Erineum Vitis Schrad. — Funk. Crypt. Gew. Fichteg. n. 285.

In pagina inferiori foliorum *Vitis viniferae* in Helvetia, Germania.

Habitus Erinei; sed sub micr. comp. patet hujus esse Generis. Flocci torti pelucidi.

317. *Phyllerium pyrinum*, effusum crassiusculum, floccis laxè intricatis variis aureis.

Erineum pyrineum. Auct.?

In foliis *Pyri Mali*, *Pruni spinosa* &c.

Colore ab *E. pyrino* Auct. differre videtur; sed loco et reliquis notis convenit.

Specimina in *Pruno spinosa* parum tenuiora. Æstate viget.

Color in *Phylleriis* et *Erineis* maxime variabilis.

318. *Phyllum Gei*, floccis subliberis tenuissimis rectiusculis filiformibus albidis.

In foliis *Gei rivalis*, æstate. (In monte Kullaberg tantum legi).

Folio immersum. Filis rectiusculis parum intricatis et habitu a reliquis maxime recedit.

CRONARTIUM. Fr.

Thallus e floccis aggregatis simplicibus cylindricis rectis non septatis, apicibus æqualibus.

319. *Cronartium asclepiadeum*, rufescens.

Erineum asclepiadeum. *Funk. Crypt. Fichtelgeb.*

In foliis *Asclepiades*. Ex horto Upsalensi dedit Cel. *Swartz*.

Obs. Superficiale, varie effusum. Flocci aggregati non cæspit. si nudo oculo facile distinguuntur.

ERINEUM. P.

Rubigo. Link.

320. *Erineum purpureum*, effusum superficiale supra-foliaceum rufo fuscum dein sanguineo purpureum, cupulis stipitatis scutellatis, margine erecto, stipite crassiusculo brevioribus.

E. betulinum. *Alb. et Schw.* l. c. p. 370.

E. roseum. *Sculze Starg.*

In pagina superiori foliorum *Betulae albae*, vere et aestate.

Omnes Erinei species sub microscopio composito determinandæ. Color enim maxime ludit. In hac vero specie non nisi in prima infantia pallide roseum vidi; sed semper fere purpureum, quale etiam descripserunt *Albertini* et *Schweinitz*.

221. *Erineum betulinum*, conglomeratum subimmersum infra-foliaceum albidum dein rufo fuscum, cupulis stipitatis pyxidatis stipite deorsum angustato dupplo brevioribus.

E. betulinum. *Schum. Sall.* II. p. 445.

In pagina inferiori foliorum *Betulae albae* delaps. l. semiviv., autumnus.

Maxime a priori differt; tempore et loco natali, colore, forma &c.; sed præcipue cu-

pularum forma; in hac specie pyxidata l. turbinata, in illa scutellata l. rotundato-excavata.

322. *Erineum padineum*, effusum infrafoliaceum superficiale albidum mox spadiceorum, cupulis stipitatis urceolatis, marginem versus constrictis.

In foliis *Pruni Padi* et pagina inferiori autumnno.

Ad formam juniorem pertinet dubia species *E. Padi* apud *Alb. et Schw. l. c.* — Cupularum forma admodum differt a reliquis hac species.

323. *Erineum acerinum*, subeffusum infrafoliaceum submersum, primo albidum, dein roseo-rubrum, demum rufo-fuscum, cupulis stipitatis late cupulatis stipitem subæquantibus.

E. acerinum. *Pers. syn. p. 700. Disp. meth. p. 43. Schum. Saell. l. c. Alb. et Schw. p. 370. Link. in Berl. Mag. 1809. t. 1. f. 36. (optime).*

Mucor ferrugineus. *Bull. t. 5^σ 4. f. 12.*

In foliis acerinis vivis meum fungum legi, in delapsis *Pers. — Schum. — Albert. — Aestate autumnno et vere.*

Cupulæ dilatatæ.

324. *Erineum populinum*, conglomeratum imbricatum flavescens, mox rufescens, demum spadiceum, cupulis minutis scutellatis breviter stipitatis, stipite crasso.

E. populinum. Pers. Syn. p. 700. Obs. Myc. I. p. 100. Schum. Saell. II. p. 446. Alb. et Schwein. p. 371.

In inferiori pagina *Populi tremulae* foliorum. Aestate viget; sed perennat.

Folia et viva et sicca amat. Cupula folii cum cupulis floccorum, quas in diagnosi intellexi, non confundenda.

325. *Erineum alneum*, conglomeratum superficiale pulvinatum fulvum, cupulis stipitatis subconvexis tuberculosis.

Er. alneum. Pers. Syn. p. 701. Schum. l. c. Albert et Schwein. c. l.

In foliis dejectis *Alni glutinosae* autumnis.

Paginam inferiorem eligit. Stipes crassiusculus; cupulae convexae tuberculosae, forma *Helotii tuberculati*. Fl. Dan.

326. *Erineum negundineum*, effusum superficiale infrafoliaceum tenue luteolum, cupulis turbinatis cum stipite confluentibus.

E. tiliaceum. Schleich. sec. spec. Helv.

In foliis *Aceris Negundinis* in Helvetia.

A *Phyll. tiliaceo* quam maxime differt;
nec alii affine. Adinodum tenue et paginam
inferiorem incolit.

327. *Erineum platanoides*, latissime
effusum superficiale infrafoliaceum tenue cine-
reum, cupulis irregularibus variis.

In foliis *Aceris platanoides* in Suecia, vere
et astate.

Erythra species primo obtutu esset, quoad
crustam, nisi flocci sub micr. comp. ejusdem
essent naturæ ac reliquarum specierum. For-
mam vere distinctam et regularem detegere non
potui. *E. acerino* minime affine.

388. *Erineum extentum*, conglomerato-
lineare seriatum convexum infrafoliaceum su-
perficiale crassiusculum, cupulis minutis vix
distinctis, rufum.

E. extentum. Achar.

E. Guinea in foliis arboris novi generis.
Afzelins.

Nec formam cupularum hujus speciei cer-
tam determinare potui.

Obs. *E. fagineum*. P. nondum vidi.

AREGMA. Fr.

Pucciniæ Heterocl. Link.

Uredo. 3. c. von Straur.

Sporidia cylindrica clausa septulis pluribus distincta stipitata in plantis vivis parasitica.

329. *Aregma obtusata*, sporidiis obtusis, septulis tribus, pedicello attenuato.

Uredo obtusa. Straur in Acc. Vetteraw.

2. p. 107. t. II. f. 49.

Puccinia potentillæ. Pers. syn. p. 229. et alior.

In foliis *Potentillæ argenteæ* frequens vere et Autumno.

Etiam in *Tormentilla erecta. Straur.*

Genus distinguendum esse jam monuit. Cel. Link; non enim "forma solum sporidii" semper clausi et septulis differt; sed "toto etiam habitu et crescendi modo" a veris *Pucciniis*, quod in Generibus distinguendis maximi semper est momenti.

Aregmam, *Bullariam*, *Pucciniam*, *Uredinem* &c. egregie illustravit Lb. Bar. von *Straur*; sed haec Genera non recte conjunxit, ut e scriptis summorum Mycologorum *Persoonii* et *Linkii* patet, Eiusque de varietate sporidiorum conclusio ad hoc saltem Genus non pertinet.

330. *Aregma mucronata*, sporidiis acutis, septulis subtribus, pedicello apice fusiformi terminato.

Puccinia mucronata. Pers. syn. p. 233 a. rosæ (exclus. var. β .) Alb et Schwein. p. 134 a.

P. Rosæ. Schum. Sall. p. 235.

Uredo mucronata. *Straur* l. c. p. 103.

f. 50.

In plurimis *Rosæ* speciebus, præcipue in *Rosa coriifolia* Fr. sed etiam in speciebus foliis glaberrimis, ex. gr. in *R. canina*, *glauca* Fr. &c.

Nunc folio nunc *Uredine Rosæ* insidet.

331. *Aregma acuminata*, sporidiis acutis, septulis 3 quibus, pedicello apice elongato-incrassato acuto.

In foliis *Rubi saxatilis*, autumnis.

Media inter antecedentem et sequentem; illi vero proxima. Sporidia acervata cæspites majores villosos formant. Apex incrassatus fere fusiformis, sed multo tenuior et brevior ac in præced.

332. *Aregma bulbosa*, sporidiis acutiusculis, septulis subquatuor, pedicello apice incrassato obtuso.

Uredo bulbosa. *Straur* l. c. p. 108 f. 51.

Puccinia Rubi. *Schum.* l. c.

P. mucronata. β . *Pers.* l. c. *Albert. et Schw.* l. c.

In *Rubo* idæo alii que.

Obs. Aliam speciem magis distinctam semel vidi in foliis *Comari palustris*. Omnes *Aregmæ* species in *Plantis Senticosis*, ut *Rosa*, *Rubo*, *Potentilla*, *Tormentilla* &c. crescere videntur et in foliorum emarcidorum l. semivivorum pagina inferiori.

EXPLICATIO ICONUM

Tab. I.

Fig. 1. *Merisma clavellare*. magn. nat.

Fig. 2. *Sphaeria multiformis*, a. coarctata.

a. Specimina plura. magn. nat.

b. Specimen tibeusium. magn. nat.

c. — dissectum in us nigricans, cum sperululis periphericis. magn. auct.

Fig. 3. *Sphaeria multiformis*, β . dilata. magn. nat.

Fig. 4. *Sphaeria rhodogramma*.

a. magn. nat.

b. specimen dissectum. magn. auct.

Fig. 5. *Sphaeria repanda*.

a. spec. integrum. magn. nat.

b. spec. dissectum. magn. nat.

c. Ejusdem pars. magn. auct.

Fig. 6. *Leptostroma xylomoides*.

a. spec. elongata. magn. nat.

b. — orbiculata disco rugoso. magn. auct.

Obs. In his sub microscopio composit. nihil aliud detegi, adeoque has icones sufficientes credidi.

Tab. II.

Fig. 1. *Hysterium melaleucum*.

- a. Folium cum fungillis. magn. nat.
- b. pars folii cum fung. magn. auct.
- c. Perithecium solitarium, magis auct.
- d. ——— e latere visum. Ejusd. magn.

Fig. 2. *Hysterium pulchellum*.

- a. b. c. d. Eisdem modis ac antecedens depictum.

Fig. 3. *Sphaeria malina*.

- a. macula cum sphaerulis. magn. nat.
- b. Sphaerulae dissectae. sub. micr. comp. pos.

Fig. 4. *Sphaeria macularis*.

- a. Folium populneum cum maculis cinereis absque sphaerulis, nudo oculo inconspicuis. magn. nat.
- b. Pars maculae cum sphaerulis sub micr. comp. pos.
- c. Sphaerulae e latere visae. Sub micr. comp.

Fig. 5. *Sphaeria argillacea*. P.

- a. Cortex cum sphaeriis. magn. nat.
- b. Sphaeria solitaria. magn. auct.

Fig. 6. *Sphaeronema cladonifera*.

- a. b. Hujus initium sec. Acharium, parum a Sph. cylindrica aliisque diversum. Sub micr. comp.
- c. d. e f. Specimina variae formae. sub micr. comp.

Fig. 7. *Sphaeronema ventricosa*.

- g. h. i. k. Spec. varia sub micr. comp. pos.

Tab. III.

Fig. I. *Aetidium Acharii*.

- a. ramulus decorticatus cum fungillis magn. nat.
- b. Ejusdem pars, magn. auct.
- c. d. Perithecia e latere visa. sub. micr. comp.
- e. f. Eadem superne visa, sub micr. comp.

Fig. 2. *Sphaeria* extensa.

- a. ramulus cortice rectus sphaerulis erumpentibus, magn. nat.
- b. — — orbatus, sphaerulis stromati extenso insidentibus magn. nat.
- c. Cæspes Sphaeriarum cum parte stromatis.
- d. Ostiolum junius, apice puncto depresso.
- e. f. Ostiola adulta, impresso-perforata.
- g. Ostiolum vetustum, apertura lata submarginata.
- h. Stromata horizontaliter sectum cum sphaerulis. Omnes sub. microsc. comp.

Fig. 3. *Sphaeria* rostellata.

- a. Caulis herbae cum sphaerulis, magn. nat.
- b. Sphaeria junior hemisphaerica ostiolo papilliformi.
- c. Varietas ostiolo subpyramidato.
- d. Sphaeria adulta umbonata, ostiolo hispido.
- e. Sphaeria ostiolo deciduo, umhonata. Omnes sub microsc. comp.

Fig. 4. *Monilia* hyalina.

- a. culmus Bromi cum floccis. magn. nat.
- b. c. Flocci magis aucti.
- d. — magnopere aucti.

Tab. IV.
Fig. 1. *Thelophora* evolvens. Fr.

- a. Spec. junius clausum. magn. nat.
- b. — evolutum magn. nat.

Fig. 2. *Peziza* Lecideola.

- a. magn. nat.
- b. pars sub microsc. comp.

Fig. 3. *Sphaeria* rufo-fusca.

- a. ramulus cum cespitibus pluribus. Dupplo auct.
- b. cæspes perpendiculariter sectus cum stromato. Dupl.

c. — horizontaliter sectus cum sperulis cavis
intus albis. Duppl. auct.

d. *Sphaeria solitaria*, sub microsc. comp.
maxim. aucta.

Fig. 4. *Sphaeria nigrella*.

a. Caulis urticae cum maculo et sphaeriis, magn.
nat.

b. Idem auctus.

c. *Sphaeria solitaria* dissecta, sub micr. comp.

Fig. 5. *Sphaeria excipuliformis*.

a. a. magn. nat. cum ostiolo.

b. — — ostiolo deciduo.

c. *Sphaeria* sub. micr. comp. ostiolo breviori.

Fig. 6. *Sphaeria Hederae*, a. magn. nat.

b. magn. auct.

c. *sphaeria solitaria* sub. microsc. comp.

Fig. 7. *Xylama rubiforme* i. *Lonicerae*.

a. magn. nat. in folio vivo.

b. Pars folii auct. cum Perith.

c. transversim dissecta, sub microsc. comp.

Fig. 1.



Fig. 2.



Fig. 3.




Fig. 4.



Fig. 5.




Fig. 6.






Fig. 1.



Fig. 2.



Fig. 3.




Fig. 4.



Fig. 5.



Fig. 6.



Fig. 7.




Fig. 8.






Fig. 1.





Fig. 2.




Fig. 3.





Fig. 4.





Fig. 1.



Fig. 2.



Fig. 3.




Fig. 4.




Fig. 5.



Fig. 7.



Fig. 6.

